

CERTAIN NEW CONTINUED FRACTIONS FOR THE RATIO OF TWO ${}_3\psi_3$ SERIES

Maheshwar Pathak* & Pankaj Srivastava**

*Department of Mathematics, University of Petroleum & Energy Studies, Dehradun-248007, India
**Department of Mathematics, Motilal Nehru National Institute of Technology, Allahabad-211004, India

(Received on: 18-04-12; Accepted on: 09-05-12)

ABSTRACT

In the present paper we have developed certain new continued fractions representations for the ratios of two basic bilateral hypergeometric series ${}_3\psi_3$.

Keywords: Basic hypergeometric series, Basic bilateral hypergeometric series, Continued fraction.

AMS Subject Classification: 33D15, 11A55.

1. INTRODUCTION

Now a days continued fractions have been become center of attraction for pure and applied mathematicians. The Indian mathematician Aryabhata(d. 550 AD) [4] used a continued fraction to solve a linear indeterminate equation. The development of continued fractions theory was stopped for more than a thousand years due to the specific applications. Major impacts of continued fractions in the different field of mathematics have been noticed from the beginning of 20th century. The Indian genius Ramanujan [16, 17] had shown new facet of continued fractions in the field of q -series. Ramanujan and his works on continued fractions had inspired a number of mathematicians working in the field of q -series to carry out researches in this direction. There are a number of mathematicians namely R. P. Agarwal [1, 2], G. E. Andrews and D. Bowman [3], B. C. Berndt [5], N. A. Bhagirathi [6,7], S. Bhargava, C. Adiga and D. D. Somashekara [8], R. Y. Denis [9, 10, 11], R. Y. Denis and S. N. Singh [12], Maheshwar Pathak and Pankaj Srivastava [14], P. Rai [15], S. N. Singh [18, 19, 20], Pankaj Srivastava [21], A. Verma, R. Y. Denis and K. Srinivasa Rao [22] etc. have established several results for basic hypergeometric series ${}_2\phi_1$, ${}_3\phi_2$ and basic bilateral hypergeometric series ${}_2\psi_2$, ${}_3\psi_3$ in terms of continued fractions. In the present paper certain new results for the quotients of two ${}_3\psi_3$ series in terms of continued fractions have been established following the techniques developed by Pankaj Srivastava [21] and also certain special cases have been deduced.

2. DEFINITIONS AND NOTATIONS

We shall use the following q -symbols:

For

$$\begin{aligned} |q| < 1 \text{ and } |q^r| < 1 \\ (a; q)_n &= \prod_{s=0}^{n-1} (1 - aq^s), \quad n \geq 1 \\ (a; q^r)_n &= \prod_{s=0}^{n-1} (1 - aq^{rs}), \quad n \geq 1 \\ (a; q)_0 &= 1, \quad (a; q^r)_0 = 1 \\ (a; q^r)_{\infty} &= \prod_{s=0}^{\infty} (1 - aq^{rs}) \\ (a)_n &= (a; q)_n. \end{aligned}$$

Corresponding author: Maheshwar Pathak*

*Department of Mathematics, University of Petroleum & Energy Studies, Dehradun-248007, India

A generalized basic hypergeometric series with base q is defined as:

$${}_A\phi_{A-1} \left[\begin{matrix} a_1, & a_2, & \dots, & a_A \\ b_1, & b_2, & \dots, & b_{A-1} \end{matrix}; q; z \right] = \sum_{n=0}^{\infty} \frac{(a_1; q)_n \dots (a_A; q)_n z^n}{(b_1; q)_n \dots (b_{A-1}; q)_n (q; q)_n},$$

where $|z| < 1$, $|q| < 1$.

A generalized basic bilateral hypergeometric series with base q is defined as:

$${}_A\Psi_A \left[\begin{matrix} a_1, & a_2, & \dots, & a_A \\ b_1, & b_2, & \dots, & b_A \end{matrix}; q; z \right] = \sum_{n=-\infty}^{\infty} \frac{(a_1; q)_n \dots (a_A; q)_n z^n}{(b_1; q)_n \dots (b_A; q)_n},$$

where $|b_1 b_2 \dots b_A / a_1 a_2 \dots a_A| < |z| < 1$, $|q| < 1$.

A finite continued fraction is an expression of the type:

$$\frac{a_1}{a_2 + a_4 + a_6 + a_8 + a_{10} + a_{12} + \dots} \frac{a_3}{a_n} \frac{a_5}{a_7} \frac{a_9}{a_{11}} \frac{a_{11}}{a_{n-1}}.$$

where $a_1, a_2, a_3, a_4, \dots$ are real or complex numbers.

It is an infinite continued fraction when $n \rightarrow \infty$:

In order to establish main results, we shall make use of the following known results:

S. N. Singh [20] has established the following general transformation formula between basic and basic bilateral hypergeometric series:

$$\begin{aligned} {}_{r+3}\psi_{r+3} \left[\begin{matrix} \frac{a}{b}, & cq, & dq, & \frac{b_1}{b} q^{m_1}, & \dots, & \frac{b_r}{b} q^{m_r} \\ \frac{q}{b}, & c, & d, & \frac{b_1}{b}, & \dots, & \frac{b_r}{b} \end{matrix}; q; z \right] &= \frac{\left(q, \frac{bq}{az}, \frac{az}{b}, \frac{q}{a}; q \right)_{\infty}}{\left(\frac{q}{b}, \frac{q}{az}, az, \frac{bq}{a}; q \right)_{\infty}} \\ &\times \frac{(1-bc)(1-bd)(b_1; q)_{m_1} \dots (b_r; q)_{m_r}}{(1-c)(1-d)(b_1/b; q)_{m_1} \dots (b_r/b; q)_{m_r}} \times {}_{r+3}\phi_{r+2} \left[\begin{matrix} a, & bcq, & bdq, & b_1 q^{m_1}, & \dots, & b_r q^{m_r} \\ bc, & bd, & b_1, & \dots, & b_r \end{matrix}; q; z \right] \quad (1) \end{aligned}$$

Bhargava, Adiga and Somashekara [8] have established the following results for the quotients of two ${}_3\phi_2$ series, which are as follows:

$$\begin{aligned} \frac{{}_3\phi_2 \left[\begin{matrix} a, & b, & cq \\ dq, & e \end{matrix}; q; de/abc \right]}{{}_3\phi_2 \left[\begin{matrix} a, & b, & cq \\ d, & e \end{matrix}; q; de/abc \right]} &= \frac{(1-d)}{(1-d) + (1-dq)(1-e/cq) +} \\ &\frac{F_0}{(1-dq^2) +} \cdots \frac{E_n}{(1-dq^{2n+1})(1-e/cq) +} \frac{F_n}{(1-dq^{2n+2}) +} \cdots \quad (2) \end{aligned}$$

where

$$E_n = (deq^n/abc)(1-aq^n)(1-bq^n)(dq^n - c), \quad n = 0, 1, 2, \dots$$

and

$$F_n = (e/cq)[1 - (dq^{n+1}/a)][1 - (dq^{n+1}/b)](1-cq^{n+1}), \quad n = 0, 1, 2, \dots$$

$$\frac{{}_3\phi_2 \left[\begin{matrix} a, & b, & c \\ dq, & e & \end{matrix}; q; \frac{deq}{abc} \right]}{{}_3\phi_2 \left[\begin{matrix} a, & b, & c \\ d, & e & \end{matrix}; q; \frac{de}{abc} \right]} = \frac{(1-d)[1-(de/abc)]}{(1-d)[1-(de/abc)] + (1-dq) +} \frac{D_0}{(1-dq) +}$$

$$\frac{C_0}{(1-dq^2)[1-(de/abc)] +} \cdots \frac{D_n}{(1-dq^{2n+1}) +} \frac{C_n}{(1-dq^{2n+2})[1-(de/abc)] +} \cdots \quad (3)$$

where

$$C_n = -eq^n[1-(dq^{n+1}/a)][1-(dq^{n+1}/b)][1-(dq^{n+1}/c)], \quad n=0,1,2,\dots$$

and

$$D_n = (de/abc)(1-aq^n)(1-bq^n)(1-cq^n), \quad n=0,1,2,\dots$$

$$\frac{{}_3\phi_2 \left[\begin{matrix} aq, & bq, & c \\ d, & eq & \end{matrix}; q; \frac{de}{abcq} \right]}{{}_3\phi_2 \left[\begin{matrix} a, & b, & c \\ d, & e & \end{matrix}; q; \frac{de}{abc} \right]} = \frac{(1-e)/[1-(de/abcq)]}{1 +} \frac{c[(de/abcq)-(e/c)][1-(d/c)]}{(1-d)[1-(de/abcq)] +} \frac{(de/abcq)(1-aq)(1-bq)(1-c)}{(1-aq) +}$$

$$\frac{B_1}{[1-(de/abcq)] + (1-aq) +} \frac{A_1}{...} \frac{B_n}{[1-(de/abcq)] + (1-aq) +} \frac{A_n}{(1-aq) +} \cdots \quad (4)$$

where

$$A_n = (de/abcq)(1-bq^{n+1})(1-cq^n), \quad n=1,2,3,\dots$$

and

$$B_n = aq[1-(dq^{n-1}/a)][1-(eq^{n-1}/a)], \quad n=1,2,3,\dots$$

The identity due to Gasper and Rahman [13, III.9, pp. 241] is:

$${}_3\phi_2 \left[\begin{matrix} a, & b, & c \\ d, & e & \end{matrix}; q; \frac{de}{abc} \right] = \frac{(e/a, de/bc; q)_\infty}{(e, de/abc; q)_\infty} {}_3\phi_2 \left[\begin{matrix} a, & d/b, & d/c \\ d, & de/bc & \end{matrix}; q; \frac{e/a}{d} \right] \quad (5)$$

3. MAIN RESULTS

We have established the following results which are as follows:

$$\frac{{}_3\psi_3 \left[\begin{matrix} a/b, & b_1q^{m_1}/b, & b_2q^{m_2+1}/b \\ q/b, & b_1/b, & b_2/b & \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]}{{}_3\psi_3 \left[\begin{matrix} a/b, & b_1q^{m_1}/b, & b_2q^{m_2}/b \\ q/b, & b_1/b, & b_2/b & \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]} = \frac{(b_2q;q)_{m_2}(b_2/b;q)_{m_2}}{(b_2q/b;q)_{m_2}(b_2;q)_{m_2}} \times \left[\frac{(1-b_2)}{(1-b_2) +} \right.$$

$$\left. \frac{E_0}{(1-b_2q)(1-b_1/b_2q^{m_2+1}) + (1-b_2q^2) +} \frac{F_0}{(1-b_2q^{2n+1})(1-b_1/b_2q^{m_2+1}) + (1-b_2q^{2n+2}) +} \frac{E_n}{(1-b_2q^{2n+1})(1-b_1/b_2q^{m_2+1}) + (1-b_2q^{2n+2}) +} \frac{F_n}{(1-b_2q^{2n+2}) +} \cdots \right] \quad (6)$$

where

$$E_n = (q^n/aq^{m_1+m_2})(1-aq^n)(1-b_1q^{m_1+n})(b_2q^n-b_2q^{m_2}), \quad n=0,1,2,\dots$$

and

$$F_n = (b_1/b_2q^{m_2+1})(1-b_2q^{n+1}/a)(1-b_2q^{n+1}/b_1q^{m_1})(1-b_2q^{m_2+n+1}), \quad n=0,1,2,\dots$$

$$\frac{{}_3\psi_3 \left[\begin{matrix} a/b, b_1 q^{m_1+1}/b, b_2 q^{m_2+1}/b \\ q/b, b_1 q/b, b_2 q/b \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]}{{}_3\psi_3 \left[\begin{matrix} a/b, b_1 q^{m_1}/b, b_2 q^{m_2}/b \\ q/b, b_1/b, b_2/b \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]} = \frac{(b-b_1)(b-b_2)(1-b_1 q^{m_1})}{(1-b_1)(1-b_2)(b-b_1 q^{m_1})} \times \frac{(1-b_2 q^{m_2})a(1-b_2)}{(b-b_2 q^{m_2})(a-b_2)}$$

$$\left[\frac{(1-b_1)(a-b_2)}{(1-b_1)(a-b_2)+(1-b_1 q)+(1-b_1 q^2)(1-b_2/a)+(1-b_1 q^{2n+1})+(1-b_1 q^{2n+2})(1-b_2/a)+\dots} \frac{D_0}{C_0} \dots \frac{D_n}{C_n} \dots \right] \quad (7)$$

where

$$C_n = -q^{n-(m_1+m_2)}(1-b_1 q^{n+1}/a)(1-b_1 q^{n+m_1+1})(1-b_2 q^{n+m_2+1}), \quad n=0,1,2,\dots$$

and

$$D_n = (b_2/a)(1-aq^n)(1-q^{n-m_1})(1-b_1 q^{n-m_2}/b_2), \quad n=0,1,2,\dots$$

$$\frac{{}_3\psi_3 \left[\begin{matrix} aq/b, b_1 q^{m_1+1}/b, b_2 q^{m_2}/b \\ q/b, b_1 q/b, b_2/b \end{matrix}; q; \frac{1}{aq^{m_1+m_2+1}} \right]}{{}_3\psi_3 \left[\begin{matrix} a/b, b_1 q^{m_1}/b, b_2 q^{m_2}/b \\ q/b, b_1/b, b_2/b \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]} = \frac{(a-1)(1-b_1 q^{m_1})(b-b_1)}{(a-b)(b-b_1 q^{m_1})(1-b_1)}$$

$$\left[\frac{(1-b_1)(1-1/aq^{m_1+m_2+1})}{1+} \frac{b_2 q^{m_2}(1/aq^{m_1+m_2+1}-b_1 q^{-m_2}/b_2)(1-q^{-m_2})}{(1-b_2)(1-1/aq^{m_1+m_2+1})+} \frac{(1/aq^{m_1+m_2+1})(1-aq)(1-b_1 q^{m_1+1})(1-b_2 q^{m_2})}{(1-aq)+} \right.$$

$$\left. \frac{B_1}{(1-1/aq^{m_1+m_2+1})+} \frac{A_1}{(1-aq)+} \dots \frac{B_n}{(1-1/aq^{m_1+m_2+1})+} \frac{A_n}{(1-aq)+} \dots \right], \quad m_2 \neq 0 \quad (8)$$

where

$$A_n = (1/aq^{m_1+m_2+1})(1-b_1 q^{m_1+n+1})(1-b_2 q^{m_2+n}), \quad n=1,2,3,\dots$$

and

$$B_n = aq(1-b_2 q^{n-1}/a)(1-b_1 q^{n-1}/a), \quad n=1,2,3,\dots$$

4. PROOF OF MAIN RESULTS

The proof of main results 6, 7 and 8 are given below:

Proof of (6): Taking $r = 2, c = d = 0$ in transformation formula (1), we get

$$\begin{aligned} {}_3\psi_3 \left[\begin{matrix} a/b, b_1 q^{m_1}/b, b_2 q^{m_2}/b \\ q/b, b_1/b, b_2/b \end{matrix}; q; z \right] \\ = \frac{\left(q, \frac{bq}{az}, \frac{az}{b}, \frac{q}{a}; q \right)_\infty}{\left(\frac{q}{b}, \frac{q}{az}, \frac{bq}{a}; q \right)_\infty} \times \frac{(b_1; q)_{m_1} (b_2; q)_{m_2}}{(b_1/b; q)_{m_1} (b_2/b; q)_{m_2}} \times {}_3\phi_2 \left[\begin{matrix} a, b_1 q^{m_1}, b_2 q^{m_2} \\ b_1, b_2 \end{matrix}; q; z \right] \end{aligned} \quad (9)$$

replacing b_2 by $b_2 q$ and putting $z = 1/aq^{m_1+m_2}$ in (9), further taking $z = 1/aq^{m_1+m_2}$ in (9) and taking ratio of these two and making use of the result (2) and after simplification, we obtain the result (6).

Proof of (7): Taking $z = 1/aq^{m_1+m_2}$ in (9), we get

$$\begin{aligned} {}_3\psi_3 \left[\begin{matrix} a/b, & b_1 q^{m_1} / b, & b_2 q^{m_2} / b \\ q/b, & b_1 / b, & b_2 / b \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right] &= \frac{\left(q, bq^{m_1+m_2+1}, 1/bq^{m_1+m_2}, q/a; q \right)_\infty}{\left(q/b, q^{m_1+m_2+1}, 1/q^{m_1+m_2}, bq/a; q \right)_\infty} \\ &\times \frac{(b_1; q)_{m_1} (b_2; q)_{m_2}}{(b_1 / b; q)_{m_1} (b_2 / b; q)_{m_2}} \times {}_3\phi_2 \left[\begin{matrix} a, & b_1 q^{m_1}, & b_2 q^{m_2} \\ b_1, & b_2 \end{matrix}; q; \frac{1}{a q^{m_1+m_2}} \right] \end{aligned} \quad (10)$$

making use of result (5) in (10) and then replace b_1, b_2 by $b_1 q, b_2 q$ in that result. Again we use the result (5) in (10) and take the ratio of these two results, finally making use of the result (3) and after simplification; we get the result (7).

Proof of (8): Replacing a by aq and b_1 by $b_1 q$ in (10), we get

$$\begin{aligned} {}_3\psi_3 \left[\begin{matrix} aq/b, & b_1 q^{m_1+1} / b, & b_2 q^{m_2} / b \\ q/b, & b_1 q / b, & b_2 / b \end{matrix}; q; \frac{1}{aq^{m_1+m_2+1}} \right] &= \frac{\left(q, bq^{m_1+m_2+1}, 1/bq^{m_1+m_2}, 1/a; q \right)_\infty}{\left(q/b, q^{m_1+m_2+1}, 1/q^{m_1+m_2}, b/a; q \right)_\infty} \\ &\times \frac{(b_1 q; q)_{m_1} (b_2; q)_{m_2}}{(b_1 q / b; q)_{m_1} (b_2 / b; q)_{m_2}} \times {}_3\phi_2 \left[\begin{matrix} aq, & b_1 q^{m_1+1}, & b_2 q^{m_2} \\ b_1 q, & b_2 \end{matrix}; q; \frac{1}{a q^{m_1+m_2+1}} \right] \end{aligned} \quad (11)$$

taking the ratio of (11) and (10) and making use of result (4) and after simplification, we get the result (8).

5. SPECIAL CASES:

In this section, we shall consider certain applications of the main results obtained in the §3.

(1) Putting $m_1 = 0$ in result (6), we get

$$\begin{aligned} \frac{{}_2\psi_2 \left[\begin{matrix} a/b, & b_2 q^{m_2+1} / b \\ q/b, & b_2 / b \end{matrix}; q; \frac{1}{aq^{m_2}} \right]}{{}_2\psi_2 \left[\begin{matrix} a/b, & b_2 q^{m_2} / b \\ q/b, & b_2 / b \end{matrix}; q; \frac{1}{aq^{m_2}} \right]} &= \frac{(b_2 q; q)_{m_2} (b_2 / b; q)_{m_2}}{(b_2 q / b; q)_{m_2} (b_2; q)_{m_2}} \times \left[\frac{(1-b_2)}{(1-b_2)+} \frac{E_o}{(1-b_2 q)(1-b_1/b_2 q^{m_2+1})} + \right. \\ &\left. \frac{F_0}{(1-b_2 q^2)+} \cdots \frac{E_n}{(1-b_2 q^{2n+1})(1-b_1/b_2 q^{m_2+1})+} \frac{F_n}{(1-b_2 q^{2n+2})+} \cdots \right] \end{aligned} \quad (12)$$

where

$$E_n = (q^n / aq^{m_2}) (1 - aq^n) (1 - b_1 q^n) (b_2 q^n - b_2 q^{m_2}), \quad n = 0, 1, 2, \dots$$

and

$$F_n = (b_1 / b_2 q^{m_2+1}) (1 - b_2 q^{n+1} / a) (1 - b_2 q^{n+1} / b_1) (1 - b_2 q^{m_2+n+1}), \quad n = 0, 1, 2, \dots$$

(2) Putting $m_2 = 0$ in result (11) and making use of the Ramanujan ${}_1\psi_1$ summation formula [13, (5.2.1), pp. 126], we get

$$\begin{aligned}
 & {}_2\psi_2 \left[\begin{matrix} a/b, & b_2q/b \\ q/b, & b_2/b \end{matrix}; q; \frac{1}{a} \right] \\
 &= \frac{\left(q, q/a, 1/b, qb; q \right)_\infty}{\left(q/b, qb/a, 1/a, b^2; q \right)_\infty} \times \left[\frac{(1-b_2)}{(1-b_2) +} \frac{G_0}{(1-b_2q)(1-b_1/b_2q) +} \right. \\
 &\quad \left. \frac{H_o}{(1-b_2q^2) +} \dots \frac{G_n}{(1-b_2q^{2n+1})(1-b_1/b_2q) +} \frac{H_n}{(1-b_2q^{2n+2}) +} \dots \right] \tag{14}
 \end{aligned}$$

where

$$G_n = (q^n/a)(1-aq^n)(1-b_1q^n)(b_2q^n - b_2), \quad n=0,1,2,\dots$$

and

$$H_n = (b_1/b_2q)(1-b_2q^{n+1}/a)(1-b_2q^{n+1}/b)(1-b_2q^{n+1}), \quad n=0,1,2,\dots$$

(3) Putting $b_1 = a$ and $b = bq$ in result (7), we get

$$\begin{aligned}
 & {}_2\psi_2 \left[\begin{matrix} aq^{m_1}/b, & b_2q^{m_2}/b \\ 1/b, & b_2/b \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right] = \frac{(bq-a)(bq-b_2)(1-aq^{m_1})}{(1-a)(1-b_2)(bq-aq^{m_1})} \times \frac{(1-b_2q^{m_2})a(1-b_2)}{(b-qb_2q^{m_2})(a-b_2)} \\
 & {}_2\psi_2 \left[\begin{matrix} aq^{m_1-1}/b, & b_2q^{m_2-1}/b \\ 1/b, & b_2/bq \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right] = \frac{(1-a)(a-b_2)}{(1-a)(a-b_2) +} \frac{D_0}{(1-aq) +} \frac{C_0}{(1-aq^2)(1-b_2/a) +} \dots \frac{D_n}{(1-aq^{2n+1}) +} \frac{C_n}{(1-aq^{2n+2})(1-b_2/a) +} \dots \tag{14}
 \end{aligned}$$

where

$$C_n = -q^{n-(m_1+m_2)}(1-q^{n+1})(1-aq^{n+m_1+1})(1-b_2q^{n+m_2+1}), \quad n=0,1,2,\dots$$

and

$$D_n = (b_2/a)(1-aq^n)(1-q^{n-m_1})(1-aq^{n-m_2}/b), \quad n=0,1,2,\dots$$

(4) Putting $b = 1$ in result (6), we get

$$\begin{aligned}
 & {}_3\phi_2 \left[\begin{matrix} a, & b_1q^{m_1}, & b_2q^{m_2+1} \\ b_1, & b_2 & \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right] = \left[\frac{(1-b_2)}{(1-b_2) +} \frac{E_o}{(1-b_2q)(1-b_1/b_2q^{m_2+1}) +} \right. \\
 & {}_3\phi_2 \left[\begin{matrix} a, & b_1q^{m_1}, & b_2q^{m_2} \\ b_1, & b_2 & \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right] \left. \right]
 \end{aligned}$$

$$\left[\frac{F_0}{(1-b_2q^2)+} \cdots \frac{E_n}{(1-b_2q^{2n+1})(1-b_1/b_2q^{m_2+1})+} \frac{F_n}{(1-b_2q^{2n+2})+} \cdots \right] \quad (15)$$

where

$$E_n = (q^n/aq^{m_1+m_2})(1-aq^n)(1-b_1q^{m_1+n})(b_2q^n-b_2q^{m_2}), \quad n=0,1,2,\dots$$

and

$$F_n = (b_1/b_2q^{m_2+1})(1-b_2q^{n+1}/a)(1-b_2q^{n+1}/b_1q^{m_1})(1-b_2q^{m_2+n+1}), \quad n=0,1,2,\dots$$

(5) Putting $a = \alpha q$ and $b = 1$ in result (7), we get

$$\begin{aligned} {}_3\phi_2 \left[\begin{matrix} \alpha q, & b_1q^{m_1+1}, & b_2q^{m_2+1} \\ b_1q, & b_2q & \end{matrix}; q; \frac{1}{\alpha q^{m_1+m_2+1}} \right] &= \frac{\alpha q(1-b_2)}{(\alpha q-b_2)} \left[\frac{(1-b_1)(\alpha q-b_2)}{(1-b_1)(\alpha q-b_2)+} \right. \\ &\quad {}_3\phi_2 \left[\begin{matrix} \alpha q, & b_1q^{m_1}, & b_2q^{m_2} \\ b_1, & b_2 & \end{matrix}; q; \frac{1}{\alpha q^{m_1+m_2+1}} \right] \\ &\quad \left. \frac{D_o}{(1-b_1q)+} \frac{C_0}{(1-b_1q^2)(1-b_2/\alpha q)+} \cdots \frac{D_n}{(1-b_1q^{2n+1})+} \frac{C_n}{(1-b_1q^{2n+2})(1-b_2/\alpha q)+} \cdots \right] \quad (16) \end{aligned}$$

where $|\alpha| > 1$

$$C_n = -q^{n-(m_1+m_2)}(1-b_1q^n/\alpha)(1-b_1q^{n+m_1+1})(1-b_2q^{n+m_2+1}), \quad n=0,1,2,\dots$$

and

$$D_n = (b_2/\alpha q)(1-\alpha q^{n+1})(1-q^{n-m_1})(1-b_1q^{n-m_2}/b_2), \quad n=0,1,2,\dots$$

(6) Putting $b_1 = 0$ in result (8), we get

$$\begin{aligned} {}_2\psi_2 \left[\begin{matrix} aq/b, & b_2q^{m_2}/b \\ q/b, & b_2/b & \end{matrix}; q; \frac{1}{aq^{m_1+m_2+1}} \right] &= \frac{(a-1)}{(a-b)} \times \left[\frac{1/(1-1/aq^{m_1+m_2+1})}{1+} \frac{(b_2q^{-(m_1+1)}/a)(1-q^{-m_2})}{(1-b_2)(1-1/aq^{m_1+m_2+1})+} \right. \\ &\quad \left. \frac{(1-1/aq^{m_1+m_2+1})(1-aq)(1-b_2q^{m_2})}{(1-aq)+} \frac{B_1}{(1-1/aq^{m_1+m_2+1})+} \frac{A_1}{(1-aq)+} \cdots \frac{B_n}{(1-1/aq^{m_1+m_2+1})+} \frac{A_n}{(1-aq)+} \cdots \right], \quad m_2 \neq 0 \quad (17) \end{aligned}$$

where

$$A_n = (1/aq^{m_1+m_2+1})(1-b_2q^{m_2+n}), \quad n=1,2,3,\dots$$

and

$$B_n = aq(1-b_2q^{n-1}/a), \quad n=1,2,3,\dots$$

(7) Putting $b = 1$ and $b_1 = 0$ in result (8), we get

$$\frac{{}_2\phi_1 \left[\begin{matrix} aq, & b_2 q^{m_2} \\ b_2 & \end{matrix}; q; \frac{1}{aq^{m_1+m_2+1}} \right]}{{}_2\phi_1 \left[\begin{matrix} a, & b_2 q^{m_2} \\ b_2 & \end{matrix}; q; \frac{1}{aq^{m_1+m_2}} \right]} = \left[\frac{1/(1-1/aq^{m_1+m_2+1})}{1+} \frac{b_2 q^{-(m_1+1)} / a(1-q^{-m_2})}{(1-b_2)(1-1/aq^{m_1+m_2+1})+} \frac{(1/aq^{m_1+m_2+1})(1-aq)(1-b_2 q^{m_2})}{(1-aq)+} \right. \\ \left. \frac{B_1}{(1-1/aq^{m_1+m_2+1})+} \frac{A_1}{(1-aq)+} \dots \frac{B_n}{(1-1/aq^{m_1+m_2+1})+} \frac{A_n}{(1-aq)+} \dots \right], \quad m_2 \neq 0 \quad (18)$$

where

$$A_n = (1/aq^{m_1+m_2+1})(1-b_2 q^{m_2+n}), \quad n = 1, 2, 3, \dots$$

and

$$B_n = aq(1-b_2 q^{n-1}/a), \quad n = 1, 2, 3, \dots$$

REFERENCES

- [1] Agarwal, R. P., "Pade approximants continued fractions and Heine's q -series", J. Math. Phys. Sci. 3 26 (1992), 281-290.
- [2] Agarwal, R. P., "Resonance of Ramanujan's mathematics Vol.III", New Age International Pvt. Ltd. Publishers, New Delhi, 1998.
- [3] Andrews, G.E. and Bowman, D., "A full extension of the Rogers-Ramanujan continued fraction", Proc. Amer. Math. Soc. (11) 123 (1995), 3343-3350.
- [4] Brezinski, Claude, "History of continued fractions and Pade approximants", Springer-Verlag, New York, 1980.
- [5] Berndt, B. C., "Ramanujan's Notebook part III", Springer-Verlag, 1991.
- [6] Bhagirathi, N. A., "On certain investigations in q -series and continued fractions", Math. Student 1-4 56 (1988), 158-170.
- [7] Bhagirathi, N. A., "On basic bilateral hypergeometric series and continued fractions", Math. Student 1-4 56 (1988), 135-141.
- [8] Bhargava, S., Adiga, C., Somashekara, D. D., "On certain continued fractions related to ${}_3\phi_2$ basic hypergeometric functions", J. Math. Phys. Sci. 21 (1987), 613-629.
- [9] Denis, R. Y., "On basic hypergeometric function and continued fractions", Math. Student 1-4 52 (1984), 129-136.
- [10] Denis, R. Y., "On certain q -series and continued fractions", Math. Student 44 (1983), 70-76.
- [11] Denis, R. Y., "On generalization of continued fraction of Gauss", Internat. J. Math. & Math. Sci. (4) 13 (1990), 741-746.
- [12] Denis, R. Y. and Singh, S. N., "Certain result involving q -series and continued fractions", Far East J. Math. Sci. (FJMS) (5) 3 (2001), 731-737.
- [13] Gasper, G. and Rahman, M., "Basic Hypergeometric Series", Cambridge University Press, New York, 1991.
- [14] Pathak, Maheshwar and Srivastava, Pankaj, "A note on continued fractions and ${}_3\psi_3$ series", Ital. J. Pure Appl. Math. 27 (2010), 191-200.
- [15] Rai, P., "Certain bilateral extensions of the Rogers-Ramanujan continued fraction", Ganita (1) 57 (2006), 73-87.
- [16] Ramanujan, S., "Note book of Ramanujan II", T.I.F.R. Bombay, 1957.

- [17] Ramanujan, S., “*The lost note book and other unpublished papers*”, Narosa, New Delhi 1988.
- [18] Singh, S. N., “*On q-hypergeometric function and continued fractions*”, Math. Student 1-4 56 (1988), 81-84.
- [19] Singh, S. N., “*Basic hypergeometric series and continued fractions*”, Math. Student 1-4 56 (1988), 91-96.
- [20] Singh, S. N., “*Certain transformation formulae for basic and bibasic hypergeometric series*”, Proc. Nat. Acad. Sci. India (III) 65(A) (1995), 319-329.
- [21] Srivastava, Pankaj, “*Certain continued fractions for quotients of two ${}_3\psi_3$ series*”, Proc. Nat. Acad. Sci. India (IV) 78(A) (2008), 327-330.
- [22] Verma, A., Denis, R. Y. and Srinivasa Rao, K., “*New continued fractions involving basic hypergeometric ${}_3\phi_2$ function*”, Jour. Math. Phy. Sci. 6 21 (1987), 585-592.

Source of support: Nil, Conflict of interest: None Declared