

## NEW FAMILIES OF 3-TOTAL PRODUCT CORDIAL GRAPHS

R. Ponraj<sup>1\*</sup>, M. Sivakumar<sup>2</sup> and M. Sundaram<sup>3</sup>

<sup>1</sup>Department of Mathematics, Sri Paramakalyani College, Alwarkurchi-627412, India

<sup>2</sup>Department of Mathematics, Unnamalai Institute of Technology, Kovilpatti-628502, India

<sup>3</sup>Associate Professor (Rtd.), Department of Mathematics, Sri Paramakalyani College  
Alwarkurchi-627412, India

(Received on: 14-05-12; Accepted on: 31-05-12)

### ABSTRACT

Let  $f$  be a map from  $V(G)$  to  $\{0, 1, \dots, k-1\}$  where  $k$  is an integer,  $2 \leq k \leq |V(G)|$ . For each edge  $uv$  assign the label  $f(u)f(v)(\text{mod } k)$ .  $f$  is called a  $k$ - Total Product cordial labeling if  $|f(i) - f(j)| \leq 1, i, j \in \{0, 1, \dots, k-1\}$ , where  $f(x)$  denotes the total number of vertices and edges labelled with  $x (x=0, 1, 2, \dots, k-1)$ . A graph that admits a  $k$ - Total Product cordial labelling is called a  $k$ - Total Product cordial graph. In this paper we investigate 3- Total Product cordial labeling behaviour of some standard graphs like Wheels, Helms, Dragons, etc.

**Keywords:** Wheel, Helms, Dragon,  $C_n \odot 2K_1$ .

**Mathematics Subject Classification (2000):** 05C78.

### 1. INTRODUCTION

The graphs considered here are finite, undirected and simple. The vertex set and edge set of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$  respectively. The following definitions are used here.

- The corona  $G_1 \odot G_2$  of two graphs  $G_1$  and  $G_2$  is defined as the graph  $G$  obtained by taking one copy of  $G_1$  (which has  $p_1$  vertices) and  $p_1$  copies of  $G_2$  and then joining the  $i^{\text{th}}$  vertex of  $G_1$  to every vertex in the  $i^{\text{th}}$  copy  $G_2$ .
- The graph  $W_n = C_n + K_1$  is called a wheel.
- The Helms  $H_n$  is the graph obtained from wheel by attaching a pendant edge at each vertex of the cycle  $C_n$ .
- A Dragon is formed by identifying the end vertex of the path to the vertex of a cycle.
- $m$  copies of the graph  $G$  is denoted by  $mG$ .

The notion of  $k$ -Product cordial labeling of graph was introduced in [2] where the  $k$ -Product cordial labeling behaviour of some standard graphs was studied. Also  $k$ -Total Product labeling of graphs was introduced in [4]. Obviously 2-Total Product cordial labeling is simply a Total Product cordial labeling [5]. Also 3-Total Product cordial labeling behaviour of some standard graphs was studied in [3]. In this paper we investigate 3-Total Product cordial labeling behaviour of Helms, Wheel, Dragon,  $C_n \odot 2K_1$  and some standard graphs. Terms not defined here are used in the sense of Harary [1].

### 2. k-TOTAL PRODUCT CORDIAL LABELING

#### Definition 2.1:

Let  $f$  be a function from  $V(G)$  to  $\{0, 1, \dots, k-1\}$  where  $k$  is an integer,  $2 \leq k \leq |V(G)|$ . For each edge  $uv$ , assign the label  $f(u)f(v)(\text{mod } k)$ .  $f$  is called a  $k$ - Total Product cordial labeling of  $G$  if  $|f(i) - f(j)| \leq 1, i, j \in \{0, 1, \dots, k-1\}$  where  $f(x)$  denotes the total number of vertices and edges labelled with  $x (x=0, 1, 2, \dots, k-1)$ . A graph with a  $k$ - Total Product cordial labelling is called a  $k$ - Total Product cordial graph.

**Theorem 2.2:** Let  $G$  be a  $(p, p)$  graph. Then  $mG$  is 3-Total Product cordial where  $m \equiv 0 (\text{mod } 3)$ .

*Corresponding author: R. Ponraj<sup>1\*</sup>*

<sup>1</sup>Department of Mathematics, Sri Paramakalyani College, Alwarkurchi-627412, India

**Proof:** let  $m=3t$ . Clearly  $mG$  has  $3pt$  vertices and  $3pt$  edges. Assign the label 2 to all the vertices of first  $2t$  copies of  $G$ . Then assign 0 to all the vertices of remaining  $t$  copies of  $G$ .

Then  $f(0) = f(1) = f(2) = 2pt$ . Therefore  $f$  is a 3-Total Product cordial labeling.

**Corollary 2.3:** If  $m \equiv 0 \pmod{3}$ , then  $mC_n$  is 3- Total Product cordial.

**Notation:** Let  $G$  be any graph. Then the graph obtained from  $G$  by identifying the central vertex of  $K_{1,p}$  to any vertex of  $G$  is denoted by  $G * K_{1,p}$ .

**Theorem 2.4:** If  $G$  is a  $(p, p)$  graph.

(i) If  $p$  is even then  $G * K_{1, \frac{p}{2}}$  is 3-Total Product cordial.

(ii) If  $p$  is odd then  $G * K_{1, \frac{p-1}{2}}$  and  $G * K_{1, \frac{p+1}{2}}$  are 3-Total Product cordial .

**Proof: Case (i):**  $p$  is even.

Assign the label 2 to all the vertices of  $G$  and 0 to all pendant vertices of  $K_{1, \frac{p}{2}}$ . Clearly  $f(0) = f(1) = f(2) = p$ .

**Case (ii):**  $p$  is odd.

Assign label as in case (i) Clearly  $f(0) = p - 1$ , Therefore  $f(1) = f(2) = p$  for the graph  $G * K_{1, \frac{p-1}{2}}$ . and  $f(0) = p + 1$ ,

$f(1) = f(2) = p$  for the graph  $G * K_{1, \frac{p+1}{2}}$ . Therefore  $f$  is a 3-Total Product cordial labeling.

**Theorem 2.5:** The Wheel  $W_n$  is 3-Total Product Cordial.

**Proof:** Let  $C_n$  be the cycle  $u_1u_2 \dots u_nu_1$  and let  $V(W_n) = V(C_n) \cup \{u\}$   $E(W_n) = E(C_n) \cup \{uu_i : 1 \leq i \leq n\}$ .

Define  $f(u) = 0$ ,

$f(u_i) = 2, 1 \leq i \leq n$ .

Here  $f(0) = n + 1$  and  $f(1) = f(2) = n$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Theorem 2.6:** The Helms  $H_n$  is 3-Total Product cordial.

**Proof:** Let the vertex set and edge set of the wheel  $W_n$  be defined as in theorem 2.5.

Let  $V(H_n) = V(W_n) \cup \{v_i, 1 \leq i \leq n\}$  and  $E(H_n) = E(W_n) \cup \{u_i v_i : 1 \leq i \leq n\}$ .

**Case (i):**  $n \equiv 0 \pmod{3}$

Let  $n=3t$

Define  $f(u) = 0$ ,

$f(u_i) = 2, 1 \leq i \leq n$

$f(v_i) = 2, 1 \leq i \leq 2t$

$f(v_{2t+i}) = 0, 1 \leq i \leq t$

Here  $f(0) = 5t + 1$  and  $f(1) = f(2) = 5t$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (ii):**  $n \equiv 1 \pmod{3}$

Let  $n=3t+1$ . Assign labels to the vertices  $u, u_i, v_i$  ( $1 \leq i \leq n-1$ ) as in case(i). Then assign the labels 2, 2 to the vertices  $u_n, v_n$  respectively. Here  $f(0) = f(1) = f(2) = 5t + 2$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (iii):**  $n \equiv 2 \pmod{3}$

Let  $n=3t+2$ . Assign labels to the vertices  $u, u_i, v_i$  ( $1 \leq i \leq n-1$ ) as in case (ii). Then assign the labels 2, 2 to the vertices  $u_n, v_n$  respectively. Here  $f(0) = 5t + 3$  and  $f(1) = f(2) = 5t + 4$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Illustration 2.7:** A 3-Total Product cordial labeling of  $H_7$  is




Figure (i)

**Notation 2.8:** Let  $C_n$  be the cycle  $u_1, u_2, \dots, u_n, u_1$ . Let  $G_n$  denotes the graph with  $V(G_n) = V(C_n) \cup \{v_i, w_i, 1 \leq i \leq n\}$  and  $E(G_n) = E(C_n) \cup \{u_i v_i, u_i w_i, v_i w_i : 1 \leq i \leq n\}$

**Theorem 2.9:**  $G_n$  is 3-Total Product cordial.

**Proof:** Let the vertex set and edge set of the graph  $G_n$  be as defined above.

**Case (i):**  $n \equiv 0 \pmod{3}$

Let  $n=3t$ . Define  $f(v_i) = f(c_i) = 0, 1 \leq i \leq t$

$$f(w_i) = 0, 1 \leq i \leq t-1$$

$$f(w_t) = 0.$$

$$f(u_{t+i}) = f(v_{t+i}) = f(w_{t+i}) = 2, \quad 1 \leq i \leq t$$

$$f(u_{2t+i}) = 2, \quad 1 \leq i \leq t$$

$$f(v_{2t+i}) = 1, \quad 1 \leq i \leq t$$

$$f(w_{2t+i}) = 2, \quad 1 \leq i \leq t$$

Then  $f(0) = f(1) = f(2) = 7t$ . Hence  $f$  is a 3-Total Product cordial labeling.


**Case (ii):**  $n \equiv 1 \pmod{3}$

Let  $n=3t+1$ . Assign labels to the vertices  $u_i, v_i, w_i, 1 \leq i \leq n-1$  as in case (i). Then assign the labels 2, 2 and 0 to the vertices  $u_n, v_n, w_n$  respectively. Here  $f(0) = 7t + 3$  and  $f(1) = f(2) = 7t + 2$ . Hence  $f$  is 3-Total Product cordial labeling.

**Case (iii):**  $n \equiv 2 \pmod{3}$

Let  $n=3t+2$ . Assign labels to the vertices  $u_i, v_i, w_i, 1 \leq i \leq n-2$  as in case (i). Then assign the labels 0, 2, 0, 2, 2 and 2 to the vertices  $v_{n-1}, v_n, w_{n-1}, w_n, u_{n-1}, u_n$  respectively. Here  $f(0) = 7t + 5$  and  $f(1) = f(2) = 7t + 4$ . Hence  $f$  is 3-Total Product cordial labeling.

**Illustration 2.10:** A 3-Total Product cordial labelling of  $G_6$  is



**Figure (ii)**

**Theorem 2.11:**  $C_n \Theta 2K_1$  is 3-Total Product cordial.

**Proof:** Let  $V(C_n \Theta 2K_1) = \{u_i, v_i, w_i, 1 \leq i \leq n, \}$  and

$E(C_n \Theta 2K_1) = \{u_i u_{i+1}, u_n u_1 : 1 \leq i \leq n-1\} \cup \{u_i v_i, u_i w_i, 1 \leq i \leq n\}$

**Case (i):**  $n$  is even.

Define

$$f(u_i) = 2, 1 \leq i \leq n$$

$$f(v_i) = f(w_i) = 0, 1 \leq i \leq \frac{n}{2}$$

$$f(v_{\frac{n}{2}+i}) = f(w_{\frac{n}{2}+i}) = 2, 1 \leq i \leq \frac{n}{2}$$

Then  $f(0) = f(1) = f(2) = 2n$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (ii):**  $n$  is odd.

Define  $f(u_i) = 2, 1 \leq i \leq n, f(v_i) = 0, 1 \leq i \leq \frac{n+1}{2}$

$$f(w_i) = 0, 1 \leq i \leq \frac{n-1}{2}, f(v_{\frac{n+1}{2}+i}) = 2, 1 \leq i \leq \frac{n-1}{2}$$

$$f(w_{\frac{n-1}{2}+i}) = 2, 1 \leq i \leq \frac{n+1}{2}$$

Then  $f(0) = f(1) = f(2) = 2n$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Illustration 2.12:** A 3-Total Product cordial labeling of  $C_7 \odot 2K_1$  is




Figure (iii)

**Theorem 2.13:** The Dragon  $C_m @ P_n$  is 3-Total Product cordial.

**Proof:** Let  $C_m$  be the cycle  $u_1, u_2, \dots, u_m, u_1$  and  $P_n$  be the path  $v_1, v_2, \dots, v_n$ . identify the vertex  $u_1$  with  $v_1$ .

**Case (i):**  $m \equiv 0 \pmod{3}$  and  $n \equiv 0 \pmod{3}$

Let  $m = 3t_1$  and  $n = 3t_2$

Define

$$f(u_i) = 0, 1 \leq i \leq t_1$$

$$f(u_{t_1+i}) = 2, 1 \leq i \leq 2t_1$$

$$f(v_i) = 0, 2 \leq i \leq t_2 - 1$$

$$f(v_{t_2}) = f(v_{t_2+1}) = 1,$$

$$f(v_{t_2+1+i}) = 2, 1 \leq i \leq 2t_2 - 1$$

Then  $f(0) = 2t_1 + 2t_2$  and  $f(1) = f(2) = 2t_1 + 2t_2 - 1$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (ii):**  $m \equiv 0 \pmod{3}$  and  $n \equiv 1 \pmod{3}$

Let  $m = 3t_1$  and  $n = 3t_2 + 1$ . Assign label 1 to the vertex  $v_n$  and assign labels to all the remaining vertices as in case (i). In this case  $f(0) = f(1) = f(2) = 2t_1 + 2t_2$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (iii):**  $m \equiv 0 \pmod{3}$  and  $n \equiv 2 \pmod{3}$

Let  $m = 3t_1$  and  $n = 3t_2 + 2$ . Assign labels 2, 1 to the vertices  $v_{n-1}, v_n$  respectively. Then assign the labels to all the remaining vertices as in case(i). Here  $f(0) = 2t_1 + 2t_2$  and  $f(1) = f(2) = 2t_1 + 2t_2 + 1$ .

Hence  $f$  is a 3-Total Product cordial labeling.

**Case (iv):**  $m \equiv 1 \pmod{3}$  and  $n \equiv 0 \pmod{3}$

Let  $m = 3t_1 + 1$  and  $n = 3t_2$ . Assign label 1 to the vertex  $u_m$ . Then assign the labels to all the remaining vertices as in case (i). In this case  $f(0) = f(1) = f(2) = 2t_1 + 2t_2$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (v):**  $m \equiv 1 \pmod{3}$  and  $n \equiv 1 \pmod{3}$

Let  $m=3t_1+1$  and  $n=3t_2+1$ . Assign label 1 to the vertex  $v_n$ . Then assign the labels to all the remaining vertices as in case (iv). Here  $f(0) = 2t_1 + 2t_2$  and  $f(1) = f(2) = 2t_1 + 2t_2 + 1$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (vi):**  $m \equiv 1 \pmod{3}$  and  $n \equiv 2 \pmod{3}$

Let  $m=3t_1+1$  and  $n=3t_2+2$ . Assign label 0 to the vertex  $v_n$ . Then assign the labels to all the remaining vertices as in case (v). In this case  $f(0) = 2t_1 + 2t_2 + 2$  and  $f(1) = f(2) = 2t_1 + 2t_2 + 1$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (vii):**  $m \equiv 2 \pmod{3}$  and  $n \equiv 0 \pmod{3}$

Let  $m=3t_1+2$  and  $n=3t_2$ . Assign labels 2, 0 to the vertices  $u_{m-1}, u_m$  respectively. Then assign the labels to all the remaining vertices as in case (i). In this case  $f(0) = 2t_1 + 2t_2$  and  $f(1) = f(2) = 2t_1 + 2t_2 + 1$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (viii):**  $m \equiv 2 \pmod{3}$  and  $n \equiv 1 \pmod{3}$

Let  $m=3t_1+2$  and  $n=3t_2+1$ . Assign label 0 to the vertex  $v_n$ . Then assign the labels to all the remaining vertices as in case (vii). Here  $f(0) = 2t_1 + 2t_2 + 2$  and  $f(1) = f(2) = 2t_1 + 2t_2 + 1$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Case (ix):**  $m \equiv 2 \pmod{3}$  and  $n \equiv 2 \pmod{3}$

Let  $m=3t_1+2$  and  $n=3t_2+2$ . Assign labels 2, 0 to the vertex  $v_{n-1}, v_n$  respectively. Then assign the labels to all the remaining vertices as in case (vii). Here  $f(0) = f(1) = f(2) = 2t_1 + 2t_2 + 2$ . Hence  $f$  is a 3-Total Product cordial labeling.

**Illustration 2.14:** A 3-Total Product cordial labeling of the dragon  $C_{12} @ P_8$  is




Figure (iv)

## REFERENCES:

- [1] Harary, F., Graph Theory, Addison wisely, New Delhi.
- [2] R. Ponraj, M. Sivakumar, M. Sundaram, k-Product cordial labeling of graphs, Int. J. Contemp. Math. Sciences, Vol. 7, 2012, no. 15, 733 - 742
- [3] R. Ponraj, M. Sivakumar, M. Sundaram, On 3-Total Product cordial graphs International Mathematical Forum, Vol. 7, 2012, no. 31, 1537 – 1546
- [4] R. Ponraj, M. Sivakumar, M. Sundaram, k -Total Product cordial labeling of graphs (Communicated).
- [5] M. Sundaram, R. Ponraj, S. Somasundaram, Total Product cordial labeling of graphs, Bulletin of Pure and Applied Sciences, 25E(2005), 199-203.

Source of support: Nil, Conflict of interest: None Declared