International Journal of Mathematical Archive-3(6), 2012, 2187-2193

On sg-closed sets

S. Balasubramanian*¹, C. Sandhya² and P. Aruna Swathi Vyjayanthi³

¹Department of Mathematics, Govt. Arts College (A), Karur – 639 005, (T.N.), India ²Department of Mathematics, C.S.R. Sarma College, Ongole–523 001, (A.P.), India ³Research Scholar, Dravidian University, Kuppam – 517 425, (A.P.), India

(Received on: 27-03-12; Revised & Accepted on: 17-04-12)

ABSTRACT

T he object of the present paper is to study the notions of minimal sg-closed set, maximal sg-open set, minimal sg-open set and maximal sg-closed set and their basic properties are studied.

Keywords: sg-closed set and minimal sg-closed set, maximal sg-open set, minimal sg-open set and maximal sg-closed set

1. INTRODUCTION:

Norman Levine introduced the concept of generalized closed sets in topological spaces. After him many authors concentrated in this direction and defined more than 25 types of generalized closed sets. Nakaoka and Oda have introduced minimal open sets and maximal open sets, which are subclasses of open sets. A. Vadivel and K. Vairamanickam introduced minimal $rg\alpha$ -open sets and maximal $rg\alpha$ -open sets in topological spaces. S. Balasubramanian and P.A.S. Vyjayanthi introduced minimal v-open sets and maximal v-open sets; minimal v-closed sets and maximal v-closed sets in topological spaces. Recently S. Balasubramanian introduced minimal vg-open sets and maximal vg-open sets; minimal vg-open sets and maximal vg-open sets; minimal vg-open sets and maximal sg-closed sets in topological spaces. Inspired with these developments we further study a new type of closed and open sets namely minimal sg-closed sets, maximal sg-open sets, minimal sg-closed sets is denoted by SGC(X). For any subset A of X its complement, interior, closure, sg-interior, sg-closure are denoted respectively by the symbols A^c , A^o , A^- , $sg(A)^0$ and $sg(A)^-$.

2. PRELIMINARIES:

Definition 2.1: $A \subset X$ is called

- (i) closed [resp: semi closed; v-closed] if its complement if open[resp:semi open; v-open].
- (ii) ra-open [*v*-open] if $\exists U \in \alpha O(X)[RO(X)]$ such that $U \subset A \subset \alpha cl(U)[U \subset A \subset cl(U)]$.
- (iii) semi- θ -open if it is the union of semi-regular sets and its complement is semi- θ -closed.
- (iv) r-closed[α -closed; pre-closed] if A = cl(A^o)[(cl(A^o))^o \subseteq A; cl(A^o) \subseteq A; cl((cl(A))^o) \subseteq A].
- (v) g-closed [rg-closed] if cl $A \subseteq U$ whenever $A \subseteq U$ and U is open[r-open] in X.
- (vi) sg-closed [gs-closed] if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open {open } in X.
- (vii) $rg\alpha$ -closed if $\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $r\alpha$ -open in X.
- (viii) *vg*-closed if $vcl(A) \subseteq U$ whenever $A \subseteq U$ and U is *v*-open in X.

Definition 2.02: Let A⊂X.

- (i) A point $x \in A$ is the *sg*-interior point of A iff $\exists G \in SGO(X, \tau)$ such that $x \in G \subset A$.
- (ii) A point $x \in X$ is said to be an *sg*-limit point of A iff for each $U \in SGO(X)$, $U \cap (A \setminus \{x\}) \neq \phi$.
- (iii) A point $x \in A$ is said to be *sg*-isolated point of A if $\exists U \in SGO(X)$ such that $U \cap A = \{x\}$.

Definition 2.03: Let $A \subset X$.

- (i) Then A is said to be *sg*-discrete if each point of A is *sg*-isolated point of A. The set of all *sg*-isolated points of A is denoted by $I_{sg}(A)$.
- (ii) For any $A \subset X$, the intersection of all *sg*-closed sets containing A is called the *sg*-closure of A and is denoted by $sg(A)^{-}$.
- (iii) For any $A \subset X$, $A \sim sg(A)^0$ is said to be *sg*-border or *sg*-boundary of A and is denoted by $B_{sg}(A)$.
- (iv) For any $A \subset X$, $sg[sg(X \sim A)^{-}]^0$ is said to be the sg-exterior $A \subset X$ and is denoted by $sg(A)^e$.

Corresponding author: S. Balasubramanian*

¹Department of Mathematics, Govt. Arts College (A), Karur – 639 005, (T.N.), India

International Journal of Mathematical Archive- 3 (6), June – 2012

S. Balasubramanian*¹, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/ On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

Definition 2.04: The set of all *sg*-interior points A is said to be *sg*-interior of A and is denoted by $sg(A)^0$.

Theorem 2.01: (i) Let $A \subseteq Y \subseteq X$ and Y is regularly open subspace of X then $A \in SGO(Y, \tau_{Y})$ iff Y is *sg*-open in X (ii) Let $Y \subseteq X$ and A is a *sg*-neighborhood of x in Y. Then A is *sg*-neighborhood of x in Y iff Y is *sg*-open in X.

Theorem 2.02: Arbitrary intersection of *sg*-closed sets is *sg*-closed. More Precisely, Let $\{A_i: i \in I\}$ be a collection of *sg*-closed sets, then $\bigcap_{i \in I} A_i$ is again *sg*-closed.

Note 2: Finite union and finite intersection of *sg*-closed sets is not *sg*-closed in general.

Theorem 2.03: Let $X = X_1 \times X_2$. Let $A_1 \in SGC(X_1)$ and $A_2 \in SGC(X_2)$, then $A_1 \times A_2 \in SGC(X_1 \times X_2)$.

3. Minimal sg-open Sets and Maximal sg-closed Sets:

We now introduce minimal sg-open sets and maximal sg-closed sets in topological spaces as follows.

Definition 3.1: A proper nonempty sg-open subset U of X is said to be a **minimal** sg-open set if any sg-open set contained in U is ϕ or U.

Remark 1: Every Minimal open set is a minimal sg-open set but converse is not true:

Example 1: Let $X = \{a, b, c, d\}$; $\tau = \{\phi, \{a\}, \{a, b, c\}, X\}$. $\{a\}$ is both Minimal open set and Minimal *sg*-open set but $\{b\}$; $\{c\}$ and $\{d\}$ are Minimal *sg*-open but not Minimal open.

Remark 2: From the above example and known results we have the following implications

Theorem 3.1:

(i) Let U be a minimal *sg*-open set and W be a *sg*-open set. Then $U \cap W = \phi$ or $U \subset W$. (ii) Let U and V be minimal *sg*-open sets. Then $U \cap V = \phi$ or U = V.

Proof:

(i) Let U be a minimal *sg*-open set and W be a *sg*-open set. If $U \cap W = \phi$, then there is nothing to prove.

If $U \cap W \neq \phi$. Then $U \cap W \subset U$. Since U is a minimal *sg*-open set, we have $U \cap W = U$. Therefore $U \subset W$.

(ii) Let U and V be minimal sg-open sets. If $U \cap V \neq \phi$, then $U \subset V$ and $V \subset U$ by (i). Therefore U = V.

Theorem 3.2: Let U be a minimal *sg*-open set. If $x \in U$, then $U \subset W$ for any regular open neighborhood W of x.

Proof: Let U be a minimal *sg*-open set and x be an element of U. Suppose \exists a regular open neighborhood W of x such that $U \not\subset W$. Then $U \cap W$ is a *sg*-open set such that $U \cap W \subset U$ and $U \cap W \neq \phi$. Since U is a minimal *sg*-open set, we have $U \cap W = U$. That is $U \subset W$, which is a contradiction for $U \not\subset W$. Therefore $U \subset W$ for any regular open neighborhood W of x.

Theorem 3.3: Let U be a minimal sg-open set. If $x \in U$, then $U \subset W$ for some sg-open set W containing x.

Theorem 3.4: Let U be a minimal *sg*-open set. Then $U = \bigcap \{W: W \in SGO(X, x)\}$ for any element x of U.

Proof: By theorem [3.3] and U is sg-open set containing x, we have $U \subset \cap \{W: W \in SGO(X, x)\} \subset U$.

Theorem 3.5: Let U be a nonempty sg-open set. Then the following three conditions are equivalent.

(i) U is a minimal *sg*-open set

(ii) $U \subset sg(S)^-$ for any nonempty subset S of U

(iii) $sg(U)^{-} = sg(S)^{-}$ for any nonempty subset S of U.

Proof: (i) \Rightarrow (ii) Let $x \in U$; U be minimal *sg*-open set and $S \neq \phi \subset U$. By theorem[3.3], for any *sg*-open set W containing x, $S \subset U \subset W \Rightarrow S \subset W$. Now $S = S \cap U \subset S \cap W$. Since $S \neq \phi$, $S \cap W \neq \phi$. Since W is any *sg*-open set containing x, by theorem [5.03], $x \in sg(S)^-$. That is $x \in U \Rightarrow x \in sg(S)^- \Rightarrow U \subset sg(S)^-$ for any nonempty subset S of U.

(ii) \Rightarrow (iii) Let S be a nonempty subset of U. That is $S \subset U \Rightarrow sg(S)^- \subset sg(U)^- \rightarrow (1)$. Again from (ii) $U \subset sg(S)^-$ for any $S(\neq \phi) \subset U \Rightarrow sg(U)^- \subset sg(sg(S)^-)^- = sg(S)^-$. That is $sg(U)^- \subset sg(S)^- \rightarrow (2)$. From (1) and (2), we have $sg(U)^- = sg(S)^-$ for any nonempty subset S of U.

S. Balasubramanian*¹, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

(iii) \Rightarrow (i) From (3) we have $sg(U)^- = sg(S)^-$ for any nonempty subset S of U. Suppose U is not a minimal sg-open set.

Then \exists a nonempty *sg*-open set V such that $V \subset U$ and $V \neq U$. Now \exists an element a in U such that $a \notin V \Rightarrow a \in V^c$. That is $sg(\{a\})^- \subset sg(V^c)^- = V^c$, as V^c is *sg*-closed set in X. It follows that $sg(\{a\})^- \neq sg(U)^-$. This is a contradiction for $sg(\{a\})^- = sg(U)^-$ for any $\{a\}(\neq \phi) \subset U$. Therefore U is a minimal *sg*-open set.

Theorem 3.6: Let V be a nonempty finite *sg*-open set. Then \exists at least one (finite) minimal *sg*-open set U such that U \subset V.

Proof: Let V be a nonempty finite *sg*-open set. If V is a minimal *sg*-open set, we may set U = V. If V is not a minimal *sg*-open set, then \exists (finite) *sg*-open set V₁ such that $\phi \neq V_1 \subset V$. If V₁ is a minimal *sg*-open set, we may set $U = V_1$. If V₁ is not a minimal *sg*-open set, then \exists (finite) *sg*-open set V₂ such that $\phi \neq V_2 \subset V_1$. Continuing this process, we have a sequence of *sg*-open sets $V \supset V_1 \supset V_2 \supset V_3 \supset \dots \supset V_k \supset \dots$. Since V is a finite set, this process repeats only finitely. Then finally we get a minimal *sg*-open set $U = V_n$ for some positive integer n.

[A topological space X is said to be locally finite space if each of its elements is contained in a finite open set.]

Corollary 3.1: Let X be a locally finite space and V be a nonempty *sg*-open set. Then \exists at least one (finite) minimal *sg*-open set U such that $U \subset V$.

Proof: Let X be a locally finite space and V be a nonempty *sg*-open set. Let x in V. Since X is locally finite space, we have a finite open set V_x such that x in V_x . Then $V \cap V_x$ is a finite *sg*-open set. By Theorem 3.6 \exists at least one (finite) minimal *sg*-open set U such that $U \subset V \cap V_x$. That is $U \subset V \cap V_x \subset V$. Hence \exists at least one (finite) minimal *sg*-open set U such that $U \subset V \cap V_x$.

Corollary 3.2: Let V be a finite minimal open set. Then \exists at least one (finite) minimal *sg*-open set U such that $U \subset V$.

Proof: Let V be a finite minimal open set. Then V is a nonempty finite *sg*-open set. By Theorem 3.6, \exists at least one (finite) minimal *sg*-open set U such that $U \subset V$.

Theorem 3.7: Let U; U_{λ} be minimal *sg*-open sets for any element $\lambda \in \Gamma$. If U $\subset \cup_{\lambda \in \Gamma} U_{\lambda}$, then \exists an element $\lambda \in \Gamma$ such that U = U_{λ}.

Proof: Let $U \subset \bigcup_{\lambda \in \Gamma} U_{\lambda}$. Then $U \cap (\bigcup_{\lambda \in \Gamma} U_{\lambda}) = U$. That is $\bigcup_{\lambda \in \Gamma} (U \cap U_{\lambda}) = U$. Also by theorem[3.1] (ii), $U \cap U_{\lambda} = \phi$ or $U = U_{\lambda}$ for any $\lambda \in \Gamma$. It follows that \exists an element $\lambda \in \Gamma$ such that $U = U_{\lambda}$.

Theorem 3.8: Let U; U_{λ} be minimal *sg*-open sets for any $\lambda \in \Gamma$. If U = U_{λ} for any $\lambda \in \Gamma$, then $(\cup_{\lambda \in \Gamma} U_{\lambda}) \cap U = \phi$.

Proof: Suppose that $(\bigcup_{\lambda \in \Gamma} U_{\lambda}) \cap U \neq \phi$. That is $\bigcup_{\lambda \in \Gamma} (U_{\lambda} \cap U) \neq \phi$. Then \exists an element $\lambda \in \Gamma$ such that $U \cap U_{\lambda} \neq \phi$. By theorem 3.1(ii), we have $U = U_{\lambda}$, which contradicts the fact that $U \neq U_{\lambda}$ for any $\lambda \in \Gamma$. Hence $(\bigcup_{\lambda \in \Gamma} U_{\lambda}) \cap U = \phi$.

We now introduce maximal sg-closed sets in topological spaces as follows.

Definition 3.2: A proper nonempty *sg*-closed $F \subset X$ is said to be **maximal** *sg***-closed set** if any *sg*-closed set containing F is either X or F.

Remark 3: Every Maximal closed set is maximal sg-closed set but not conversely

Example 2: In Example 1, {b, c, d} is Maximal closed and Maximal *sg*-closed but {a, b, c}, {a, b, d} and {a, c, d} are Maximal *sg*-closed but not Maximal closed.

Remark 4: From the known results and by the above example we have the following implications:

Theorem 3.9: A proper nonempty subset F of X is maximal sg-closed set iff X-F is a minimal sg-open set.

Proof: Let F be a maximal *sg*-closed set. Suppose X-F is not a minimal *sg*-open set. Then \exists *sg*-open set $U \neq X$ -F such that $\phi \neq U \subset X$ -F. That is $F \subset X$ -U and X-U is a *sg*-closed set which is a contradiction for F is a minimal *sg*-open set.

Conversely let X-F be a minimal *sg*-open set. Suppose F is not a maximal *sg*-closed set. Then \exists *sg*-closed set $E \neq F$ such that $F \subset E \neq X$. That is $\phi \neq X \cdot E \subset X$ -F and X-E is a *sg*-open set which is a contradiction for X-F is a minimal *sg*-open set. Therefore F is a maximal *sg*-closed set.

S. Balasubramanian*¹, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

Theorem 3.10:

(i) Let F be a maximal *sg*-closed set and W be a *sg*-closed set. Then $F \cup W = X$ or $W \subset F$. (ii) Let F and S be maximal *sg*-closed sets. Then $F \cup S = X$ or F = S.

Proof: (i) Let F be a maximal *sg*-closed set and W be a *sg*-closed set. If $F \cup W = X$, then there is nothing to prove.

Suppose $F \cup W \neq X$. Then $F \subset F \cup W$. Therefore $F \cup W = F \Rightarrow W \subset F$.

(ii) Let F and S be maximal *sg*-closed sets. If $F \cup S \neq X$, then we have $F \subset S$ and $S \subset F$ by (i). Therefore F = S.

Theorem 3.11: Let F be a maximal *sg*-closed set. If x is an element of F, then for any *sg*-closed set S containing x, $F \cup S = X$ or $S \subset F$.

Proof: Let F be a maximal *sg*-closed set and x is an element of F. Suppose $\exists sg$ -closed set S containing x such that $F \cup S \neq X$. Then $F \subset F \cup S$ and $F \cup S$ is a *sg*-closed set, as the finite union of *sg*-closed sets is a *sg*-closed set. Since F is a *sg*-closed set, we have $F \cup S = F$. Therefore $S \subset F$.

Theorem 3.12: Let F_{α} , F_{β} , F_{δ} be maximal *sg*-closed sets such that $F_{\alpha} \neq F_{\beta}$. If $F_{\alpha} \cap F_{\beta} \subset F_{\delta}$, then either $F_{\alpha} = F_{\delta}$ or $F_{\beta} = F_{\delta}$

Proof: Given that $F_{\alpha} \cap F_{\beta} \subset F_{\delta}$. If $F_{\alpha} = F_{\delta}$ then there is nothing to prove.

If $F_{\alpha} \neq F_{\delta}$ then we have to prove $F_{\beta} = F_{\delta}$. Now $F_{\beta} \cap F_{\delta} = F_{\beta} \cap (F_{\delta} \cap X) = F_{\beta} \cap (F_{\delta} \cap (F_{\alpha} \cup F_{\beta})(by \text{ thm. } 3.10 (ii)) = F_{\beta} \cap ((F_{\delta} \cap F_{\alpha}) \cup (F_{\delta} \cap F_{\beta})) = (F_{\beta} \cap F_{\delta} \cap F_{\alpha}) \cup (F_{\beta} \cap F_{\delta} \cap F_{\beta}) = (F_{\alpha} \cap F_{\beta}) \cup (F_{\delta} \cap F_{\beta}) (by F_{\alpha} \cap F_{\beta} \subset F_{\delta}) = (F_{\alpha} \cup F_{\delta}) \cap F_{\beta} = X \cap F_{\beta}$ (Since F_{α} and F_{δ} are maximal *sg*-closed sets by theorem[3.10](ii), $F_{\alpha} \cup F_{\delta} = X) = F_{\beta}$. That is $F_{\beta} \cap F_{\delta} = F_{\beta} \Rightarrow F_{\beta} \subset F_{\delta}$ Since F_{β} and F_{δ} are maximal *sg*-closed sets, we have $F_{\beta} = F_{\delta}$. Therefore $F_{\beta} = F_{\delta}$.

Theorem 3.13: Let F_{α} , F_{β} and F_{δ} be different maximal *sg*-closed sets to each other. Then $(F_{\alpha} \cap F_{\beta}) \not\subset (F_{\alpha} \cap F_{\delta})$.

Proof: Let $(F_{\alpha} \cap F_{\beta}) \subset (F_{\alpha} \cap F_{\delta}) \Rightarrow (F_{\alpha} \cap F_{\beta}) \cup (F_{\delta} \cap F_{\beta}) \subset (F_{\alpha} \cap F_{\delta}) \cup (F_{\delta} \cap F_{\beta}) \Rightarrow (F_{\alpha} \cup F_{\delta}) \cap F_{\beta} \subset F_{\delta} \cap (F_{\alpha} \cup F_{\beta})$. Since by theorem 3.10(ii), $F_{\alpha} \cup F_{\delta} = X$ and $F_{\alpha} \cup F_{\beta} = X \Rightarrow X \cap F_{\beta} \subset F_{\delta} \cap X \Rightarrow F_{\beta} \subset F_{\delta}$ From the definition of maximal *sg*-closed set it follows that $F_{\beta} = F_{\delta}$, which is a contradiction to the fact that F_{α} , F_{β} and F_{δ} are different to each other. Therefore $(F_{\alpha} \cap F_{\beta}) \not\subset (F_{\alpha} \cap F_{\delta})$.

Theorem 3.14: Let F be a maximal *sg*-closed set and x be an element of F. Then $F = \bigcup \{S: S \text{ is a } sg\text{-closed set containing x such that } F \cup S \neq X\}.$

Proof: By theorem 3.12 and fact that F is a *sg*-closed set containing x, we have $F \subset \bigcup \{S: S \text{ is a } sg\text{-closed set containing x such that } F \cup S \neq X\} - F$. Therefore we have the result.

Theorem 3.15: Let F be a proper nonempty cofinite *sg*-closed set. Then \exists (cofinite) maximal *sg*-closed set E such that $F \subset E$.

Proof: If F is maximal *sg*-closed set, we may set E = F. If F is not a maximal *sg*-closed set, then \exists (cofinite) *sg*-closed set F₁ such that $F \subset F_1 \neq X$. If F₁ is a maximal *sg*-closed set, we may set $E = F_1$. If F₁ is not a maximal *sg*-closed set, then \exists a (cofinite) *sg*-closed set F₂ such that $F \subset F_1 \subset F_2 \neq X$. Continuing this process, we have a sequence of *sg*-closed, $F \subset F_1 \subset F_2 \subset ... \subset F_k \subset ...$ Since F is a cofinite set, this process repeats only finitely. Then, finally we get a maximal *sg*-closed set $E = E_n$ for some positive integer n.

Theorem 3.16: Let F be a maximal *sg*-closed set. If x is an element of X-F. Then $X-F \subset E$ for any *sg*-closed set E containing x.

Proof: Let F be a maximal *sg*-closed set and x in X-F. $E \not\subset F$ for any *sg*-closed set E containing x. Then $E \cup F = X$ by theorem 3.10(ii). Therefore X-F $\subset E$.

4. Minimal sg-Closed set and Maximal sg-open set:

We now introduce minimal sg-closed sets and maximal sg-open sets in topological spaces as follows.

Definition 4.1: A proper nonempty *sg*-closed subset F of X is said to be a **minimal** *sg*-closed set if any *sg*-closed set contained in F is ϕ or F.

S. Balasubramanian^{*1}, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/ On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

Remark 5: Every Minimal closed set is minimal *sg*-closed set but not conversely:

Example 3: Let $X = \{a, b, c, d\}$; $\tau = \{\phi, \{b\}, \{a, b\}, \{b, c\}, \{a, b, c\}, X\}$. {d} is both Minimal closed set and Minimal *sg*-closed set but {a}, {b} and {c} are Minimal *sg*-closed but not Minimal closed.

Definition 4.2: A proper nonempty *sg*-open $U \subset X$ is said to be a **maximal** *sg*-open set if any *sg*-open set containing U is either X or U.

Remark 6: Every Maximal open set is maximal *sg*-open set but not conversely.

Example 4: In Example 3. $\{a, b, c\}$ is Maximal open set and maximal *sg*-open set but $\{a, b, d\}$, $\{a, c, d\}$ and $\{b, c, d\}$ are Maximal *sg*-open but not maximal open.

Theorem 4.1: A proper nonempty subset U of X is maximal sg-open set iff X-U is a minimal sg-closed set.

Proof: Let U be a maximal sg-open set. Suppose X-U is not a minimal sg-closed set. Then $\exists sg$ -closed set $V \neq X$ -U such that $\phi \neq V \subset X$ -U. That is $U \subset X$ -V and X-V is a sg-open set which is a contradiction for U is a minimal sg-closed set. Conversely let X-U be a minimal sg-closed set. Suppose U is not a maximal sg-open set. Then $\exists sg$ -open set $E \neq U$ such that $U \subset E \neq X$. That is $\phi \neq X$ -E $\subset X$ -U and X-E is a sg-closed set which is a contradiction for X-U is a minimal sg-closed set. Therefore U is a maximal sg-closed set.

Lemma 4.1:

(i) Let U be a minimal *sg*-closed set and W be a *sg*- closed set. Then $U \cap W = \phi$ or U subset W. (ii) Let U and V be minimal *sg*- closed sets. Then $U \cap V = \phi$ or U = V.

Proof: (i) Let U be a minimal *sg*-closed set and W be a *sg*-closed set. If $U \cap W = \phi$, then there is nothing to prove.

If $U \cap W \neq \phi$. Then $U \cap W \subset U$. Since U is a minimal *sg*-closed set, we have $U \cap W = U$. Therefore $U \subset W$.

(ii) Let U and V be minimal sg-closed sets. If $U \cap V \neq \phi$, then $U \subset V$ and $V \subset U$ by (i). Therefore U = V.

Theorem 4.2: Let U be a minimal sg-closed set. If $x \in U$, then $U \subset W$ for any regular open neighborhood W of x.

Proof: Let U be a minimal *sg*-closed set and x be an element of U. Suppose \exists an regular open neighborhood W of x such that $U \not\subset W$. Then $U \cap W$ is a *sg*-closed set such that $U \cap W \subset U$ and $U \cap W \neq \phi$. Since U is a minimal *sg*-closed set, we have $U \cap W = U$. That is $U \subset W$, which is a contradiction for $U \not\subset W$. Therefore $U \subset W$ for any regular open neighborhood W of x.

Theorem 4.3: Let U be a minimal sg-closed set. If $x \in U$, then $U \subset W$ for some sg-closed set W containing x.

Theorem 4.4: Let U be a minimal *sg*-closed set. Then $U = \bigcap \{W: W \in SGO(X, x)\}$ for any element x of U.

Proof: By theorem[4.3] and U is *sg*-closed set containing x, we have $U \subset \cap \{W: W \in SGO(X, x)\} \subset U$.

Theorem 4.5: Let U be a nonempty sg-closed set. Then the following three conditions are equivalent.

(i) U is a minimal *sg*-closed set

(ii) $U \subset sg(S)^-$ for any nonempty subset S of U

(iii) $sg(U)^{-} = sg(S)^{-}$ for any nonempty subset S of U.

Proof: (i) \Rightarrow (ii) Let $x \in U$; U be minimal *sg*-closed set and $S \neq \phi \subset U$. By theorem[4.3], for any *sg*-closed set W containing x, $S \subset U \subset W \Rightarrow S \subset W$. Now $S = S \cap U \subset S \cap W$. Since $S \neq \phi$, $S \cap W \neq \phi$. Since W is any *sg*-closed set containing x, by theorem [4.3], $x \in sg(S)^-$. That is $x \in U \Rightarrow x \in sg(S)^- \Rightarrow U \subset sg(S)^-$ for any nonempty subset S of U.

(ii) \Rightarrow (iii) Let S be a nonempty subset of U. That is $S \subset U \Rightarrow sg(S)^- \subset sg(U)^- \rightarrow (1)$. Again from (ii) $U \subset sg(S)^-$ for any $S(\neq \phi) \subset U \Rightarrow sg(U)^- \subset sg(sg(S)^-)^- = sg(S)^-$. That is $sg(U)^- \subset sg(S)^- \rightarrow (2)$. From (1) and (2), we have $sg(U)^- = sg(S)^-$ for any nonempty subset S of U.

(iii) \Rightarrow (i) From (3) we have $sg(U)^- = sg(S)^-$ for any nonempty subset S of U. Suppose U is not a minimal sg-closed set. Set. Then \exists a nonempty sg-closed set V such that $V \subset U$ and $V \neq U$. Now \exists an element a in U such that $a \notin V \Rightarrow a \in V^c$. That is $sg(\{a\})^- \subset sg(V^c)^- = V^c$, as V^c is sg-closed set in X. It follows that $sg(\{a\})^- \neq sg(U)^-$. This is a contradiction for $sg(\{a\})^- = sg(U)^-$ for any $\{a\}(\neq \phi) \subset U$. Therefore U is a minimal sg-closed set.

S. Balasubramanian^{*1}, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/ On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

Theorem 4.6: Let V be a nonempty finite *sg*-closed set. Then \exists at least one (finite) minimal *sg*-closed set U such that $U \subset V$.

Proof: Let V be a nonempty finite *sg*-closed set. If V is a minimal *sg*-closed set, we may set U = V. If V is not a minimal *sg*-closed set, then \exists (finite) *sg*-closed set V_1 such that $\phi \neq V_1 \subset V$. If V_1 is a minimal *sg*-closed set, we may set $U = V_1$. If V_1 is not a minimal *sg*-closed set, then \exists (finite) *sg*-closed set V_2 such that $\phi \neq V_2 \subset V_1$. Continuing this process, we have a sequence of *sg*-closed sets $V \supset V_1 \supset V_2 \supset V_3 \supset \dots \supset V_k \supset \dots$. Since V is a finite set, this process repeats only finitely. Then finally we get a minimal *sg*-closed set $U = V_n$ for some positive integer n.

Corollary 4.1: Let X be a locally finite space and V be a nonempty *sg*-closed set. Then \exists at least one (finite) minimal *sg*-closed set U such that $U \subset V$.

Proof: Let X be a locally finite space and V be a nonempty *sg*-closed set. Let x in V. Since X is locally finite space, we have a finite open set V_x such that x in V_x . Then $V \cap V_x$ is a finite *sg*-closed set. By Theorem 4.6 \exists at least one (finite) minimal *sg*-closed set U such that $U \subset V \cap V_x$. That is $U \subset V \cap V_x \subset V$. Hence \exists at least one (finite) minimal *sg*-closed set U such that $U \subset V \cap V_x$.

Corollary 4.2: Let V be a finite minimal open set. Then \exists at least one (finite) minimal *sg*-closed set U such that U \subset V.

Proof: Let V be a finite minimal open set. Then V is a nonempty finite *sg*-closed set. By Theorem 4.6, \exists at least one (finite) minimal *sg*-closed set U such that U \subset V.

Theorem 4.7: Let U; U_{λ} be minimal *sg*-closed sets for any element $\lambda \in \Gamma$. If U $\subset \cup_{\lambda \in \Gamma} U_{\lambda}$, then \exists an element $\lambda \in \Gamma$ such that U = U_{λ}.

Proof: Let $U \subset \bigcup_{\lambda \in \Gamma} U_{\lambda}$. Then $U \cap (\bigcup_{\lambda \in \Gamma} U_{\lambda}) = U$. That is $\bigcup_{\lambda \in \Gamma} (U \cap U_{\lambda}) = U$. Also by lemma[4.1] (ii), $U \cap U_{\lambda} = \phi$ or $U = U_{\lambda}$ for any $\lambda \in \Gamma$. It follows that \exists an element $\lambda \in \Gamma$ such that $U = U_{\lambda}$.

Theorem 4.8: Let U; U_{λ} be minimal *sg*-closed sets for any $\lambda \in \Gamma$. If U = U_{λ} for any $\lambda \in \Gamma$, then ($\cup_{\lambda \in \Gamma} U_{\lambda}$) \cap U = ϕ .

Proof: Suppose that $(\bigcup_{\lambda \in \Gamma} U_{\lambda}) \cap U \neq \phi$. That is $\bigcup_{\lambda \in \Gamma} (U_{\lambda} \cap U) \neq \phi$. Then \exists an element $\lambda \in \Gamma$ such that $U \cap U_{\lambda} \neq \phi$. By lemma [4.1](ii), we have $U = U_{\lambda}$, which contradicts the fact that $U \neq U_{\lambda}$ for any $\lambda \in \Gamma$. Hence $(\bigcup_{\lambda \in \Gamma} U_{\lambda}) \cap U = \phi$.

Theorem 4.9: A proper nonempty subset F of X is maximal sg-open set iff X-F is a minimal sg-closed set.

Proof: Let F be a maximal *sg*-open set. Suppose X-F is not a minimal *sg*-open set. Then \exists *sg*-open set $U \neq X$ -F such that $\phi \neq U \subset X$ -F. That is $F \subset X$ -U and X-U is a *sg*-open set which is a contradiction for F is a minimal *sg*-closed set.

Conversely let X-F be a minimal *sg*-open set. Suppose F is not a maximal *sg*-open set. Then \exists *sg*-open set $E \neq F$ such that $F \subset E \neq X$. That is $\phi \neq X$ -E $\subset X$ -F and X-E is a *sg*-open set which is a contradiction for X-F is a minimal *sg*-closed set. Therefore F is a maximal *sg*-open set.

Theorem 4.10:

(i) Let F be a maximal *sg*-open set and W be a *sg*-open set. Then $F \cup W = X$ or $W \subset F$. (ii) Let F and S be maximal *sg*-open sets. Then $F \cup S = X$ or F = S.

Proof: (i) Let F be a maximal *sg*-open set and W be a *sg*-open set. If $F \cup W = X$, then there is nothing to prove. Suppose $F \cup W \neq X$. Then $F \subset F \cup W$. Therefore $F \cup W = F \Rightarrow W \subset F$. (ii) Let F and S be maximal *sg*-open sets. If $F \cup S \neq X$, then we have $F \subset S$ and $S \subset F$ by (i). Therefore F = S.

Theorem 4.11: Let F be a maximal *sg*-open set. If x is an element of F, then for any *sg*-open set S containing x, $F \cup S = X$ or $S \subset F$.

Proof: Let F be a maximal *sg*-open set and x is an element of F. Suppose $\exists sg$ -open set S containing x such that $F \cup S \neq X$. Then $F \subset F \cup S$ and $F \cup S$ is a *sg*-open set, as the finite union of *sg*-open sets is a *sg*-open set. Since F is a *sg*-open set, we have $F \cup S = F$. Therefore $S \subset F$.

Theorem 4.12: Let F_{α} , F_{β} , F_{δ} be maximal *sg*-open sets such that $F_{\alpha} \neq F_{\beta}$. If $F_{\alpha} \cap F_{\beta} \subset F_{\delta}$, then either $F_{\alpha} = F_{\delta}$ or $F_{\beta} = F_{\delta}$

Proof: Given that $F_{\alpha} \cap F_{\beta} \subset F_{\delta}$. If $F_{\alpha} = F_{\delta}$ then there is nothing to prove.

S. Balasubramanian^{*1}, C. Sandhya² and P. Aruna Swathi Vyjayanthi³/On sg-closed sets/ IJMA- 3(6), June-2012, Page: 2187-2193

If $F_{\alpha} \neq F_{\delta}$ then we have to prove $F_{\beta} = F_{\delta}$. Now $F_{\beta} \cap F_{\delta} = F_{\beta} \cap (F_{\delta} \cap X) = F_{\beta} \cap (F_{\delta} \cap (F_{\alpha} \cup F_{\beta})(by thm. 4.10 (ii)) = F_{\beta} \cap ((F_{\delta} \cap F_{\alpha}) \cup (F_{\delta} \cap F_{\beta})) = (F_{\beta} \cap F_{\delta} \cap F_{\alpha}) \cup (F_{\beta} \cap F_{\delta} \cap F_{\beta}) = (F_{\alpha} \cap F_{\beta}) \cup (F_{\delta} \cap F_{\beta}) (by F_{\alpha} \cap F_{\beta} \subset F_{\delta}) = (F_{\alpha} \cup F_{\delta}) \cap F_{\beta} = X \cap F_{\beta}$ (Since F_{α} and F_{δ} are maximal *sg*-open sets by theorem[4.10](ii), $F_{\alpha} \cup F_{\delta} = X) = F_{\beta}$. That is $F_{\beta} \cap F_{\delta} = F_{\beta} \Rightarrow F_{\beta} \subset F_{\delta}$ Since F_{β} and F_{δ} are maximal *sg*-open sets, we have $F_{\beta} = F_{\delta}$ Therefore $F_{\beta} = F_{\delta}$

Theorem 4.13: Let F_{α} , F_{β} and F_{δ} be different maximal *sg*-open sets to each other. Then $(F_{\alpha} \cap F_{\beta}) \not\subset (F_{\alpha} \cap F_{\delta})$.

Proof: Let $(F_{\alpha} \cap F_{\beta}) \subset (F_{\alpha} \cap F_{\delta}) \Rightarrow (F_{\alpha} \cap F_{\beta}) \cup (F_{\delta} \cap F_{\beta}) \subset (F_{\alpha} \cap F_{\delta}) \cup (F_{\delta} \cap F_{\beta}) \Rightarrow (F_{\alpha} \cup F_{\delta}) \cap F_{\beta} \subset F_{\delta} \cap (F_{\alpha} \cup F_{\beta})$. Since by theorem 4.10(ii), $F_{\alpha} \cup F_{\delta} = X$ and $F_{\alpha} \cup F_{\beta} = X \Rightarrow X \cap F_{\beta} \subset F_{\delta} \cap X \Rightarrow F_{\beta} \subset F_{\delta}$ From the definition of maximal *sg*-open set it follows that $F_{\beta} = F_{\delta}$, which is a contradiction to the fact that F_{α} , F_{β} and F_{δ} are different to each other. Therefore $(F_{\alpha} \cap F_{\beta}) \not\subset (F_{\alpha} \cap F_{\delta})$.

Theorem 4.14: Let F be a maximal *sg*-open set and x be an element of F. Then $F = \bigcup \{S: S \text{ is a } sg\text{-open set containing x such that } F \cup S \neq X \}.$

Proof: By theorem 4.12 and fact that F is a *sg*-open set containing x, we have $F \subset \bigcup \{S: S \text{ is a } sg$ -open set containing x such that $F \cup S \neq X\} - F$. Therefore we have the result.

Theorem 4.15: Let F be a proper nonempty cofinite *sg*-open set. Then \exists (cofinite) maximal *sg*-open set E such that F \subset E.

Proof: If F is maximal *sg*-open set, we may set E = F. If F is not a maximal *sg*-open set, then \exists (cofinite) *sg*-open set F_1 such that $F \subseteq F_1 \neq X$. If F_1 is a maximal *sg*-open set, we may set $E = F_1$. If F_1 is not a maximal *sg*-open set, then \exists a (cofinite) *sg*-open set F_2 such that $F \subseteq F_1 \subseteq F_2 \neq X$. Continuing this process, we have a sequence of *sg*-open, $F \subseteq F_1 \subseteq F_2 \subseteq ... \subseteq F_k \subseteq ... \subseteq F_k \subseteq ...$ Since F is a cofinite set, this process repeats only finitely. Then, finally we get a maximal *sg*-open set $E = E_n$ for some positive integer n.

Theorem 4.16: Let F be a maximal sg-open set. If x is an element of X-F. Then X-F \subset E for any sg-open set E containing x.

Proof: Let F be a maximal *sg*-open set and x in X-F. $E \not\subset F$ for any *sg*-open set E containing x. Then $E \cup F = X$ by theorem 4.10(ii). Therefore X-F $\subset E$.

Conclusion

The Author is thankful to the referees for their critical comments and suggestions for the development of the paper.

REFERENCES:

[1] S. Balasubramanian, C. Sandhya and P. A. S. Vyjayanthi, On v-closed sets (In press) Acta Ciencia Indica

- [2] S. Balasubramanian, On vg-closed sets, Inter. J. Math. Archive. Vol. 2, No. 10 (2011)1909 1915.
- [3] F. Nakaoka and N. Oda, Some Properties of Maximal Open Sets, Int. J. Math. Sci. 21,(2003)1331-1340.

[4] F. Nakaoka and N. Oda, Some Applications of Minimal Open Sets, Int. J. Math. Sci. 27-8, (2001), 471-476.

[5] F. Nakaoka and N. Oda, On Minimal Closed Sets, Proceeding of Topological spaces Theory and its Applications, 5(2003), 19-21.

[6] A. Vadivel and K. Vairamanickam, Minimal $rg\alpha$ -open Sets and Maximal $rg\alpha$ -closed Sets, Submitted to Journal of Tripura Mathematical Society.

Source of support: Nil, Conflict of interest: None Declared