

## APPLICATION OF THE MELLIN TYPE INTEGRAL TRANSFORM IN THE RANGE $[0, 1/a]$

Dr. Rachana Mathur<sup>1</sup> & Sarita Poonia<sup>2\*</sup>

<sup>1</sup>Department of Mathematics, Govt. Dungar College, Bikaner, Rajasthan, India

<sup>2</sup>Research Scholar, Govt. Dungar College, Bikaner, Rajasthan, India

(Received on: 24-05-12; Revised & Accepted on: 11-06-12)

### ABSTRACT

In this paper Laplace operators are used to solve the Mellin Type Integral Transform which can be a technique for solving boundary and initial value problems. This Transforms is applicable in finite interval. This work intends to understand how Laplace operators leads to properties and relations with the Mellin Type Integral Transform. The main aim of this work is to find the relation between Laplace transform and the Mellin Type Integral Transform in  $[0, 1/a]$ , the result have been modified by applying suitable functions which leads to the results of Mellin Type Integral Transform in the interval  $0 \leq t \leq 1/a$ . We illustrate the advantages and use of this transformation Cauchy's Differential Equation have been solved. We have also studied graphical representation of Mellin Type Integral Transform using MATLAB.

**Keywords:** Laplace Transform, Mellin Transform, Integral Transform, Finite Transform.

**AMS Mathematical Classification:** 65D32, 33C90, 26D07, 26D99(2000), 41A60, 44A35.

### 1. INTRODUCTION

The Laplace Transform is used to study the properties of Mellin Type Integral Transform in the range 0 to  $1/a$  and also to show the validity of properties like Linear Property, Scaling Property, Power Property, theorems like Inversion Theorem, Convolution Theorem, Parseval's Theorem, First and Second Shifting Theorems. We obtain the Mellin Type Integral Transform of the  $n^{\text{th}}$  order derivative of  $f(t)$  with respect to  $t$  and Cauchy's Linear Differential equation  $L_2(F(t)) = t^2 f''(t) + t f'(t) + f(t) = 0$  is solved by using this Integral Transform. The solution of differential equation is graphically presented by MATLAB.

### 2. PRELIMINARY RESULTS

Let  $f(x)$  be a given function of  $x$  which is defined for all  $x \geq 0$  and 's' is parameter, then Laplace

$$\text{Transform is } L[f(x)] = \int_0^\infty e^{-sx} f(x) dx \quad (1)$$

Considering  $x = \log(1/at)$  then  $L[f(t)] = \int_0^{1/a} a^s t^{s-1} f(t) dt$ , denoted by  $M[f(t), s, 0, 1/a]$

$$M[f(t), s, 0, 1/a] = \int_0^{1/a} a^s t^{s-1} f(t) dt \quad (2)$$

Thus obtained is a The Mellin Type Integral Transform in  $[0, 1/a]$ ,  $a > 0$

### 3. PROPERTIES

#### 3.1 LINEAR PROPERTY

The Mellin Type Integral Transform is a Linear operator, that is for any functions  $f(t)$  and  $g(t)$ , We have

$$M[\alpha f(t) + \beta g(t), s, 0, \frac{1}{a}] = \alpha M[f(t), s, 0, \frac{1}{a}] + \beta M[g(t), s, 0, \frac{1}{a}] \quad (3)$$

**Corresponding author:** Sarita Poonia<sup>2</sup>, <sup>2</sup>Research Scholar, Govt. Dungar College, Bikaner, Rajasthan, India

### 3.2 SCALING PROPERTY

The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt$$

$$\text{Then } M\left[f(bt), s, 0, \frac{1}{a}\right] = (b^{-s}) \int_0^{1/a} a^s t^{s-1} f(t) dt = (b^{-s}) M\left[f(t), s, 0, \frac{b}{a}\right] \quad (4)$$

### 3.3 POWER PROPERTY

The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt$$

$$\text{Then } M\left[f(t^b), s, 0, \frac{1}{a}\right] = \frac{1}{b} M\left[f(t), \frac{s}{b}, 0, \frac{1}{ab}\right] \quad (5)$$

## 4. MAIN RESULTS

### 4.1 INVERSION THEOREM

**Theorem:** The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt$$

$$\text{Then } f(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{t^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] ds$$

**Proof:** Assume that  $M\left[f(t), s, 0, \frac{1}{a}\right]$  is a regular equation in the strip  $|Re(s)| < r$  ('r' to real number) of the s-plane and that  $0 < c < v$ ,  $c-i\infty \leq s \leq c+i\infty$ , where c is constant then

$$\begin{aligned} M\left[f(t), s, 0, \frac{1}{a}\right] &= \int_0^{1/a} a^s t^{s-1} f(t) dt \\ &= \int_0^{1/a} a^s t^{s-1} \left[ \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^{-s} M\left[f(t), s, 0, \frac{1}{a}\right] ds \right] dt \\ &= \left[ \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^{-s} M\left[f(t), s, 0, \frac{1}{a}\right] ds \right] \int_0^{1/a} a^s t^{s-1} dt \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] ds \\ f(x) &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] ds \end{aligned} \quad (6)$$

### 4.2 CONVOLUTION THEOREM

**Theorem:** The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt \quad \text{and} \quad M\left[g(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} g(t) dt$$

$$\text{Then } M^{-1}\left[f(t), s, 0, \frac{1}{a}\right] = f(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] ds$$

$$M^{-1}\left[g(t), s, 0, \frac{1}{a}\right] = g(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[g(t), s, 0, \frac{1}{a}\right] ds$$

$$M\left[f(t)g(x-t), s, 0, \frac{1}{a}\right] = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] M\left[g(x-t), s, \frac{1}{a}-t, t\right] ds \quad (7)$$

### 4.3 PARSEVALS THEOREM

**Theorem:** The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt \quad \text{and} \quad M\left[g(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} g(t) dt$$

$$\text{Then } M^{-1}\left[f(t), s, 0, \frac{1}{a}\right] = f(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] ds$$

$$M^{-1}\left[g(t), s, 0, \frac{1}{a}\right] = g(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[g(t), s, 0, \frac{1}{a}\right] ds$$

$$M\left[f(t)g(t), s, 0, \frac{1}{a}\right] = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{-s}}{s} M\left[f(t), s, 0, \frac{1}{a}\right] M\left[g(t), s, \frac{1}{a} - t, t\right] ds \quad (8)$$

### 4.4 DEFINITIONS

Heviside Unit Step Function

For  $U(t-a)=H(t-a) = 1$ , when  $t > a$

$$= 0, \text{ when } t < a$$

$U(t-a)$  or  $H(t-a)$  is known as the Heviside Unit Step Function.

### 4.5 FIRST SHIFTING THEOREM

The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt$$

$$\text{Then } M\left[t^n f(t), s, 0, \frac{1}{a}\right] = M[f(t), s + n, 0, \frac{1}{a}] \quad (9)$$

### 4.6 SECOND SHIFTING THEOREM

The Mellin Type Integral Transform in  $[0, 1/a]$  is

$$M\left[f(t), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t) dt$$

$$\text{Then } M\left[f(t-b)U(t-b), s, 0, \frac{1}{a}\right] = M[f(u), s, -b, \frac{1}{a} - b]$$

$$\textbf{Proof: } M\left[f(t-b)U(t-b), s, 0, \frac{1}{a}\right] = \int_0^{1/a} a^s t^{s-1} f(t-b)U(t-b) dt$$

$$\text{Let } u=t-b \text{ then} \quad = \int_{-b}^{\frac{1}{a}-b} a^s (u+b)^{s-1} f(u)U(u) du$$

$$\text{If } U(t-b)=1, \text{ when } t>b \quad = \int_{-b}^{\frac{1}{a}-b} a^s (u+b)^{s-1} f(u) du$$

$$= M[f(u), s, -b, \frac{1}{a} - b] \quad \text{where kernel is } (u+b)^{s-1} \quad (10)$$

## 5. DERIVATIVES

### 5.1 The Mellin Type Integral Transforms of first order derivative of $f(t)$ w.r.t. $t$

Suppose that  $f(t)$  is continuous for all  $t \geq 0$  satisfying (2) for some value  $\gamma$  and  $m$  and it has derivative  $f'(t)$  which is piecewise continuous on every finite interval in the range of  $t \geq 0$ . Then the Mellin Type (finite) Integral Transforms of the derivative of  $f(t)$  exists when  $s > \gamma$  and  $|f(t)| \leq m e^{\gamma t}$  for all  $t \geq 0$  for some constants.

**Proof:** Considering the case when  $f'(t)$  is continuous for all  $t \geq 0$ , then integrating by parts, we get

$$\begin{aligned} M \left[ f'(t), s, 0, \frac{1}{a} \right] &= \int_0^{1/a} a^s t^{s-1} f'(t) dt \\ &= a f \left( \frac{1}{a} \right) - (s-1) M[f(t), s-1, 0, \frac{1}{a}] \end{aligned} \quad (11)$$

## 5.2 The Mellin Type Integral Transforms of $n^{th}$ order derivative of $f(t)$ w.r.t. $t$

$$\begin{aligned} M \left[ f''(t), s, 0, \frac{1}{a} \right] &= \int_0^{1/a} a^s t^{s-1} f''(t) dt \\ &= a f' \left( \frac{1}{a} \right) - (s-1) a^2 f \left( \frac{1}{a} \right) + (s-1)(s-2) M[f(t), s-2, 0, \frac{1}{a}] \end{aligned} \quad (12)$$

$$\begin{aligned} \text{Similarly, } M \left[ f'''(t), s, 0, \frac{1}{a} \right] &= a f'' \left( \frac{1}{a} \right) - (s-1) a^2 f' \left( \frac{1}{a} \right) + (s-1)(s-2) a^3 f \left( \frac{1}{a} \right) \\ &\quad - (s-1)(s-2)(s-3) M[f(t), s-3, 0, \frac{1}{a}] \end{aligned} \quad (13)$$

$$M \left[ f^n(t), s, 0, \frac{1}{a} \right] = a f^{n-1} \left( \frac{1}{a} \right) - (s-1) a^2 f^{n-2} \left( \frac{1}{a} \right) \dots \dots \dots - (s-1)(s-2) \dots \dots (s-n) M[f(t), s, 0, \frac{1}{a}] \quad (14)$$

## 6. APPLICATION

Consider Cauchy differential equation  $L_2(F(t)) = t^2 f''(t) + t f'(t) + f(t)$ ,

$$\begin{aligned} M \left[ t f'(t), s, 0, \frac{1}{a} \right] &= \int_0^{1/a} a^s t^s f'(t) dt \\ &= f \left( \frac{1}{a} \right) - s M[f(t), s, 0, \frac{1}{a}] \\ M \left[ t^2 f''(t), s, 0, \frac{1}{a} \right] &= \int_0^{1/a} a^s t^{s+1} f''(t) dt \\ &= \frac{1}{a} f' \left( \frac{1}{a} \right) - (s+1) f \left( \frac{1}{a} \right) + s(s+1) M[f(t), s, 0, \frac{1}{a}] \end{aligned}$$

$$\begin{aligned} M[L_2(F(t)), s, 0, \frac{1}{a}] &= M[t^2 f''(t) + t f'(t) + f(t), s, 0, \frac{1}{a}] \\ &= \frac{1}{a} f' \left( \frac{1}{a} \right) - s f \left( \frac{1}{a} \right) + (s^2 + 1) M[f(t), s, 0, \frac{1}{a}] \end{aligned}$$

$$\text{If } L_2(F(t)) = 0 \text{ then } M[L_2(F(t)), s, 0, \frac{1}{a}] = 0$$

$$\frac{1}{a} f' \left( \frac{1}{a} \right) - s f \left( \frac{1}{a} \right) + (s^2 + 1) M[f(t), s, 0, \frac{1}{a}] = 0$$


$$M \left[ f(t), s, 0, \frac{1}{a} \right] = \frac{1}{s^2 + 1} \left[ s f \left( \frac{1}{a} \right) - \frac{1}{a} f' \left( \frac{1}{a} \right) \right] \quad (15)$$

## 7. GRAPHICAL REPRESENTATION:


Consider the equation

$$M \left[ f(t), s, 0, \frac{1}{a} \right] = \frac{1}{s^2 + 1} \left[ s f \left( \frac{1}{a} \right) - \frac{1}{a} f' \left( \frac{1}{a} \right) \right]$$


(a)  $f(x)=x^2$  then  $f(a)=a^2$  and let  $a=1$



(b)  $f(x)=e^x$  then  $f(a)=e^a$  and let  $a=1$



(c)  $f(x)=\log(1+x)$  then  $f(a)=\log(1+a)$  and let  $a=1$



## 8. CONCLUSION:

We have seen how the properties of the Laplace operator hold good for the Mellin Type Integral Transform in  $[0, 1/a]$ . The Cauchy Differential Equation is solved and we have given graphical representation of the solution of differential equation by MATLAB. It is observed that Mellin Type Integral Transform in  $[0, 1/a]$  gives better results than other Integral Transforms.

## 9. REFERENCES

- [1] A .E. Gracc and M. Spann, A comparison between Fourier-Mellin descriptors and Moment based features for invariant object recognition using neural network, *Pattern Recog. Lett.*, 12 (1991), 635-643.
- [2] A. H. Zemanian, Generalized Integral Transformation, Interscience Publication, New York (1968)
- [3] A. Z .Zemanian, The distributional Laplace and Mellin transformations, *J. SIAM*, 14(1), 1908.
- [4] Bruce Littlefield, Mastering Matlab, Prentice Hall, Upper saddle River, New York
- [5] C. Fox, Applications of Mellin's Transformation to the integral equations, (1933)
- [6] C.Fox Application of Mellin Transformation to Integral Equation, 3rd March, 1934, 495-502.
- [7] Derek Naylor, On a Mellin Type Integral Transforms, *Journal of Mathematics and Mechanics*, 12(2) (1963).
- [8].Ian N. Sneddon, The use of Integral Transforms, TMH edition 1974.
- [9] I. S. Reed, The Mellin Type Double Integral, Cambridge, London.
- [10] Jean M. Tchuenche and Nyimvua S. Mhare, An applications of double Sumudu Transform, *Applied Mathematical Sciences*, 1 (2007), 31-39.
- [11] J. M. Mendez and J. R.Negrin, On the finite Hankel-Schwartz Transformation of Distributions, *Ganita*, 39(1) (1988).
- [12] Rugra Pratap, Getting Started With Matlab, Oxford University Press, NewYork, 2003.
- [13] Stephane Derrode and Faouzi Ghorbel, Robust and Efficient Fourier-Mellin Transform Appoximations for Gray-Level Image Reconsrruction and Complete Invarient Description, *Computer Vision and Image Understandind* 83 (2001), 57-78.
- [14] S. M. Khairnar, R.M. Pise and J. N. Salunke , Applications of the Mellin type integral transform in the range  $(1/a, \infty)$ , IJMSA (Accepted).
- [15] S. M. Khairnar, R.M. Pise and J. N. Salunke, Study Of The Sumudu Mellin Integral Transform and Its Applications, *Int. J. Mat. Sci. and Engg. Appl.*, 4(IV) (2010), 307-325.
- [16] S. M. Khairnar, R.M. Pise and J. N. Salunke, Bilateral Laplace-Mellin Integral Transform and its Applications, *International Journal of Pure and Applied Sciences and Technology* (Accepted).
- [17] S. M. Khairnar, R.M. Pise and J. N. Salunke, Application Of Mellin Type Integral Transform In The Range  $(1/a, \infty)$ , *International Journal of Mathematical Science and applications* (Accepted).

**Source of support: Nil, Conflict of interest: None Declared**