

COMMONFIXED POINT THEOREM IN BANACH SPACE

Neena Vijaywargiand & Shyam Patkar*

**Department of Mathematics, Truba Institute of Engineering & Information Technology, Bhopal, India*

(Received on: 25-07-12; Accepted on: 18-08-12)

ABSTRACT

In the present paper we establish a fixed point theorem in Banach space taking new rational expression, which satisfies the well-known results.

Key words: Self mapping, Continuous mappings, Non contraction mappings, Banach Space.

AMS subject classification: 47H10, 54H25,

INTRODUCTION&PRELIMINARIES

The study of Non-Contraction mapping concerning the existence of fixed point draws attention of various authors in non linear analysis dealing with the study of Non –expansive mapping and the existence of fixed points.

It is well known that the differential and integral equations that arise in the physical problems are generally non linear, therefore the fixed point methods specially “Banach contraction Principle ([1],1922) provides a powerful tool for obtaining the solution of their equations which were very difficult to solve by any other methods.

It is also true that some qualitative properties of the solution of related equations are proved by functional analysis approach. Many authors have presented valuable results with non contraction mapping ([27], 2007) in Banach space.

Definition 2.1: (Banach space) A Banach space $(X, \|\cdot\|)$ is a normed vector space such that X is complete under the metric induced by the norm $\|\cdot\|$

Example 2.1: The set of continuous functions on closed interval of real line with the norm $\|\cdot\|$ of function f given by

$$\|f\| = \sup_{x \in X} |f(x)|$$

is a Banach space ,where sup denotes the supremum.

Definition2.2: (Normed linear space) let $\|\cdot\|$ denotes a function from a linear space X into R that satisfies the following axioms

- i) $\forall x \in X, \|x\| \geq 0, \|x\| = 0$ iff $x = 0$
- ii) $\forall x, y \in X, \|x + y\| \leq \|x\| + \|y\|$
- iii) $\forall x \in X, \alpha \in R, \|\alpha x\| = |\alpha| \|x\|$

$\|x\|$ is called the norm of x and $(X, \|\cdot\|)$ is called a Normed linear space.

Example -2.2.1: $\langle R^n, \|\cdot\|_p \rangle, \forall x \in R^n, \|x\|_\infty = \max_{i=1}^n |x_i|$

Example -2.2.2: $\langle l_p, \|\cdot\|_p \rangle, 1 \leq p \leq \infty, \forall x \in l_p = \{x: x \in R^\infty, \sum_{i=1}^\infty |x_i|^p < \infty\}, \|x\| = (\sum_{i=1}^\infty |x_i|^p)^{1/p}$

Definition 2.3: Asequence $\{x_n\}$ in a normed space is said to be a Cauchy sequence if $\|x_n - x_m\| \rightarrow 0$ as $m, n \rightarrow \infty$ i.e. given $\epsilon > 0$,there exist an integer N such that $\|x_n - x_m\| < \epsilon$,for all $m, n > N$

*Corresponding author: Shyam Patkar**

**Department of Mathematics, Truba Institute of Engineering & Information Technology, Bhopal, India*

3. MAIN RESULT

Theorem 3.1: Let f be a mapping of a Banach space X into itself. If f satisfies the following conditions;

$$f^2 = I, \text{ where } I \text{ is identity mapping.} \quad (3.1.1)$$

$$\begin{aligned} \|f(x) - f(y)\| &\leq \alpha \left[\frac{\|x-y\|(\|x-f(x)\| + \|x-f(x)\| \|x-f(y)\| + \|x-f(y)\| \|y-f(x)\|)}{\|x-y\| + \|y-f(y)\|} \right] \\ &\quad + \gamma (\|x-f(x)\| + \|y-f(y)\|) + \delta (\|x-f(y)\| + \|y-f(x)\|) + \eta \|x-y\| \end{aligned} \quad (3.1.2)$$

For every $x, y \in X$, where $\alpha, \delta, \eta > 0$ and $5\alpha + 4\gamma + 2\delta + \eta < 2$, then f has a fixed point.

If $\alpha + 2\delta + \eta < 1$ then f has an unique fixed point.

Proof: Suppose x is a point in the Banach space X .

Taking $y = \frac{1}{2}(f + I)(x), z = f(y)$ and $u = 2y - z$ we have

$$\begin{aligned} \|z-x\| &= \|f(y) - f^2(x)\| = \|f(y) - f(f(x))\| \\ &\leq \alpha \left[\frac{\|y-f(x)\|(\|y-f(y)\| + \|y-f(y)\| \|y-f^2(x)\| + \|y-f^2(x)\| \|f(x)-f(y)\|)}{\|y-f(x)\| + \|f(x)-f^2(x)\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|f(x) - f^2(x)\|) + \delta (\|y-f^2(x)\| + \|f(x) - f(y)\|) + \eta \|y-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(x)\|(\|y-f(y)\| + \|y-f(y)\| \|x-y\| + \|x-y\| \|f(x)-f(y)\|)}{\|y-f(x)\| + \|x-f(x)\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta (\|x-y\| + \|f(x) - f(y)\|) + \eta \|y-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(y)\|(\|y-f(x)\| + \|x-y\|) + \|x-y\| \|f(x)-f(y)\|}{\|x-y\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta (\|x-y\| + \|f(x) - f(y)\|) + \eta \|y-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(y)\|(\|x-f(x)\| + \|x-y\|) + \|x-y\| \|f(x)-f(y)\|}{\|x-y\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta (\|x-y\| + \|f(x) - f(y)\|) + \eta \|y-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(y)\|(\|x-f(x)\| + \|x-y\|) + \|x-y\| \|f(x)-f(y)\|}{\|x-y\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta (\|x-y\| + \|f(x) - f(y)\|) + \eta \|y-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(y)\|(\|x-f(x)\| + \|x-y\|) + \|x-y\| \|f(x)-f(y)\|}{\|x-\frac{1}{2}(f+I)(x)\|} \right] + \gamma (\|y-f(y)\| + \|x-f(x)\|) \\ &\quad + \delta \left[\left\| x - \frac{1}{2}(f+I)(x) \right\| + \left\| f(x) - f\left(\frac{1}{2}(f+I)(x)\right) \right\| \right] + \eta \left\| \frac{1}{2}(f+I)(x) - f(x) \right\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[\frac{\|y-f(y)\|(\|x-f(x)\| + \frac{1}{2}\|x-y\|) + \frac{1}{2}\|x-y\| \|f(x)-f(y)\|}{\frac{1}{2}\|x-f(x)\|} \right] \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta \left[\frac{1}{2}\|x-f(x)\| + \frac{1}{2}\|x-y\| \right] + \eta \frac{1}{2}\|x-f(x)\| \end{aligned}$$

$$\begin{aligned} \|z-x\| &\leq \alpha \left[2\|y-f(y)\| + \frac{1}{2}\|x-f(x)\| \right] + \beta \frac{1}{2}\|x-f(x)\| \\ &\quad + \gamma (\|y-f(y)\| + \|x-f(x)\|) + \delta (\|x-f(x)\|) + \eta \frac{1}{2}\|x-f(x)\| \end{aligned}$$

$$\|z-x\| \leq \left(\frac{\alpha}{2} + \gamma + \delta + \frac{\eta}{2} \right) \|x-f(x)\| + (2\alpha + \gamma) \|y-f(y)\| \quad (3.1.3)$$

Also

$$\|u-x\| = \|2y-z-x\| = \left\| \frac{1}{2}(f+I)(x) - z - x \right\| = \|f(x) - z\| = \|f(x) - f(y)\|$$

$$\|u - x\| \leq \alpha \left[\frac{\|x - y\| \|x - f(x)\| + \|x - f(x)\| \|x - f(y)\| + \|x - f(y)\| \|y - f(x)\|}{\|x - y\| + \|y - f(y)\|} \right] \\ + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta [\|x - f(y)\| + \|y - f(x)\|] + \eta \|x - y\|$$

$$\|u - x\| \leq \alpha \left[\frac{\|x - f(x)\| (\|x - y\| + \|x - f(y)\|) + \|x - f(y)\| \|y - f(x)\|}{\|x - f(y)\|} \right] \\ + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta [\|x - f(y)\| + \|y - f(x)\|] + \eta \|x - y\|$$

$$\|u - x\| \leq \alpha \left[\frac{\|x - f(x)\| \|y - f(y)\|}{\|x - f(y)\|} + \|y - f(x)\| \right] \\ + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta [\|x - f(y)\| + \|y - f(x)\|] + \eta \|x - y\|$$

$$\|u - x\| \leq \alpha \left[\frac{\|x - f(x)\| \|y - f(y)\|}{\|x - f(y)\|} + \|y - f(x)\| \right] \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta [\|x - f(y)\| + \|y - f(x)\|] + \eta \|x - y\|$$

$$\|u - x\| \leq \alpha \left[\frac{\|x - f(x)\| \|y - f(y)\|}{\|x - f(\frac{1}{2}(f+I)(x))\|} + \left\| \frac{1}{2}(f+I)(x) - f(x) \right\| \right] \\ + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta \left[\left\| x - f\left(\frac{1}{2}(f+I)(x)\right) \right\| + \left\| \frac{1}{2}(f+I)(x) - f(x) \right\| \right] \\ + \eta \left\| x - \frac{1}{2}(f+I)(x) \right\| \|u - x\|$$

$$\leq \alpha \left[\frac{\|x - f(x)\| \|y - f(y)\|}{\frac{1}{2}\|x - f(x)\|} + \frac{1}{2} \|x - f(x)\| \right] \\ + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta \left[\frac{1}{2} \|x - f(x)\| + \frac{1}{2} \|x - f(x)\| \right] + \eta \frac{1}{2} \|x - f(x)\|$$

$$\|u - x\| \leq \alpha [2\|y - f(y)\| + \frac{1}{2}\|x - f(x)\|] + \gamma [\|x - f(x)\| + \|y - f(y)\|] + \delta \|x - f(x)\| + \eta \frac{1}{2} \|x - f(x)\|$$

$$\|u - x\| \leq (\frac{\alpha}{2} + \gamma + \delta + \frac{\eta}{2}) \|x - f(x)\| + (2\alpha + \gamma) \|y - f(y)\| \quad (3.1.4)$$

Now,

$$\|z - u\| = \|(z - x) - (x - u)\| \leq (\|z - x\| + \|x - u\|)$$

$$\leq [(\frac{\alpha}{2} + \gamma + \delta + \frac{\eta}{2}) \|x - f(x)\| + (2\alpha + \gamma) \|y - f(y)\|] + [(\frac{\alpha}{2} \gamma + \delta + \frac{\eta}{2}) \|x - f(x)\| + (2\alpha + \gamma) \|y - f(y)\|]$$

$$\|z - u\| \leq (\alpha + 2\gamma + 2\delta + \eta) \|x - f(x)\| + (4\alpha + 2\gamma) \|y - f(y)\|$$

Also

$$\begin{aligned} \|z - u\| &= \|f(y) - (2y - z)\| \\ &= \|f(y) - 2y - f(y)\| \\ &= 2\|y - f(y)\| \quad (3.1.5) \end{aligned}$$

From (3.1.5)

$$\begin{aligned} 2\|y - f(y)\| &\leq (\alpha + \beta + 2\lambda + 2\mu + 2\gamma + 2\delta + \eta) \|x - f(x)\| + (4\alpha + 2\lambda + 2\gamma) \|y - f(y)\| \\ &\quad [2 - (4\alpha + 2\gamma + 2\lambda)] \|y - f(y)\| \\ &\leq (\alpha + \beta + 2\lambda + 2\mu + 2\gamma + 2\delta + \eta) \|x - f(x)\| \end{aligned}$$

$$\|y - f(y)\| \leq q \|x - f(x)\|$$

$$\text{Where } q = \frac{(\alpha + \beta + 2\lambda + 2\mu + 2\gamma + 2\delta + \eta)}{[2 - (4\alpha + 2\lambda + 2\gamma)]} < 1$$

Since $5\alpha + 4\gamma + 2\delta + \eta < 2$

Let $g = \frac{1}{2}(f + I)$ then for every $x \in X$

$$\begin{aligned} \|g^2(x) - g(x)\| &= \|g(y) - y\| \\ &= \left\| \frac{1}{2}(f + I)y - y \right\| \\ &= \frac{1}{2}\|y - f(y)\| \\ &\leq \frac{\eta}{2}\|x - f(x)\| \end{aligned}$$

By the definition of η , we claim that $\{g^n(x)\}$ is a Cauchy sequence in X .

By the completeness, $\{g^n(x)\}$ converges to some element x_0 in X

i.e. $\lim_{n \rightarrow \infty} g^n(x) = x_0$

which implies that $g(x_0) = x_0$

hence $f(x_0) = x_0$

i.e. x_0 is fixed point of f

For the uniqueness:

If possible let $y_0 (\neq x_0)$ be another fixed point of f then

$$\begin{aligned} \|x_0 - y_0\| &= \|f(x_0) - f(y_0)\| \\ &\leq \alpha \left[\frac{\|x_0 - y_0\| \|x_0 - f(x_0)\| + \|x_0 - f(x_0)\| \|x_0 - f(y_0)\| + \|x_0 - f(y_0)\| \|y_0 - f(x_0)\|}{\|x_0 - y_0\| + \|y_0 - f(y_0)\|} \right] \\ &\quad + \gamma [\|x_0 - f(x_0)\| + \|y_0 - f(y_0)\|] + \delta [\|x_0 - f(y_0)\| + \|y_0 - f(x_0)\|] + \eta \|x_0 - y_0\| \\ \|x_0 - y_0\| &\leq \alpha \frac{\|x_0 - y_0\|^2}{\|x_0 - y_0\|} + 2\delta \|x_0 - y_0\| + \eta \|x_0 - y_0\| \\ &\leq \alpha \|x_0 - y_0\| + 2\delta \|x_0 - y_0\| + \eta \|x_0 - y_0\| \\ &\leq \alpha \|x_0 - y_0\| + 2\delta \|x_0 - y_0\| + \eta \|x_0 - y_0\| \\ \|x_0 - y_0\| &= (\alpha + 2\delta + \eta) \|x_0 - y_0\| \end{aligned}$$

Since $\alpha + 2\delta + \eta < 1$

$$\|x_0 - y_0\| = 0$$

$$x_0 = y_0$$

This complete the proof

REFERENCES

1. Ahmad A. and Shakil, M. "Some fixed point theorems in Banach spaces" Nonlinear Funct. Anal. And Appl. 11(2006) 343-349.
2. Banach S. "Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales" Fund. Math. 3(1922) 133-181.
3. Badshah V.H. and Gupta, O.P. "Fixed point theorems in Banach and 2 – Banach spaces" Jnanabha 35(2005) 73-78.
4. Browder F.E. "Non-expansive non-linear operators in Banach spaces" Proc. Nat. Acad. Sci. U.S.A. 54 (1965) 1041-1044.
5. Datson W.G. Jr. "Fixed point of quasi non-expansive mappings" J. Austral. Math. Soc. 13 (1972) 167-172.
6. Gohde D. "Zum Prinzip der kontraktiven Abbildung" Math. Nachr 30 (1965) 251-258.

7. Goebel, K. "An elementary proof of the fixed point theorem of Browder and Kirk" Michigan Math. J. 16 (1969) 381-383.
8. Goebel K. and Zlotkiewics, E. "Some fixed point theorems in Banach spaces" Colloq Math 23(1971) 103-106.
9. Goebel K. Kirk, W.A. and Shmidt, T.N. "A fixed point theorem in uniformly convex spaces" Boll. Un. Math. Italy 4 (1973) 67-75.
10. Gahlar S. "2 – Metrcheraume and ihre topologiscche structure" Math. Nadh. 26 (1963-64) 115-148.
11. Isekey K. "Fixed point theorem in Banach space" Math Sem. Notes, Kobe University 2(1974) 111-115.
12. Jong S.J. "Viscosity approximation methods for a family of finite non expansive in Banach spaces" nonlinear Analysis 64(2006) 2536-2552.
13. Khan M.S. "Fixed points and their approximation in Banach spaces for certain commuting mappings" Glasgow Math. Jour. 23(1982) 1-6.
14. Khan M.S. and Imdad, M. "Fixed points of certain involutions in Banach spaces" J. Austral. Math. Soc. 37 (1984) 169-177.
15. Kirk W.A. "A fixed point theorem mappings do not increase distance" Amer. Math. Monthly 72 (1965) 1004-1006.
16. Kirk W.A. "A fixed point theorem for non-expansive mappings" Lecture notes in Math. Springer-Verlag, Berlin and New York 886 (1981) 111-120.
17. Kirk W.A. "Fixed point theorem for non-expansive mappings" Contem Math. 18(1983) 121-140.
18. Pathak H.K. and Maity, A.R. "A fixed point theorem in Banach space" Acta CienciaIndica 17 (1991) 137-139.
19. Qureshi N.A. and Singh, B. "A fixed point theorem in Banach space" Acta CienciaIndica 17 (1995) 282-284.
20. Rajput S.S. and Naroliya, N. "Fixed point theorem in Banach space" Acta CienciaIndica 17 (1991) 469-474.
21. Sgarma P.L. abd Rajput, S.S. "Fixed point theorem in Banach space" Vikram Mathematical Journal 4 (1983) 35-38.
22. Singh M.R. and Chattergee, A.K. "Fixed point theorem in Banach space" Pure Math. Manuscript 6 (19870) 53-61.
23. Sharma S. and Bhagwan, A. "Common fixed point theorems on Normed space" ActaCienciaIndica 31 (2003) 20-24.
24. Shahzad N and Udomene, A. "Fixed point solutions of variational inequalities for asymptotically non-expansive mappings in Banach spaces" Nonlinear Analysis 64(2006) 558-567.
25. Verma B.P. "Application of Banach fixed point theorem to solve non linear equations and its generalization" Jnanabha 36 (2006) 21-23.
26. Yadava R.N., Rajput, S.S. and Bhardwaj, R.K. "Some fixed point and common fixed point theorems in Banach spaces" ActaCienciaIndica 33 No 2 (2007) 453-460.
27. Yadava R.N. Rajput, S.S., Choudhary, S. And Bhardwaj, R.K. "Some fixed point and common fixed point theorems for non-contraction mapping on 2 – Banach spaces" Acta CienciaIndica 33 No. 3)2007) 737-744.
28. Sabhakant Dwivedi, ramakant Bhardwaj, Rajesh Shivastava "Common fixed point theorems for two mappings in 2-Banach spaces" Int. Jour. of Math. Analysis 3(2009) 889-896
29. R. Shrivastav, B. Dwivedi, S.S. rajput "Some common fixed point theorems in Banach spaces" Int. Jour. Of Math Sci. &Engg. Appl.5 (2011).

Source of support: Nil, Conflict of interest: None Declared