

SUM CORDIAL LABELING FOR SOME GRAPHS

J. SHIAMA*

Assistant professor, Department of Mathematics, Cms College of Science and Commerce,
Coimbatore -641049, (TN), india

(Received on: 18-08-12; Revised & Accepted on: 11-09-12)

ABSTRACT

In this paper I investigate a new labeling called the Sum Cordial labeling and some Sum Cordial graphs. A binary vertex labeling of a graph G with induced edge labeling $f^* : E(G) \rightarrow \{0,1\}$ defined by $f^*(uv) = (f(u) + f(v)) \pmod{2}$ is called a sum cordial labeling if $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$. Any graph which satisfies the sum cordial labeling is called a Sum Cordial graph. Here I prove some graphs like path P_n , cycle C_n , star $K_{1,n}$, bistar $B_{n,n}$ and the graphs obtained by joining apex vertices of k copies of double stars to a new vertex are Sum cordial graphs.

Keywords: Sum cordial labeling, Sum Cordial graphs,

INTRODUCTION

All graphs in this paper are finite and undirected. For all other terminology and notations I follow Harary [3]. Let $G(V, E)$ be a graph where the symbols $V(G)$ and $E(G)$ denotes the vertex set and the edge set. The cardinality of the vertex set is called the order of the graph G and it is denoted by p . The cardinality of the edge set is called the size of the graph G and it is denoted by q . Hence the graph is denoted by $G(p, q)$. If the vertices or edges or both of the graph are assigned values subject to certain conditions it is known as graph labeling. A dynamic survey on graph labeling is regularly updated by Gallian [4] and it is published by Electronic Journal of Combinatorics. Some basic concepts are taken from [5], [6] and [14]. This labeling can also be investigated for the graphs in the papers [7], [8], [9], [10] and [11]. Labeled graphs have variety of applications in graph theory, particularly for missile guidance code, design good radar type codes and convolution codes with optimal autocorrelation properties. Labeled graph plays vital role in the study of X-ray crystallography, communication network and to determine optimal circuit layouts. A detailed study on variety of applications on graph labeling is carried out in Bloom and Golomb [1].

Definition-1: Consider k copies of stars $K_{1,n}^{(1)}, K_{1,n}^{(2)}, K_{1,n}^{(3)}, \dots, K_{1,n}^{(k)}$ then $G = \langle K_{1,n}^{(1)} : K_{1,n}^{(2)} : K_{1,n}^{(3)} : \dots : K_{1,n}^{(k)} \rangle$ is the graph obtained by joining apex vertices of each $K_{1,n}^{(p-1)}$ and $K_{1,n}^{(p)}$ to a new vertex x_{p-1} where $2 \leq p \leq k$. G has $k(2n+2) - 1$ vertices and $k(2n+2) - 2$ edges.

Definition-2: A mapping $f : V(G) \rightarrow \{0, 1\}$ is called binary vertex labeling of G and $f(v)$ is called the label of vertex v of G under f .

The induced edge labeling $f^* : E(G) \rightarrow \{0,1\}$ is given by $f^*(uv) = |f(u) - f(v)|$ for every $uv \in E(G)$. Let $v_f(0)$, $v_f(1)$ be the number of vertices of G having labels 0 and 1 respectively under f and let $e_f(0)$, $e_f(1)$ be the number of edges of G having labels 0 and 1 respectively under f .

Definition-3: A binary vertex labeling of a graph G is called cordial labeling if $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$. A graph G is cordial if it admits cordial labeling.

The concept of cordial labeling was introduced by Cahit [2] in which he investigated several results. The product cordial labeling was investigated by Sundaram [12]. Also some new graphs are investigated as product cordial graphs by Vaidya [13].

Definition-4: A binary vertex labeling of a graph G with induced edge labeling $f^* : E(G) \rightarrow \{0,1\}$ defined by $f^*(uv) = (f(u) + f(v)) \pmod{2}$ is called sum cordial labeling if $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$. A graph G is sum cordial if it admits sum cordial labeling.

Corresponding author: J. SHIAMA*

Assistant professor, Department of mathematics, Cms College of science and commerce,
Coimbatore -641049, tamilnadu, india

2. MAIN RESULTS

Theorem 2.1: The path P_n admits sum cordial labeling.

Proof: Let $G = P_n = v_1 v_2 \dots v_n$ be a path of length n with n vertices and $n-1$ edges

Define $f: V(G) \rightarrow \{0, 1\}$, we consider four cases

Case-1: When $n = 4k$

P_n : $v_1 v_2 \dots v_{4k}$ is a path of length $4k$ with $4k$ vertices and $4k-1$ edges

Define

$$f(v_{4i-3}) = f(v_{4i-2}) = 1, \text{ for } 1 \leq i \leq k \text{ and}$$

$$f(v_{4i-1}) = f(v_{4i}) = 0,$$

Case-2: When $n = 4k - 1$

P_n : $v_1 v_2 \dots v_{4k-1}$ is a path of length $4k-1$ with $4k-1$ vertices and $4k-2$ edges

Define

$$f(v_{4i-3}) = f(v_{4i-2}) = 1$$

$$f(v_{4i-1}) = 0 \text{ for } 1 \leq i \leq k \text{ and } f(v_{4i}) = 0 \text{ for } 1 \leq i \leq k-1$$

Case-3: When $n = 4k + 1$

P_n : $v_1 v_2 \dots v_{4k+1}$ is a path of length $4k+1$ with $4k+1$ vertices and $4k$ edges

Define

$$f(v_{4i-3}) = 1 \text{ for } 1 \leq i \leq k+1$$

$$f(v_{4i-2}) = 1, f(v_{4i-1}) = f(v_{4i}) = 0 \text{ for } 1 \leq i \leq k$$

Case-4: When $n = 4k + 2$

P_n : $v_1 v_2 \dots v_{4k+2}$ is a path of length $4k+2$ with $4k+2$ vertices and $4k+1$ edges

Define

$$f(v_1) = f(v_2) = f(v_5) = 1$$

$$f(v_3) = f(v_4) = f(v_6) = 0$$

$$f(v_{4i-1}) = f(v_{4i}) = 1$$

$$\text{and } f(v_{4i+1}) = f(v_{4i+2}) = 0, \text{ for } 2 \leq i \leq k$$

In all the above cases, $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$

Hence the path P_n admits sum cordial labeling.

Example 2.2: The paths P_7 and P_{10} are sum cordial graphs




Fig-1: Sum cordial labeling of the paths P_7 and P_{10}

Theorem 2.3: The Star $K_{1,n}$ admits sum cordial labeling.

Proof: Let $K_{1,n}$ be a star with $n+1$ vertices and n edges.

Define $f: V(G) \rightarrow \{0, 1\}$, we consider two cases

Case- 1: n is even

Define

$$f(u_i) = 1, \quad 1 \leq i \leq \frac{n}{2}$$

$$= 0, \text{ otherwise}$$

$$\text{and } f(c) = 0$$

Case-2: n is odd

Define'

$$f(u_i) = 1, \quad 1 \leq i \leq \lceil \frac{n}{2} \rceil$$

$$= 0, \text{ otherwise}$$

$$\text{and } f(c) = 0$$

In both the cases

$$|v_f(0) - v_f(1)| \leq 1 \text{ and } |e_f(0) - e_f(1)| \leq 1$$

Hence the star $K_{1,n}$ admits sum cordial labeling.

Example 2.4: The stars $K_{1,6}$ and $K_{1,7}$ are sum cordial graphs




Fig-2: Sum cordial labeling of the stars $K_{1,6}$ and $K_{1,7}$

Theorem 2.5: The bistar $B_{n,n}$ admits sum cordial labeling.

Proof: Let $u, u_1, u_2, \dots, u_n, v, v_1, v_2, \dots, v_n$ be the vertices of the bistar $B_{n,n}$.

Let $G = B_{n,n}$. Then $|V(G)| = 2n+2$ and $|E(G)| = 2n+1$

Define $f: V(G) \rightarrow \{0, 1\}$

Case-i: n is even

Define

$$f(u) = 0$$

$$f(v) = 1$$

$$\text{and } f(u_i) = f(v_i) = 0 \text{ for } 1 \leq i \leq \frac{n}{2}$$

$$= 1 \text{ for } \frac{n}{2} + 1 \leq i \leq n$$

Case-ii: n is odd

Define

$$f(u) = 0$$

$$f(v) = 1$$

$$\text{and } f(u_i) = 0 \text{ for } 1 \leq i \leq \lceil \frac{n}{2} \rceil$$

$$= 1 \text{ for } \lceil \frac{n}{2} \rceil + 1 \leq i \leq n$$

$$\text{and } f(v_i) = 0 \text{ for } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor$$

$$= 1 \text{ for } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n$$

In both the cases $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$

Hence any bistar $B_{n,n}$ admits sum cordial labeling

Example 2.6: The bistars $B_{3,3}$ and $B_{6,6}$ are sum cordial graphs






Fig-3: Sum cordial labeling of the bistars $B_{3,3}$ and $B_{6,6}$

Theorem 2.7: The double star $K_{1,n,n}$ is a sum cordial graph

Proof: Let $u, u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n$ be the vertices of the double star $K_{1,n,n}$.

Let $G = K_{1,n,n}$. Then $|V(G)| = 2n+1$ and $|E(G)| = 2n$

Define $f: V(G) \rightarrow \{0, 1\}$ by

$f(u) = 1$

$f(u_i) = 1, 1 \leq i \leq n$ and

$f(v_i) = 0, 1 \leq i \leq n$

Then $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$

Therefore f is a sum cordial labeling.

Hence any double star $K_{1,n,n}$ admits sum cordial labeling.

Example 2.8: The double stars $K_{1,4,4}$ and $K_{17,7}$ are sum cordial graphs




Fig-4: Sum cordial labeling of the double stars $K_{1,4,4}$ and $K_{17,7}$

Theorem 2.9: Graph $G = \langle K_{1,n,n}^{(1)} : K_{1,n,n}^{(2)} : K_{1,n,n}^{(3)} : \dots : K_{1,n,n}^{(k)} \rangle$ is sum cordial.

Proof: Let $u_1^{(1)}, u_2^{(1)}, u_3^{(1)}, \dots, u_n^{(1)}, v_1^{(1)}, v_2^{(1)}, v_3^{(1)}, \dots, v_n^{(1)}$ be the vertices of $K_{1,n,n}^{(1)}$, $u_1^{(2)}, u_2^{(2)}, u_3^{(2)}, \dots, u_n^{(2)}, v_1^{(2)}, v_2^{(2)}, v_3^{(2)}, \dots, v_n^{(2)}$ be the vertices of $K_{1,n,n}^{(2)}$ and $u_1^{(k)}, u_2^{(k)}, u_3^{(k)}, \dots, u_n^{(k)}, v_1^{(k)}, v_2^{(k)}, v_3^{(k)}, \dots, v_n^{(k)}$ be the vertices of $K_{1,n,n}^{(k)}$. Let c_1, c_2, \dots, c_k be the apex vertices of $K_{1,n,n}^{(1)}, K_{1,n,n}^{(2)}, \dots, K_{1,n,n}^{(k)}$ respectively and c_{p-1} and c_p are adjacent to a new vertex $x_{p-1}, 2 \leq p \leq k$. Let $G = \langle K_{1,n,n}^{(1)} : K_{1,n,n}^{(2)} : K_{1,n,n}^{(3)} : \dots : K_{1,n,n}^{(k)} \rangle$. Here $|V(G)| = 2k(n+1) - 1$ and $|E(G)| = 2k(n+1) - 2$

Define sum cordial labeling $f: V(G) \rightarrow \{0, 1\}$ as follows.

For $1 \leq i \leq n$

$f(u_i^{(p)}) = 1$, if p is odd and $1 \leq p \leq k$

$f(v_i^{(q)}) = 1$, if q is even $1 \leq q \leq n$

$f(v_i^{(p)}) = 0$, if p is odd and $1 \leq p \leq k$

$f(u_i^{(q)}) = 0$, if q is even and $1 \leq p \leq k$

$f(c_i) = 1$; if i is odd, $1 \leq i \leq k$

$f(c_i) = 0$; if i is even, $1 \leq i \leq k$

$f(x_{p-1}) = 1$; if p is even and $2 \leq p \leq k$

$f(x_{p-1}) = 0$; if p is odd and $2 \leq p \leq k$

Also $|v_f(0) - v_f(1)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$

Hence $G = \langle K_{1,n,n}^{(1)} : K_{1,n,n}^{(2)} : K_{1,n,n}^{(3)} : \dots : K_{1,n,n}^{(k)} \rangle$ admits sum cordial labeling.

Example 2.10: $G = \langle K_{1,7,7}^{(1)}, K_{1,7,7}^{(2)}, K_{1,7,7}^{(3)} \rangle$ is a sum cordial.




Fig-5: Sum cordial labeling of the graph $G = \langle K_{1,7,7}^{(1)}, K_{1,7,7}^{(2)}, K_{1,7,7}^{(3)} \rangle$

Example 2.10: $G = \langle K_{1,6,6}^{(1)}, K_{1,6,6}^{(2)} \rangle$ is a sum cordial.




Fig-6: Sum cordial labeling of the graph $G = \langle K_{1,6,6}^{(1)}, K_{1,6,6}^{(2)} \rangle$

CONCLUSION

In this paper some graphs like path, star, bistar, doublestar and the graphs obtained by joining apex vertices of k copies of double stars to a new vertex are investigated for the sum cordial labeling. This labeling can be verified for some other graphs.

REFERENCES

- [1] G.S Bloom and S.W Golomb "Applications of numbered undirected graph "Proceedings of IEEE, 65(4), pp.562-570. 1977.
- [2] Cahit I," Cordial Graphs: A weaker version of graceful and harmonious graphs" Ars Combinatoria vol.23, pp.201-207, 1987.
- [3] Frank Harry, Graph theory, Narosa Publishing House- (2001).
- [4] J A Gallian, A dynamic survey of graph labeling, The Electronics journal of Combinatorics, 17(2010) # DS6.
- [5] Gary Chartrand, and Lesniak, Graphs and digraphs, fourth edition, Chapman & HALL/CRC.
- [6] Gary Chartrand, Ping Zhang, Introduction to Graph theory, McGraw- Hill International Edition.
- [7] J. Shiama" Square sum labeling for some middle and total graphs" International Journal of Computer Applications (0975-08887) Volume 37- No.4 January 2012.

- [8] J. Shiama "Permutation labeling for some shadow graphs" International Journal of Computer Application (0975-8887) Volume 40- No.6 February 2012.
- [9] J. Shiama "Permutation labeling for some splitting graphs" Proceedings of the National Conference on Mathematical Modeling and Simulation. NCMS'12 11th Feb.2012, by Vinayaka missions.
- [10] J. Shiama" Square difference labeling for some graphs" International Journal of Computer Applications (0975-08887) Volume 44- No.4, April 2012.
- [11] J. Shiama "Some Special types of Square difference graphs" International Journal of Mathematical archives- 3(6), 2012,2369-2374 ISSN 2229-5046.
- [12] Sundaram M,Ponraj and Somasundaram S "Product cordial labeling of graphs". Bull. Pure and Applied Science(Mathematics and Statistics) ,Vol.23E, pp. 155-163,2004.
- [13] Vaidya and Dani N.A "Some new product cordial graphs", journal of App. Comp. Sci. Math. Vol8, No.4, pp.62-65, 2010
- [14] D B West, Introduction to Graph Theory, Prentice-Hall, India, 2001.

Source of support: Nil, Conflict of interest: None Declared