

## COMMUTANT OF COMPOSITE INTEGRAL OPERATORS

Anupama Gupta\*

Govt. College for Women, Parade. Jammu, J&K, India

(Received on: 29-09-12; Revised & Accepted on: 10-11-12)

### ABSTRACT

*In this paper we study the composite integral operators on  $L^p$ -spaces. The conditions for composite integral operators to be bounded are investigated. The commutants of composite integral operators and Volterra composition operators are computed.*

**Keywords:** Randon-Nikodym derivative, conditional expectation operator, Commutant, Contraction, Fixed point

**AMS Subject Classification (2005):** Primary 47B38, Secondary 47B399.

### 1. INTRODUCTION AND PRELIMINARIES

Let  $(X, S, \mu)$  be a  $\sigma$ -finite measure space and let  $\phi: X \rightarrow X$  be a non-singular measurable transformation ( $\mu(E) = 0 \Rightarrow \mu\phi^{-1}(E) = 0$ ). Then a composition transformation  $C_\phi: L^p(\mu) \rightarrow L^p(\mu)$  is defined by the equation

$$C_\phi f = f \circ \phi \text{ for every } f \in L^p(\mu).$$

In case  $C_\phi$  is continuous, we call it a composition operator induced by  $\phi$ .

For each  $f \in L^p(\mu)$ ,  $1 \leq p < \infty$ , there exists a unique  $\phi^{-1}(S)$  measurable function  $E(f)$  such that

$$\int g f d\mu = \int g E(f) d\mu$$

for every  $\phi^{-1}(S)$  measurable function  $g$  for which the left integral exists. The function  $E(f)$  is called conditional expectation of  $f$  with respect to the sub  $\sigma$ - algebra  $\phi^{-1}(S)$ .  $E$  has the property that for  $f \in L^p(\mu)$ ,  $E(f) = g \circ \phi$  for exactly one  $S$ -measurable function  $g$ . We shall write  $g = E(f) \circ \phi^{-1}$ , which is well-defined measurable function. For more details about expectation operator, we refer to Parathasarthy [8].

Let  $K: X \times X \rightarrow \mathbb{C}$  be a measurable function. Then a linear transformation  $I: L^p(\mu) \rightarrow L^p(\mu)$  defined by

$$(I f)(x) = \int K(x, y) f(y) d\mu(y) \quad \text{for all } f \in L^p(\mu)$$

is known as integral operator. The composite integral operator  $I_\phi$  is a bounded linear operator  $I_\phi: L^p(\mu) \rightarrow L^p(\mu)$  defined by

$$(I_\phi f)(x) = \int K(x, y) f(\phi(y)) d\mu(y) \tag{1}$$

The equation (1) can also be written as

$$(I_\phi f)(x) = \int E(K_x) \circ \phi^{-1}(y) f_o(y) d\mu(y),$$

where  $K_x(y) = K(x, y)$  and  $f_o = \frac{d\mu\phi^{-1}}{d\mu}$ , the Randon-Nikodym derivative of the measure  $\mu\phi^{-1}$  with respect to the measure  $\mu$ .

**Corresponding author:** Anupama Gupta\*, Govt. College for Women, Parade. Jammu, J&K, India

The Volterra composition operator is a composition of Volterra integral operator  $V$  and a composition operator  $C_\phi$  defined as

$$\begin{aligned}(V_\phi f)(x) &= (V f) \circ \phi(x) \\ &= \int_0^x f(\phi(t)) dt \quad \text{for every } f \in L^p[0,1],\end{aligned}$$

where  $\phi : [0,1] \rightarrow [0,1]$  is a measurable function.

An intensive study of composition operators is made over the past several decades. To worth mention, few of them are Singh ([10], [11]), Singh and Kumar [12], Singh and Komal [13], Singh and Manhas [14], Ridge [9] and Campbell [2]. The integral operators and composite integral operators in particular Volterra integral operator on  $L^p(\mu)$  have received considerable attention in recent years. The theory of integral operators is the source of all modern functional analysis and operator theory. Mathematicians like Halmos and Sunder [6], Setpanov ([15], [16]), Bloom and Kermen [1] have done great deal of work on integral operators. Gupta and Komal ([3], [4], [5]) also studied composite integral operators. Whitley [17] established the Lyubic's conjecture [7] and generalized it to Volterra composition operators on  $L^p[0,1]$ .

In this paper we have make an effort to explore a commutant of composite integral operators.

## 2. BOUNDED COMPOSITE INTEGRAL OPERATORS

In this section we study bounded composite integral operators.

**Theorem 2.1:** Suppose  $1 \leq p, q < \infty$ . Suppose  $I_\phi : L^p(\mu) \rightarrow L^q(\mu)$  is a linear transformation. Then  $I_\phi$  is continuous.

**Proof:** Let  $f_n \rightarrow f$  in  $L^p(\mu)$  and  $I_\phi f_n \rightarrow g$  in  $L^q(\mu)$ . Then there exists a dominated subsequence  $\{f'_n\}$  of  $\{f_n\}$  such that

$$f'_n(x) \rightarrow f(x) \quad \text{a.e.} \quad (i)$$

Again since  $I_\phi f_n \rightarrow g$  in  $L^q(\mu)$ , we can select a dominated subsequence  $\{f''_n\}$  of  $\{f'_n\}$  such that

$$(I_\phi f''_n)(x) \rightarrow g(x) \quad \text{a.e.}$$

or that

$$\int K_\phi(x, y) f''_n(y) d\mu(y) \rightarrow g(x) \quad \text{a.e.}$$

Also  $|f''_n| \leq h$  for some  $h \in L^p(\mu)$ .

It follows from (i) that

$$K_\phi(x, y) f''_n(y) \rightarrow K_\phi(x, y) f(y) \quad \text{a.e.} \quad (ii)$$

and

$$|K_\phi(x, y) f''_n(y)| \leq |K_\phi(x, y) h(y)| \quad \text{for almost every } y.$$

But the dominated subsequence  $\{K_\phi f''_n\}$  converges to  $\{K_\phi f\}$  almost everywhere. By the Lebesgue dominated convergence theorem,

$$\int K_\phi(x, y) f''_n(y) d\mu(y) \rightarrow \int K_\phi(x, y) f(y) d\mu(y) \quad (iii)$$

From (ii) and (iii), we conclude that

$$(I_\phi f)(x) = \int K_\phi(x, y) f(y) d\mu(y) = g(x)$$

which proves that the graph of  $I_\phi$  is closed. Hence, by the closed graph theorem,  $I_\phi$  is continuous.

In the following theorem, we take  $r$  such that and  $\frac{1}{p} + \frac{1}{r} = \frac{1}{q}$  and

$$S(x) = \|K_\phi(x, \cdot)\|_{r/q}$$

**Theorem 2.2:** For  $1 \leq p, q < \infty$ , let  $S \in L^{r/q}(\mu)$ . Then  $I_\phi: L^p(\mu) \rightarrow L^q(\mu)$  is a bounded composite integral operator.

**Proof:** For  $f \in L^p(\mu)$ , consider

$$\begin{aligned} \|I_\phi f\|^q &= \int_X |I_\phi f|^q dx \\ &= \int_X \left| \int_X K_\phi(x, y) f(y) dy \right|^q dx \\ &\leq \int_X \left\{ \left( \int_X |K_\phi(x, y)|^{r/q} dy \right)^{q/r} \left( \int_X |f(y)|^{p/q} dy \right)^{q/p} \right\} dx \\ &= \|S(x)\|_{r/q}^q \cdot \|f\|^p \end{aligned}$$

This proves that  $I_\phi$  is bounded composite integral operator.

In the next result we make an attempt to use composite integral operators to solve the integral equations.

**Theorem 2.3:** If  $K_\phi \in L^2(\mu \times \mu)$  and  $g \in L^2[0, 1]$ , then the integral equation

$$f(x) = g(x) + \lambda \int K_\phi(x, y) f(y) d\mu(y) \quad (1)$$

has unique solution for sufficiently small values of scalar  $\lambda$ .

**Proof:** Define  $I_\phi: L^2[0, 1] \rightarrow L^2[0, 1]$  as  $I_\phi f = h$

$$\text{where } h(x) = g(x) + \lambda \int_0^1 K_\phi(x, y) f(y) d\mu(y).$$

We first show that

$$\psi(x) = \int K_\phi(x, y) f(y) d\mu(y) \quad \text{for every } f \in L^2[0, 1].$$

Consider

$$\left| \int_0^1 K_\phi(x, y) f(y) d\mu(y) \right| \leq \left( \int_0^1 |K_\phi(x, y)|^2 d\mu(y) \right)^{1/2} \left( \int_0^1 |f(y)|^2 d\mu(y) \right)^{1/2} \quad (\text{by using Holder's inequality})$$

Therefore,

$$\begin{aligned} \int_0^1 |\psi(x)|^2 dx &\leq \int_0^1 \left( \int_0^1 |K_\phi(x, y)|^2 d\mu(y) \right) d\mu(x) \int_0^1 \left( \int_0^1 |f(y)|^2 d\mu(y) \right) d\mu(x) \\ &< \infty. \end{aligned}$$

Now

$$\begin{aligned} \|I_\phi f - I_\phi f_1\| &= \|\lambda \int_0^1 K_\phi(x, y) [f(y) - f_1(y)] d\mu(y)\| \\ &\leq |\lambda| \left( \int_0^1 \int_0^1 |K_\phi(x, y)|^2 d\mu(x) d\mu(y) \right)^{1/2} \left( \int_0^1 |f(y) - f_1(y)|^2 d\mu(y) \right)^{1/2} \\ &\leq M \|f - f_1\|, \end{aligned}$$

$$\text{where } M = |\lambda| \left( \int_0^1 \int_0^1 |K_\phi(x, y)|^2 d\mu(x) d\mu(y) \right)^{1/2}.$$

This proves that  $I_\phi$  is a contraction and hence it has a unique fixed point, say  $f^*$ . Thus  $f^*$  is a unique solution of eq. (1).

### 3. COMMUTANT OF COMPOSITE INTEGRAL OPERATOR

In this section we have made an attempt to compute the commutant of composite integral operator.

**Theorem 3.1:** Let  $I_\phi \in B(L^p(\mu))$ . Then  $M_\theta$  commutes with  $I_\phi$  if and only if  $\theta = \theta \circ \phi$  a.e.

**Proof:** For  $f \in L^p(\mu)$ ,

$$\begin{aligned}(I_\phi M_\theta f)(x) &= \int K(x, y)(M_\theta f) \circ \phi(y) \, d\mu(y) \\ &= \int E(K_x \circ \phi^{-1})(y) f_\theta(y) \theta(y) f(y) \, d\mu(y)\end{aligned}\tag{i}$$

and

$$\begin{aligned}(M_\theta I_\phi f)(x) &= \theta(x) (I_\phi f)(x) \\ &= \theta(x) \int E(K_x \circ \phi^{-1})(y) f_\theta(y) f(y) d\mu(y)\end{aligned}\tag{ii}$$

In view of (i) and (ii)

$$(M_\theta I_\phi f)(x) - (I_\phi M_\theta f)(x) = \int f_\theta(y) E(K_x \circ \phi^{-1})(y) [\theta(y) - \theta(x)] f(y) d\mu(y).$$

Hence, the result.

In the next theorem we characterize multiplication operators which commute with Volterra composite operators.

**Theorem 3.2:** Let  $M_\theta \in B(L^2(\mu))$ . Suppose  $\phi$  is an injective map. Then  $M_\theta$  commutes with  $V_\phi$  if and only if  $\theta = \theta \circ \phi$  a.e.

**Proof:** For  $f \in L^2(\mu)$ , we have

$$\begin{aligned}(M_\theta V_\phi f)(x) &= (\theta \cdot V(f \circ \phi))(x) \\ &= \theta(x) \int_0^x f(\phi(t)) dt \\ &= \theta(x) \int_0^x \chi_{[0,x]}(t) f(\phi(t)) dt \\ &= \theta(x), \quad \text{for } f = \chi_{[0,x]}\end{aligned}$$

Also we have

$$\begin{aligned}(V_\phi M_\theta f)(x) &= V(M_\theta f) \circ \phi(x) \\ &= \int_0^x \theta \circ \phi \cdot f \circ \phi(t) dt \\ &= \int_0^1 \chi_{[0,x]}(t) \theta \circ \phi(t) f(\phi(t)) dt \\ (M_\theta V_\phi f)(x) - (V_\phi M_\theta f)(x) &= \int_0^1 \chi_{[0,x]}(t) [\theta(x) - \theta \circ \phi(t)] f(\phi(t)) dt\end{aligned}$$

as  $\phi$  is injective,  $C_\phi$  has dense range

$$\chi_{[0,x]}(t) [\theta(x) - \theta \circ \phi(t)] = 0.$$

Hence the result follows using the given condition.

**Corollary:** There is a composition operator  $C_\phi \in L^2(\mu)$  such that  $V C_\phi = C_\phi V$

**Proof:** For  $f \in L^2(\mu)$ , we have

$$\begin{aligned}(V C_\phi f)(x) &= \int_0^x C_\phi f(t) dt = \int_0^x f \circ \phi(t) dt \\ (C_\phi V f)(x) &= (V f) \circ \phi(x) = \int_0^{\phi(x)} f(t) dt = \int_0^x f \circ \phi(t) dt\end{aligned}$$

Hence, the result follows.

## REFERENCES

- [1] Bloom, S. and Kerman, R., Weighted norm inequalities for operators of Hardy type, Proc. Amer. Math. Soc. 113 (1991), 135-141.
- [2] Campbell, J. and Jamison, J., On some classes of weighted composition operators, Glasgow Math. J. 32 (1990), 87-94.
- [3] Gupta, A. and Komal, B.S., Composite integral operators on  $L^2(\mu)$ , Pitman Lecture Notes in Mathematics series 377 (1997), 92-99.
- [4] Gupta, A. and Komal, B.S., Weighted Composite Integral Operators, Int. Journal of Math Analysis, Vol.3, 26 (2009), 1283-1293.
- [5] Gupta, A. and Komal, B.S., Volterra composition operators, Int. J. Contemp. Math. Sciences, vol.6, no.7 (2011), 345-351.
- [6] Halmos, P. R. and Sunder, V.S., Bounded integral operators on  $L^2$ -spaces, Springer-Verlag, New York, (1978).
- [7] Lyubich, Yu I., Composition of integration and substitution, Linear and Complex Analysis, Problem Book, Springer Lect. Notes in Math., 1043, Berlin, (1984), 249-250.
- [8] Parthasarthy, K.R., Introduction to probability and measure, MacMillan Limited, (1977).
- [9] Ridge, W.C., Spectrum of composition operators, Proc. Amer. Math. Soc. 37 (1973), 121-127.
- [10] Singh, R.K., Compact and quasinormal composition operators, Proc. Amer. Math. Soc. 45(1974), 80-82
- [11] Singh, R.K., Normal and Hermitian composition operators, Proc. Amer. Math. Soc. 47 (1975), 348-350
- [12] Singh, R.K. and Komal, B.S., Composition operators, Bull. Austral. Math. Soc. 18 (1978), 439-446.
- [13] Singh, R.K., and Kumar, A., Multiplication operators and composition operators with closed ranges, Bull. Austral. Math. Soc. 16 (1977), 247-252.
- [14] Singh, R.K., and Manhas, J.S., Composition operators on function spaces, North Holland Mathematics studies 179, Elsevier Science Publishers Amsterdam, (1993).
- [15] Stepanov, V. D., Weighted inequalities for a class of Volterra convolution operators, J. London Math, (1992), 232-242.
- [16] Stepanov, V. D., Weighted norm inequalities of Hardy type for a class of integral operators, J. London Math. Soc. (2) 50 (1994), 105-120.
- [17] Whitley, R., The spectrum of a Volterra composition operator, Integral equation and operator theory, Vol. 10 (1987), 146-149.

**Source of support: Nil, Conflict of interest: None Declared**