

FORMATION OF TWO SUMMATION FORMULAE ALLIED WITH CONTIGUOUS RELATION

SALAHUDDIN*

P.D.M College of Engineering, Bahadurgarh, Haryana, India

Emails: sludn@yahoo.com ; vsludn@gmail.com

(Received on: 04-03-11; Accepted on: 21-03-11)

ABSTRACT

The main aim of this paper is to evaluate two summation formulae involving Contiguous Relation associated with Recurrence relation and Hypergeometric function.

A.M.S. Subject Classification (2000): 33C05, 33C45, 33C60, 33C70

Key words and phrases: Pochhammer's symbol, Contiguous relation, Recurrence relation

A. INTRODUCTION:

The Pochhammer's symbol:

$$(a, k) = (a)_k = \frac{\Gamma(a+k)}{\Gamma(a)} = \begin{cases} \alpha(\alpha+1)(\alpha+2) \dots (\alpha+k-1); & \text{if } k = 1, 2, 3, \dots \\ 1; & \text{if } k = 0 \\ ; & \text{if } \alpha = 1 \end{cases} \quad (1)$$

Generalized Gaussian Hypergeometric function of one variable:

$${}_A F_B(a_1, a_2, \dots, a_A; b_1, b_2, \dots, b_B; z) = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_A)_k z^k}{(b_1)_k (b_2)_k \dots (b_B)_k k!} \quad (2)$$

or

$${}_A F_B((a_A); (b_B); z) \equiv {}_A F_B((a_j)_{j=1}^A; (b_j)_{j=1}^B; z) = \sum_{k=0}^{\infty} \frac{((a_A))_k z^k}{((b_B))_k k!} \quad (3)$$

where the parameters b_1, b_2, \dots, b_B are neither zero nor negative integers and A, B are non negative integers.

Contiguous Relations:

[Andrews p.363 (9.16), E.D. p.51 (10), H.T.F.I. p.103 (32)]

$$(a-b) {}_2 F_1(a, b; c; z) = a {}_2 F_1(a+1, b; c; z) - b {}_2 F_1(a, b+1; c; z) \quad (4)$$

[Abramowitz p.558 (15.2.19)]

$$(a-b)(1-z) {}_2 F_1(a, b; c; z) = (c-b) {}_2 F_1(a, b-1; c; z) + (a-c) {}_2 F_1(a-1, b; c; z) \quad (5)$$

A New Summation Formula:

$${}_2 F_1(a, b; \frac{a+b-1}{2}; \frac{1}{2}) = 2^{b-1} \frac{\Gamma(\frac{a+b-1}{2})}{\Gamma(b)} \left[\frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-1}{2})} \left\{ \frac{b+a-1}{a-1} \right\} + 2 \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a}{2})} \right] \quad (6)$$

Recurrence relation:

$$\Gamma(z+1) = z \Gamma(z) \quad (7)$$

***Corresponding author: SALAHUDDIN*, *E-mail: sludn@yahoo.com**

B. MAIN SUMMATION FORMULAE:

For both the formulae $a \neq b$

For $a < 1$ and $a > 13$

$$\begin{aligned}
 {}_2F_1(a, b ; \frac{a+b-13}{2}; \frac{1}{2}) = & 2^{b-1} \frac{\Gamma(\frac{a+b-13}{2})}{(a-b)\Gamma(b)} \left[\frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-13}{2})} \left\{ \frac{-135135a + 264207a^2 - 177331a^3 + 57379a^4 - 10045a^5}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \right. \\
 & + \frac{973a^6 - 49a^7 + a^8 + 135135b - 318983a^2b + 298522a^3b - 36415a^4b + 18956a^5b - 1225a^6b + 90a^7b - 264207b^2}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{318983ab^2 - 93730a^3b^2 + 54145a^4b^2 - 5733a^5b^2 + 910a^6b^2 + 177331b^3 - 298522ab^3 + 93730a^2b^3 - 5005a^4b^3}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} + \\
 & + \frac{2002a^5b^3 - 5733a^6b^3 + 86415ab^4 - 54145a^2b^4 + 5005a^5b^4 + 10045ab^5 - 18956a^2b^5 + 5733a^3b^5 - 2002a^4b^5}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{-973b^6 + 1225ab^6 - 910a^3b^6 + 49b^7 - 90ab^7 - b^8}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-1)} \left. \right\} + \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-12}{2})} \left\{ \frac{221874a - 249056a^2 + 141866a^3 - 27200a^4}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \\
 & + \frac{4886a^5 - 224a^6 + 14a^7 - 221874b + 224578a^2b - 101760a^3b + 37650a^4b - 2680a^5b + 350a^6b + 249056b^2}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & + \frac{224578ab^2 + 39260a^3b^2 - 6240a^4b^2 + 1638a^5b^2 - 141866b^3 + 101760ab^3 - 39260a^2b^3 + 1430a^4b^3 + 27200b^4}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & \left. \left. \left. + \frac{-37650ab^4 + 6240a^2b^4 - 1430a^3b^4 - 4886b^5 + 2688ab^5 - 1638a^2b^5 + 224b^6 - 350ab^6 - 14b^7}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right] \right\} \quad (8)
 \end{aligned}$$

For $a < 1$ and $a > 14$

$$\begin{aligned}
 {}_2F_1(a, b ; \frac{a+b-14}{2}; \frac{1}{2}) = & 2^{b-1} \frac{\Gamma(\frac{a+b-14}{2})}{(a-b)\Gamma(b)} \left[\frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-14}{2})} \left\{ \frac{-645120a + 1326336a^2 - 833920a^3}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \right. \\
 & + \frac{330480a^4 - 49600a^5 + 7224a^6 - 280a^7 + 16a^8 + 645120b - 489984ab - 721792a^2b + 887360a^3b - 258240a^4b}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{70256a^5b - 4088a^6b + 440a^7b - 836352b^2 + 1135488ab^2 - 351323a^2b^2 - 141440a^3b^2 + 117000a^4b^2 - 12792a^5b^2}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{2548a^6b^2 + 420224b^3 - 758208ab^3 + 268160a^2b^3 - 47849a^3b^3 - 5720a^4b^3 + 3432a^5b^3 - 108304b^4 + 165440ab^4}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{-115000a^2b^4 + 11960a^3b^4 - 1430a^4b^4 + 15680b^5 - 30352ab^5 + 9240a^2b^5 - 3640a^3b^5 - 1288b^6 + 1624ab^6}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{-1260a^2b^6 + 56b^7 - 104ab^7 - b^8}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \left. \right\} + \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-14}{2})} \left\{ \frac{-645120a + 836352a^2 - 420224a^3 + 108304a^4}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-2)} \right. \\
 & + \frac{-15680a^5 + 1288a^6 - 56a^7 + a^8 + 645120b + 489984ab - 1135488a^2b + 758208a^3b - 165440a^4b + 30352a^5b}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & + \frac{-1624a^6b + 104a^7b - 1326336b^2 + 721792ab^2 + 351323a^2b^2 - 268160a^3b^2 + 115000a^4b^2 - 7248a^5b^2}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & + \frac{1260a^6b^2 + 833920b^3 - 887360ab^3 + 141440a^2b^3 + 47840a^3b^3 - 11960a^4b^3 + 3640a^5b^3 - 230480b^4}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & + \frac{258240ab^4 - 117000a^2b^4 + 5720a^3b^4 + 1430a^4b^4 + 49600b^5 - 70256ab^5 + 12792a^2b^5 - 3432a^3b^5}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)}
 \end{aligned}$$

$$+\frac{-7224b^6+4088ab^6-2548a^2b^6+280b^7-440ab^7-15b^8}{(a-14)(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)}\}] \quad (9)$$

C. DERIVATIONS OF SUMMATION FORMULAE (8) TO (9):

Derivation of (7): Substituting $c = \frac{a+b-13}{2}$ and $z = \frac{1}{2}$ in equation (4), we get

$$(a-b) {}_2F_1(a, b; \frac{a+b-13}{2}; \frac{1}{2}) = (a-b-13) {}_2F_1(a, b-1; \frac{a+b-13}{2}; \frac{1}{2}) + (a-b+13) {}_2F_1(a-1, b; \frac{a+b-13}{2}; \frac{1}{2})$$

Now with the help of the derived result from equation (6), we get

$$\begin{aligned} L.H.S &= (a-b-13) 2^{b-2} \frac{\Gamma(\frac{a+b-13}{2})}{(a-b+1)\Gamma(b-1)} \left[\frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-11}{2})} \left\{ \frac{135135 - 118677a - 96005a^2 + 112567a^3 - 3875a^4}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \right. \\ &\quad \frac{6097a^5 - 455a^6 + 13a^7 - 264207b + 363186ab - 58929a^2b - 56004a^3b + 27183a^4b - 3822a^5b + 273a^6b}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \\ &\quad \left. \left. + \frac{177331b^2 - 274531ab^2 + 104702a^2b^2 + 858a^3b^2 - 4433a^4b^2 + 1001a^5b^2 - 57379b^3 + 91292ab^3 - 40274a^2b^3}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \right\} \right. \\ &\quad \left. + \frac{5148a^3b^3 + 429a^4b^3 + 10045b^4 - 16445ab^4 + 6279a^2b^4 - 1001a^5b^4 - 973b^5 + 1330ab^5 - 637a^2b^5 + 49b^6}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \\ &\quad \left. + \frac{-77ab^2 - b^7}{(a-11)(a-9)(a-7)(a-5)(a-3)(a-1)} + \frac{\Gamma(\frac{b-1}{2})}{\Gamma(\frac{a-12}{2})} \left\{ \frac{-211479 - 41633a - 14635b^2/a^2 + 85507/a^3 - 20923a^4}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \right. \\ &\quad \left. \left. + \frac{2205a^6 - 119a^7 + a^8 - 480059b + 292450ab + 69059a^2b - 82052a^3b + 20139a^4b - 2142a^5b + 77a^6b + 398307b^2}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right\} \right. \\ &\quad \left. + \frac{-320047ab^2 + 44590a^2b^2 + 22386a^3b^2 - 6097a^4b^2 + 637a^5b^2 - 160407b^3 + 144924ab^3 - 32890a^2b^3 + 572a^3b^3}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \\ &\quad \left. + \frac{1001a^4b^5 + 35061b^4 - 32799ab^4 + 8151a^2b^4 - 429a^3b^4 - 4641b^5 + 3458ab^5 - 1001a^2b^5 + 273b^6 - 273ab^6 - 13b^7}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right\} \right] \\ &\quad + (a-b+13) 2^{b-1} \frac{\Gamma(\frac{a-b-13}{2})}{(a-b-1)\Gamma(b)} \left[\frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-12}{2})} \left\{ \frac{-211479 + 480059a - 398307a^2 + 160407a^3 - 35061a^4 + 4641a^5}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \right. \\ &\quad \left. \left. + -273a^6 + 13a^7 + 41633b - 292450ab + 320047a^2b - 144924a^3b + 32799a^4b - 3458a^5b + 273a^6b + 144857b^2}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right\} \right. \\ &\quad \left. + \frac{-69059ab^2 - 44590a^2b^2 + 32890a^3b^2 - 8151a^4b^2 + 1001a^5b^2 - 85507b^3 + 82052ab^3 - 22386a^2b^3 - 572a^3b^3}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \\ &\quad \left. + \frac{429a^4b^5 + 20923b^4 - 20139ab^4 + 6097a^2b^4 - 1001a^3b^4 - 2205b^5 + 2142ab^5 - 637a^2b^5 + 119b^6 - 77ab^6 - b^7}{(a-12)(a-10)(a-8)(a-6)(a-4)(a-2)} \right\} \\ &\quad + \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-13}{2})} \left\{ \frac{-135135 + 264207a - 177331a^2 + 57379a^3 - 10045a^4 + 973a^5 - 49a^6 + a^7 + 118677b - 363186ab}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)} \right. \\ &\quad \left. + \frac{274531a^2b - 91292a^3b + 16443a^4b - 1330a^5b + 77a^6b + 96005b^2 + 58929ab^2 - 104702a^2b^2 + 40274a^3b^2}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)} \right\} \end{aligned}$$

$$+\frac{-6279a^4b^2+637a^5b^2-112567b^5+56004ab^5-859a^5b^2-5149a^6b^2+1001a^6b^2+38675b^4-27189ab^4}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)}$$

$$+\frac{4433a^2b^4-429a^3b^4-6097b^5+3822ab^5-1001a^2b^5+455b^6-273ab^6-13b^7}{(a-13)(a-11)(a-9)(a-7)(a-5)(a-3)} \}$$

On simplification, we get the result (8)

Similarly, we can prove the result (9).

REFERENCES:

- [1] Abramowitz, Milton., A and Stegun, Irene ; *Handbook of Mathematical Functions with Formulas , Graphs , and Mathematical Tables*. National Bureau of Standards, 1970.
- [2] Arora, Asish, Singh, Rahul , Salahuddin ; Development of a family of summation formulae of half argument using Gauss and Bailey theorems , *Journal of Rajasthan academy of Physical Sciences* .,7(2008) , 335-342
- [3] Lavoie, J.L., ; Some summation formulae for the series ${}_3F_2$, *Math. Comput.* , 49(1987),269-274
- [4] Lavoie, J.L., Grondin, F. and Rathie,A.K. ; Generalizations of Watson's theorem on the sum of a ${}_3F_2$,*Indian J. Math.*,34(1992),23-32
- [5] Lavoie, J. L., Grondin, F. and Rathie, A.K.; Generalizations of Whipple's theorem on the sum of a ${}_3F_2$, *J. Comput. Appl. Math.*, 72(1996), 293-300.
- [6] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O.I.; *Integrals and Series Vol.3: More Special Functions*. Nauka, Moscow, 1986. Translated from the Russianby G.G. Gould, Gordon and Breach Science Publishers, New York, Philadelphia, London, Paris, Montreux, Tokyo, Melbourne, 1990.
- [7] Rainville, E. D.; The contiguous function relations for ${}_pF_q$ with applications to Bateman's $J_n^{u,v}$ and Rice's $H_n(\zeta, p, v)$, *Bull. Amer. Math. Soc.*, 51(1945), 714-723.
