

## THREE SUMMATION FORMULAE USING CONTIGUOUS RELATION AND RELATED TO HYPERGEOMETRIC FUNCTION

SALAHUDDIN\*

*P.D.M College of Engineering, Bahadurgarh , Haryana , India**Emails : [sludn@yahoo.com](mailto:sludn@yahoo.com) ; [vsludn@gmail.com](mailto:vsludn@gmail.com)**(Received on: 11-04-11; Accepted on: 17-04-11)***ABSTRACT**

*The main aim of this paper is to evaluate two summation formulae involving Contiguous Relation associated with Recurrence relation and Hypergeometric function.*

**A.M.S. Subject Classification (2000) :** 33C05 , 33C20 , 33C45 , 33C60 , 33C70

**Key words and phrases:** Contiguous relation , Gaussian Hypergeometric function.

**A. INTRODUCTION:****Generalized Gaussian Hypergeometric function of one variable**

$${}_A F_B(a_1, a_2, \dots, a_A; b_1, b_2, \dots, b_B; z) = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_A)_k}{(b_1)_k (b_2)_k \dots (b_B)_k k!} z^k \quad (1)$$

where the parameters  $b_1, b_2, \dots, b_B$  are neither zero nor negative integers and  $A, B$  are non negative integers.

**Contiguous Relations**

[Andrews p.363(9.16) , E.D. p.51(10), H.T.F.I. p.103(32)]

$$(a-b) {}_2F_1(a, b; c; z) = a {}_2F_1(a+1, b; c; z) - b {}_2F_1(a, b+1; c; z) \quad (2)$$

[Abramowitz p.558 (15.2.19)]

$$(a-b)(1-z) {}_2F_1(a, b; c; z) = (c-b) {}_2F_1(a, b-1; c; z) + (a-c) {}_2F_1(a-1, b; c; z) \quad (3)$$

**A New Summation Formula**

[Ref. [2] p.337 (10)]

$${}_2F_1(a, b; \frac{a+b-1}{2}; \frac{1}{2}) = 2^{b-1} \frac{\Gamma(\frac{a+b-1}{2})}{\Gamma(b)} \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-1}{2})} \left\{ \frac{b+a-1}{a-1} \right\} + 2 \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{1}{2})} \right] \quad (4)$$

**B. MAIN SUMMATION FORMULAE:**

For all the formulae  $a \neq b$

For  $a < 1$  and  $a > 8$

$$\begin{aligned} {}_2F_1(a, b; \frac{a+b-8}{2}; \frac{1}{2}) = & 2^{b-1} \frac{\Gamma(\frac{a+b-8}{2})}{(a-b)\Gamma(b)} \left[ \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-7}{2})} \left\{ \frac{384a - 624a^2 + 396a^3 - 60a^4 + 9a^5 - 384b + 224ab + 348a^2b}{(a-7)(a-5)(a-3)(a-1)} \right. \right. + \\ & \left. \left. \frac{-192a^5b + 75a^4b + 400b^2 - 604ab^2 + 72a^3b^2 + 42a^2b^3 - 140b^4 + 160ab^3 - 90a^3b^3 + 20b^4 - 35ab^4 - b^5}{(a-7)(a-5)(a-3)(a-1)} \right\} + \right] \end{aligned}$$

**\*Corresponding author: SALAHUDDIN\*, \*E-mail: [sludn@yahoo.com](mailto:sludn@yahoo.com)**

$$\begin{aligned}
 & + \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-8}{2})} \left\{ \frac{384a - 400a^2 + 140a^3 - 20a^4 + a^5 - 334b - 224ab + 604a^2b - 160a^3b + 35a^4b + 624b^2 - 348ab^2 - 72a^2b^2}{(a-8)(a-6)(a-4)(a-2)} + \right. \\
 & \left. + \frac{90a^5b^2 - 396b^5 + 192ab^5 - 42a^2b^3 + 60b^4 - 75ab^4 - 9b^6}{(a-8)(a-6)(a-4)(a-2)} \right\} ] \quad (5)
 \end{aligned}$$

For  $a < 1$  and  $a > 9$

$$\begin{aligned}
 {}_2F_1(a, b ; \frac{a+b-9}{2}; \frac{1}{2}) = & 2^{b-1} \frac{\Gamma(\frac{a+b-9}{2})}{(a-b)\Gamma(b)} \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-9}{2})} \left\{ \frac{-945a + 1689a^2 - 950a^3 + 230a^4 - 25a^5 + a^6 + 945b - 1870a^2b}{(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \\
 & + \frac{1540a^5b - 275a^4b + 44a^5b - 1689b^2 + 1870ab^2 - 330a^5b^2 + 165a^4b^2 + 950b^5 - 1640ab^3 + 330a^2b^3 - 230b^4 + 275ab^4}{(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{-165a^5b^4 + 25b^5 - 44ab^5 - b^6}{(a-9)(a-7)(a-5)(a-3)(a-1)} \left. + \frac{\Gamma(\frac{b-1}{2})}{\Gamma(\frac{a-8}{2})} \left\{ \frac{1578a - 1456a^2 + 716a^3 - 80a^4 + 10a^5 - 1578b + 1276a^2b - 352a^5b}{(a-8)(a-6)(a-4)(a-2)} \right. \right. \\
 & \left. \left. + \frac{110a^4b + 1456b^3 - 1276ab^2 + 132a^5b^2 - 716b^5 + 352ab^3 - 132a^2b^3 + 80ab^4 - 110ab^4 - 10b^5}{(a-8)(a-6)(a-4)(a-2)} \right\} \right] \quad (6)
 \end{aligned}$$

For  $a < 1$  and  $a > 10$

$$\begin{aligned}
 {}_2F_1(a, b ; \frac{a+b-10}{2}; \frac{1}{2}) = & 2^{b-1} \frac{\Gamma(\frac{a+b-10}{2})}{(a-b)\Gamma(b)} \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-10}{2})} \left\{ \frac{-3840a + 7392a^2 - 3784a^3 + 1276a^4 - 110a^5 + 11a^6 + 3840b}{(a-9)(a-7)(a-5)(a-3)(a-1)} \right. \\
 & + \frac{-3008ab - 3432a^2b + 3872a^3b - 682a^4b + 154a^5b - 4384b^2 + 5416ab^2 - 1672a^2b^2 - 396a^5b^2 + 297a^4b^2 + 1800b^5}{(a-9)(a-7)(a-5)(a-3)(a-1)} \\
 & + \frac{-3136ab^3 + 748a^2b^3 - 132a^3b^2 - 340b^4 + 410ab^4 - 275a^2b^4 + 30b^5 - 54ab^5 - b^6}{(a-9)(a-7)(a-5)(a-3)(a-1)} + \\
 & \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-10}{2})} \left\{ \frac{-3840a + 4384a^2 - 1800a^3 + 340a^4 - 30a^5 + a^6 + 3840b + 3008ab - 5416a^2b + 3136a^3b - 410a^4b + 54a^5b}{(a-10)(a-8)(a-6)(a-4)(a-2)} \right. \\
 & + \frac{-7392b^2 + 3432ab^2 + 1672a^2b^2 - 748a^3b^2 + 275a^4b^2 + 3784b^3 - 3872ab^3 + 396a^2b^3 + 132a^3b^2 - 1276b^4 + 682ab^4}{(a-10)(a-8)(a-6)(a-4)(a-2)} \\
 & \left. + \frac{-297a^5b^4 + 110b^6 - 154ab^6 - 11b^6}{(a-10)(a-8)(a-6)(a-4)(a-2)} \right\} ] \quad (7)
 \end{aligned}$$

### C. DERIVATIONS OF THE SUMMATION FORMULAE:

**Derivation of (5):** Substituting  $c = \frac{a+b-8}{2}$  and  $z = \frac{1}{2}$  in equation (3) , we get

$$(a-b) {}_2F_1(a, b ; \frac{a+b-8}{2}; \frac{1}{2}) = (a-b-8) {}_2F_1(a, b-1 ; \frac{a+b-8}{2}; \frac{1}{2}) + (a-b+8) {}_2F_1(a-1, b ; \frac{a+b-8}{2}; \frac{1}{2})$$

Now with the help of the derived result from equation (4) , we get

$$\begin{aligned}
 \text{L.H.S} = (a-b-8) 2^{b-2} \frac{\Gamma(\frac{a+b-8}{2})}{(a-b+1)\Gamma(b-1)} \frac{\Gamma(\frac{b-1}{2})}{\Gamma(\frac{a-7}{2})} & \left\{ \frac{384 - 240a - 356a^2 + 224a^3 - 43a^4 + a^5 - 784b + 340ab}{(a-7)(a-5)(a-3)(a-1)} \right. \\
 & \left. + \frac{96a^5b - 160a^3b + 27a^4b + 540b^2 - 672ab^2 + 126a^2b^2 + 42a^3b^2 - 160b^3 + 204ab^3 - 42a^2b^3 + 21b^4 - 27ab^4 - b^5}{(a-7)(a-5)(a-3)(a-1)} \right\} \\
 & + \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-6}{2})} \left\{ \frac{-384 - 32a + 256a^2 - 88a^3 + 8a^4 + 672b - 288ab - 72a^2b + 48a^3b - 352b + 216ab^2 + 72b^3 - 48ab^3 - 8b^4}{(a-6)(a-4)(a-2)} \right\}
 \end{aligned}$$

$$\begin{aligned}
 & + (a-b+8) 2^{b-1} \frac{\Gamma(\frac{a+b-8}{2})}{\Gamma(a-1)\Gamma(b)} \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-8}{2})} \right] \left\{ \frac{-384+784a-540a^2+160a^3-21a^4+a^5+240b-340ab+672a^2b}{(a-8)(a-6)(a-4)(a-2)} \right. \\
 & \left. + \frac{-204a^5h+27a^4h+356a^2h^2-96ah^2-126a^2h^3+42a^5h^2-224h^5+180ah^5-42a^2h^5+43h^4-27ah^4-h^5}{(a-8)(a-6)(a-4)(a-2)} \right\} \\
 & + \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a-7}{2})} \left\{ \frac{384-672a+352a^2-72a^3+8a^4+32b+288ab-216a^2b+43a^5b-256b^2+72ab^2+88b^3-48ab^2-8b^4}{(a-7)(a-6)(a-5)} \right\}
 \end{aligned}$$

On simplification , we get the formula (5)

Similarly, we can prove the other formulae.

#### REFERENCES:

- [1] Abramowitz, Milton., A and Stegun, Irene ; *Handbook of Mathematical Functions with Formulas , Graphs , and Mathematical Tables*. National Bureau of Standards, 1970.
- [2] Arora, Asish, Singh, Rahul , Salahuddin ; Development of a family of summation formulae of half argument using Gauss and Bailey theorems , *Journal of Rajasthan academy of Physical Sciences.* , 7(2008), 335-342
- [3] Garg, O.P, Salahuddin , Shakeeluddin ; On Certain Summation Formulae Involving Hypergeometric Function, *International Journal of Computational Science and Mathematics*, 2(2010), 67-76
- [4] Salahuddin; Evaluation of a Summation Formula Involving Recurrence Relation, *Gen. Math. Notes*. 2(2010), 42-59.
- [5] Salahuddin; Two Summation Formulae Based On Half Argument Associated To Hypergeometric function , *Global Journal of ScienceFrontier Research.*, 10(2010), 08-19.
- [6] Salahuddin; Evaluation of Certain Summation Formulae Involving Gauss Theorem , *Global Journal of Mathematical Sciences: Theoryand Practical.*, 10(2010), 309-316
- [7] Salahuddin; A Summation Formula Related To Bailey Theorem, *Global Journal of Science Frontier Research*, 11(2011), 53-67.

\*\*\*\*\*