

ON THE COMPARATIVE GROWTH ANALYSIS
OF A SPECIAL TYPE OF DIFFERENTIAL POLYNOMIAL

Sanjib Kumar Datta^{1*} and Ritam Biswas²

¹*Department of Mathematics, University of Kalyani, Kalyani,
Dist.- Nadia, PIN- 741235, West Bengal, India.*

²*Murshidabad College of Engineering and Technology, Banjetia, Berhampore,
P. O.- Cossimbazar Raj, PIN-742102, West Bengal, India.*

(Received on: 10-08-13; Revised & Accepted on: 22-10-13)

ABSTRACT

In this paper we discuss some comparative growth estimates of composite entire and meromorphic functions and a special type of differential polynomial as considered by Bhooshnurmath and Prasad [4] and generated by one of the factors of the composition.

AMS Subject Classification (2010): 30D30, 30D35.

Keywords and phrases: Order (lower order), entire function, meromorphic function, composition, growth properties, special type of differential polynomial.

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

For any two transcendental entire functions f and g defined in the open complex plane \mathbb{C} , Clunie [5] proved that

$$\lim_{r \rightarrow \infty} \frac{T(r, f \circ g)}{T(r, f)} = \lim_{r \rightarrow \infty} \frac{T(r, f \circ g)}{T(r, g)} = \infty.$$

Singh [10] studied some comparative growth properties of $\log T(r, f \circ g)$ and $T(r, f)$. He [10] also raised the question of investigating the comparative growth of $\log T(r, f \circ g)$ and $T(r, g)$ which he was unable to solve. Lahiri [8] proved some results on the comparative growth of $\log T(r, f \circ g)$ and $T(r, g)$.

Some mathematicians like H. X. Yi [12] and many more studied the comparative growth of a meromorphic function and its derivatives.

Since the natural extension of a derivative is a differential polynomial, in this paper we extend some earlier results for a special type of linear differential polynomial of the form $F = f^n Q[f]$ where $Q[f]$ is a differential polynomial in f and $n = 0, 1, 2, \dots$ as considered by Bhooshnurmath and Prasad [4]. We do not explain the standard notations and definitions in the theory of entire and meromorphic functions because those are available in [11] and [7].

In the sequel we use the following two notations:

(i) $\log^{[k]} x = \log(\log^{[k-1]} x)$ for $k = 1, 2, 3, \dots$; $\log^{[0]} x = x$

and

(ii) $\exp^{[k]} x = \exp(\exp^{[k-1]} x)$ for $k = 1, 2, 3, \dots$; $\exp^{[0]} x = x$.

Corresponding author: Sanjib Kumar Datta^{1*}

**¹Department of Mathematics, University of Kalyani, Kalyani,
Dist.- Nadia, PIN- 741235, West Bengal, India.**

E-mail: sanjib_kr_datta@yahoo.co.in

The following definitions are well known:

Definition: 1 The order ρ_f and lower order λ_f of a meromorphic function f are defined as

$$\rho_f = \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r}$$

and

$$\lambda_f = \liminf_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r}.$$

If f is entire, one can easily verify that

$$\rho_f = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log r}$$

and

$$\lambda_f = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log r}.$$

Definition: 2 The hyper order $\bar{\rho}_f$ and hyper lower order $\bar{\lambda}_f$ of a meromorphic function f are defined as

$$\bar{\rho}_f = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log r}$$

and

$$\bar{\lambda}_f = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log r}.$$

If f is entire, then

$$\bar{\rho}_f = \limsup_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log r}$$

and

$$\bar{\lambda}_f = \liminf_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log r}.$$

Definition: 3 [9] Let f be a meromorphic function of order zero. Then the quantities ρ_f^* , λ_f^* and $\bar{\rho}_f^*$, $\bar{\lambda}_f^*$ are defined in the following way

$$\rho_f^* = \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log^{[2]} r},$$

$$\lambda_f^* = \liminf_{r \rightarrow \infty} \frac{\log T(r, f)}{\log^{[2]} r}$$

and

$$\bar{\rho}_f^* = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log^{[2]} r}$$

$$\bar{\lambda}_f^* = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log^{[2]} r}.$$

If f is entire then clearly

$$\rho_f^* = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log^{[2]} r},$$

$$\lambda_f^* = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log^{[2]} r}.$$

and

$$\bar{\rho}_f^* = \limsup_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log^{[2]} r},$$

$$\bar{\lambda}_f^* = \liminf_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log^{[2]} r}.$$

Definition: 4 The type σ_f of a meromorphic function f is defined as follows

$$\sigma_f = \limsup_{r \rightarrow \infty} \frac{T(r, f)}{r^{\rho_f}}, 0 < \rho_f < \infty.$$

When f is entire, then

$$\sigma_f = \limsup_{r \rightarrow \infty} \frac{\log M(r, f)}{r^{\rho_f}}, 0 < \rho_f < \infty.$$

Definition: 5 A meromorphic function $a \equiv a(z)$ is called small with respect to f if $T(r, a) = S(r, f)$.

2. LEMMAS

In this section we present some lemmas which will be needed in the sequel.

Lemma: 1 [5] If f and g be any two entire functions then for all sufficiently large values of r ,

$$M(r, f \circ g) \geq M\left(\frac{1}{8}M\left(\frac{r}{2}, g\right) - |g(0)|, f\right).$$

Lemma: 2 [1] Let f be meromorphic and g be entire then for all sufficiently large values of r ,

$$T(r, f \circ g) \leq \{1 + o(1)\} \frac{T(r, g)}{\log M(r, g)} T(M(r, g), f).$$

Lemma: 3 [3] Let f be meromorphic and g be entire and suppose that $0 < \mu < \rho_g \leq \infty$. Then for a sequence of values of r tending to infinity,

$$T(r, f \circ g) \geq T(\exp(r^\mu), f).$$

Lemma: 4 [4] Let $F = f^n Q[f]$ where $Q[f]$ is a differential polynomial in f . If $n \geq 1$ then $\rho_F = \rho_f$ and $\lambda_F = \lambda_f$.

Lemma: 5 Let $F = f^n Q[f]$ where $Q[f]$ is a differential polynomial in f . If $n \geq 1$ then

$$\lim_{r \rightarrow \infty} \frac{T(r, F)}{T(r, f)} = 1.$$

The proof of Lemma 5 directly follows from Lemma 4.

In the line of Lemma 4 we may prove the following lemma:

Lemma: 6 Let $F = f^n Q[f]$ where $Q[f]$ is a differential polynomial in f . If $n \geq 1$ then $\bar{\rho}_F = \bar{\rho}_f$ and $\bar{\lambda}_F = \bar{\lambda}_f$.

Lemma: 7 Let f be meromorphic and g be transcendental entire such that $\rho_f = 0$ and $\rho_g < \infty$. Then $\rho_{f \circ g} \leq \rho_f^* \cdot \rho_g$.

Proof: In view of Lemma 2 and the inequality $T(r, g) \leq \log^+ M(r, g)$, we get that

$$\begin{aligned} \rho_{f \circ g} &= \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{\log r} \leq \limsup_{r \rightarrow \infty} \frac{\log T(M(r, g), f) + o(1)}{\log r} \\ &= \limsup_{r \rightarrow \infty} \frac{\log T(M(r, g), f)}{\log [2]M(r, g)} \cdot \limsup_{r \rightarrow \infty} \frac{\log [2]M(r, g)}{\log r} = \rho_f^* \cdot \rho_g. \end{aligned}$$

This proves the lemma.

Remark: 1 The sign ' \leq ' in Lemma 7 cannot be removed by ' $<$ ' only as we see in the following example.

Example: 1 Let $f = z$ and $g = \exp z$. Then $\rho_{f \circ g} = 1, \rho_g = 1$ and $\rho_f = 0$. So

$$\rho_f^* = \limsup_{r \rightarrow \infty} \frac{\log [2]M(r, f)}{\log [2]r} = \limsup_{r \rightarrow \infty} \frac{\log [2]r}{\log [2]r} = 1.$$

Therefore

$$\rho_{f \circ g} = \rho_f^* \cdot \rho_g.$$

3. THEOREMS

In this section we present the main results of the paper.

Theorem: 1 Let f be transcendental meromorphic and g be entire satisfying the following conditions:

i) ρ_f and ρ_g are both finite,

ii) ρ_f is positive and

iii) let $F = f^n Q[f]$ for $n \geq 1$. Then for $p' > 0$ and each $\alpha \in (-\infty, \infty)$,

$$\liminf_{r \rightarrow \infty} \frac{\{\log T(r, f_o g)\}^{1+\alpha}}{\log T(\exp(r^{p'}), F)} = 0 \quad \text{if } p' > (1 + \alpha)\rho_g.$$

Proof: If $1 + \alpha \leq 0$, the theorem is trivial. So we take $1 + \alpha > 0$. Since $T(r, g) \leq \log^+ M(r, g)$, by Lemma 2, we get for all sufficiently large values of r that

$$T(r, f_o g) \leq \{1 + o(1)\}T(M(r, g), f)$$

i.e.,

$$\log T(r, f_o g) \leq \log\{1 + o(1)\} + \log T(M(r, g), f)$$

i.e.,

$$\log T(r, f_o g) \leq o(1) + (\rho_f + \varepsilon) \log M(r, g)$$

i.e.,

$$\log T(r, f_o g) \leq o(1) + (\rho_f + \varepsilon)r^{(\rho_g + \varepsilon)}$$

i.e.,

$$\log T(r, f_o g) \leq r^{(\rho_g + \varepsilon)}\{(\rho_f + \varepsilon) + o(1)\}$$

i.e.,

$$\{\log T(r, f_o g)\}^{1+\alpha} \leq r^{(\rho_g + \varepsilon)(1+\alpha)}\{(\rho_f + \varepsilon) + o(1)\}^{1+\alpha}. \quad (1)$$

Again in view of Lemma 4, we have for a sequence of values of r tending to infinity and for $\varepsilon > 0$,

$$\log T(\exp(r^{p'}), F) > (\rho_f - \varepsilon) \log(\exp(r^{p'})) = (\rho_f - \varepsilon)r^{p'}. \quad (2)$$

Now combining (1) and (2) we obtain for a sequence of values of r tending to infinity that

$$\frac{\{\log T(r, f_o g)\}^{1+\alpha}}{\log T(\exp(r^{p'}), F)} \leq \frac{r^{(\rho_g + \varepsilon)(1+\alpha)}\{(\rho_f + \varepsilon) + o(1)\}^{1+\alpha}}{(\rho_f - \varepsilon)r^{p'}}$$

from which the theorem follows because we can choose ε such that

$$0 < \varepsilon < \min\{\rho_f, \frac{p'}{1 + \alpha} - \rho_g\}.$$

This proves the theorem.

Remark: 2 The condition $p' > (1 + \alpha)\rho_g$ is essential in Theorem 1 as we see in the next example.

Example: 2 Let $f = \exp z$, $g = \exp z$, $\alpha = 0$ and $p' = 1$. Then

$$\rho_f = 1 = \rho_g$$

and

$$\sum_{a \neq \infty} \delta(a; f) + \delta(\infty; f) = 2.$$

Also let $F = f^n Q[f]$ for $n \geq 1$.

Taking $n = 1$, $A_j = 1$, $n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$; we see that $F = \exp(2z)$.

Now we have

$$\begin{aligned} \log T(r, f_o g) &= \log T(r, \exp^{[2]} z) \sim \log \left\{ \frac{\exp r}{(2\pi^3 r)^2} \right\} \quad (r \rightarrow \infty) \\ &\sim r - \frac{1}{2} \log r + O(1) \quad (r \rightarrow \infty). \end{aligned}$$

Therefore

$$\begin{aligned} \liminf_{r \rightarrow \infty} \frac{\{\log T(r, f_o g)\}^{1+\alpha}}{\log T(\exp(r^{p'}), F)} &= \liminf_{r \rightarrow \infty} \frac{\log T(r, f_o g)}{\log T(\exp r, \exp 2z)} \\ &= \liminf_{r \rightarrow \infty} \frac{r - \frac{1}{2} \log r + O(1)}{\log \left\{ \frac{\exp r}{\pi} \right\}} \end{aligned}$$

$$= \liminf_{r \rightarrow \infty} \frac{r - \frac{1}{2} \log r + O(1)}{r + O(1)} = 1.$$

which contradicts Theorem 1.

Theorem: 2 If f be meromorphic and g be transcendental entire such that $\rho_g < \infty$, $\rho_{f \circ g} = \infty$ and for $n \geq 1$, $G = g^n Q[g]$, then for every $A > 0$,

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{\log T(r^A, G)} = \infty.$$

Proof: If possible, let there exist a constant β such that for all sufficiently large values of r , we have

$$\log T(r, f \circ g) \leq \beta \log T(r^A, G). \quad (3)$$

In view of Lemma 4, for all sufficiently large values of r , we get that

$$\log T(r^A, G) \leq (\rho_G + \varepsilon) A \log r$$

$$\text{i.e., } \log T(r^A, G) \leq (\rho_g + \varepsilon) A \log r. \quad (4)$$

Now combining (3) and (4), we obtain for all sufficiently large values of r that

$$\log T(r, f \circ g) \leq \beta (\rho_g + \varepsilon) A \log r$$

i.e.,

$$\rho_{f \circ g} \leq \beta A (\rho_g + \varepsilon),$$

which contradicts the condition $\rho_{f \circ g} = \infty$. So for a sequence of values of r tending to infinity, it follows that

$$\log T(r, f \circ g) > \beta \log T(r^A, G),$$

from which the theorem follows.

Corollary: 1 Under the assumption of Theorem 2,

$$\limsup_{r \rightarrow \infty} \frac{T(r, f \circ g)}{T(r^A, G)} = \infty.$$

Proof: By Theorem 2 we obtain for all sufficiently large values of r and for $K_1 > 1$ that

$$\log T(r, f \circ g) > K_1 \log T(r^A, G)$$

$$\text{i.e., } T(r, f \circ g) > \{T(r^A, G)\}^{K_1},$$

from which the corollary follows.

Remark: 3 The condition $\rho_{f \circ g} = \infty$ is necessary in Theorem 2 and Corollary 1 which is evident from the following example.

Example: 3 Let $f = z$, $g = \exp z$ and $A = 1$. Then $\rho_g = 1 < \infty$ and $\rho_{f \circ g} = 1 < \infty$.

Let $G = g^n Q[g]$ for $n \geq 1$. Taking $n = 1$, $A_j = 1$, $n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$; we see that $G = \exp(2z)$.

Now we have

$$T(r, f \circ g) = T(r, \exp z) = \frac{r}{\pi}$$

$$\text{and } T(r^A, G) = T(r, \exp 2z) = \frac{2r}{\pi}.$$

Therefore

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{\log T(r^A, G)} = \limsup_{r \rightarrow \infty} \frac{\log r + O(1)}{\log r + O(1)} = 1$$

$$\text{and } \limsup_{r \rightarrow \infty} \frac{T(r, f_0 g)}{T(r^A, G)} = \limsup_{r \rightarrow \infty} \frac{\left(\frac{r}{\pi}\right)}{\left(\frac{2r}{\pi}\right)} = \frac{1}{2},$$

which is contrary to Theorem 2.

Remark: 4 If we take $\rho_f < \infty$ and $F = f^n Q[f]$ for $n \geq 1$ instead of $\rho_g < \infty$ and $G = g^n Q[g]$ for $n \geq 1$ respectively, then Theorem 2 and Corollary 1 remain valid with G replaced by F in the denominator as we see in the following theorem and corollary.

Theorem: 3 If f be transcendental meromorphic and g be entire such that $\rho_f < \infty, \rho_{f_0 g} = \infty$ and for $n \geq 1$, $F = f^n Q[f]$, then for every $A > 0$,

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f_0 g)}{\log T(r^A, F)} = \infty.$$

Proof: If possible let there exist a constant γ such that for all sufficiently large values of r , we have $\log T(r, f_0 g) \leq \gamma \log T(r^A, F)$.

In view of Lemma 5, for all sufficiently large values of r we get that

$$\log T(r^A, F) \leq (\rho_f + \varepsilon) A \log r.$$

Now combining the above two inequalities, we get for all sufficiently large values of r that

$$\log T(r, f_0 g) \leq \gamma (\rho_f + \varepsilon) A \log r$$

$$\text{i.e., } \rho_{f_0 g} \leq \gamma A (\rho_f + \varepsilon),$$

which contradicts the condition $\rho_{f_0 g} = \infty$. So for a sequence of values of r tending to infinity, it follows that

$$\log T(r, f_0 g) > \gamma \log T(r^A, F),$$

from which the theorem follows.

Corollary: 2 Under the assumptions of Theorem 3,

$$\limsup_{r \rightarrow \infty} \frac{T(r, f_0 g)}{T(r^A, F)} = \infty.$$

Proof: In view of Theorem 3, we obtain for all sufficiently large values of r and for $K_2 > 1$ that

$$\log T(r, f_0 g) > K_2 \log T(r^A, F)$$

$$\text{i.e., } T(r, f_0 g) > \{T(r^A, F)\}^{K_2},$$

from which the corollary follows.

Remark: 5 The condition $\rho_{f_0 g} = \infty$ is necessary in Theorem 3 and Corollary 2 which is evident from the following example.

Example: 4 Let $f = \exp z$, $g = z$ and $A = 1$. Then $\rho_f = 1 < \infty$, $\rho_{f_0 g} = 1 < \infty$ and for $n \geq 1$, $F = f^n Q[f]$. Taking $n = 1, A_j = 1, n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$; we see that $F = \exp(2z)$. Now we have

$$T(r, f_0 g) = T(r, \exp z) = \frac{r}{\pi} \text{ and } T(r^A, F) = T(r, \exp 2z) = \frac{2r}{\pi}.$$

Therefore

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f_0 g)}{\log T(r^A, F)} = \limsup_{r \rightarrow \infty} \frac{\log r + O(1)}{\log r + O(1)} = 1 \text{ and } \limsup_{r \rightarrow \infty} \frac{T(r, f_0 g)}{T(r^A, F)} = 1,$$

which contradicts Theorem 3.

Theorem: 4 Let f and g be any two entire functions with $\lambda_f > 0$ and $\rho_f < \lambda_g$. Also let f be transcendental with $F = f^n Q[f]$ for $n \geq 1$. Then

$$\lim_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f_0 g)}{\log M(r, F)} = \infty.$$

Proof: In view of Lemma 1, we get for all sufficiently large values of r that

$$M(r, f_0 g) \geq M\left(\frac{1}{16} M\left(\frac{r}{2}, g\right), f\right)$$

$$\text{i.e., } \log^{[2]} M(r, f_0 g) \geq \log^{[2]} M\left(\frac{1}{16} M\left(\frac{r}{2}, g\right), f\right)$$

$$\text{i.e., } \log^{[2]} M(r, f_0 g) \geq (\lambda_f - \varepsilon) \log \frac{1}{16} + (\lambda_f - \varepsilon) \log M\left(\frac{r}{2}, g\right)$$

$$\text{i.e., } \log^{[2]} M(r, f_0 g) \geq O(1) + (\lambda_f - \varepsilon) \left(\frac{r}{2}\right)^{(\lambda_g - \varepsilon)}. \quad (5)$$

Again for all sufficiently large values of r , we get by Lemma 5 that

$$\log M(r, F) \leq r^{(\rho_f + \varepsilon)} = r^{(\rho_f + \varepsilon)}. \quad (6)$$

Now combining (5) and (6), it follows for all sufficiently large values of r that

$$\frac{\log^{[2]} M(r, f_0 g)}{\log M(r, F)} \geq \frac{O(1) + (\lambda_f - \varepsilon) \left(\frac{r}{2}\right)^{(\lambda_g - \varepsilon)}}{r^{(\rho_f + \varepsilon)}}. \quad (7)$$

Since $\rho_f < \lambda_g$, we can choose $\varepsilon (> 0)$ in such a way that

$$\rho_f + \varepsilon < \lambda_g - \varepsilon. \quad (8)$$

Thus from (7) and (8) we obtain that

$$\liminf_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f_0 g)}{\log M(r, F)} = \infty,$$

from which the theorem follows.

Remark: 6 The condition $\rho_f < \lambda_g$ is necessary in Theorem 4 which is evident from the following two examples.

Example: 5 Let $f = \exp z$ and $g = \exp z$. Then $\lambda_f = 1 > 0$, $\rho_f = 1 = \lambda_g$ and $F = f^n Q[f]$ for $n \geq 1$. Taking $n = 1, A_j = 1, n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$; we see that $F = \exp(2z)$. Again $M(r, f_0 g) = \exp^{[2]} r$ and $M(r, F) = \exp(2r)$.

Therefore

$$\lim_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f_0 g)}{\log M(r, F)} = \lim_{r \rightarrow \infty} \frac{\log^{[2]} (\exp^{[2]} r)}{\log (\exp 2r)} = \lim_{r \rightarrow \infty} \frac{r}{2r} = \frac{1}{2},$$

which is contrary to Theorem 4.

Example: 6 Let $f = \exp z$ and $g = z$. Then $\lambda_f = 1 > 0$, $\rho_f = 1 > 0 = \lambda_g$ and let $F = f^n Q[f]$ for $n \geq 1$. Taking $n = 1, A_j = 1, n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$; we see that $F = \exp(2z)$. Again $M(r, f_0 g) = \exp r$ and $M(r, F) = \exp(2r)$.

Therefore

$$\lim_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f_0 g)}{\log M(r, F)} = \lim_{r \rightarrow \infty} \frac{\log^{[2]} (\exp r)}{\log (\exp 2r)} = \lim_{r \rightarrow \infty} \frac{\log r}{2r} = 0,$$

which contradicts Theorem 4.

Theorem: 5 If f be a transcendental meromorphic function and g be entire with $0 < \lambda_f \leq \rho_f < \infty$, $\rho_g < \infty$ and $F = f^n Q[f]$ for $n \geq 1$, then

$$\lim_{r \rightarrow \infty} \frac{T(r, f_0 g) T(r, F)}{T(\exp(r^p), F)} = 0,$$

if $p' > \rho_g$.

Proof: Since $T(r, g) \leq \log^+ M(r, g)$, for all sufficiently large values of r we get from Lemma 2 that $T(r, f_0 g) \leq \{1 + o(1)\} T(M(r, g), f)$

$$\text{i.e., } T(r, f_0 g) \leq \{1 + o(1)\} \exp\{(\rho_f + \varepsilon) r^{(\rho_g + \varepsilon)}\}. \quad (9)$$

Again by Lemma 5, we obtain for all sufficiently large values of r that

$$T(r, F) \leq r^{(\rho_F + \varepsilon)} = r^{(\rho_f + \varepsilon)}. \quad (10)$$

Now combining (9) and (10), it follows for all sufficiently large values of r that

$$T(r, f_0 g) T(r, F) \leq \{1 + o(1)\} r^{(\rho_f + \varepsilon)} \exp\{(\rho_f + \varepsilon) r^{(\rho_g + \varepsilon)}\}. \quad (11)$$

Also in view of Lemma 4, we have for all sufficiently large values of r ,

$$\log T(\exp(r^{p'}), F) \geq (\lambda_F - \varepsilon) \log\{\exp(r^{p'})\}$$

$$\text{i.e., } \log T(\exp(r^{p'}), F) \geq \exp\{(\lambda_f - \varepsilon) r^{p'}\}. \quad (12)$$

From (11) and (12) it follows for all sufficiently large values of r that

$$\frac{T(r, f_0 g) T(r, F)}{\log T(\exp(r^{p'}), F)} \leq \frac{\{1 + o(1)\} r^{(\rho_f + \varepsilon)} \exp\{(\rho_f + \varepsilon) r^{(\rho_g + \varepsilon)}\}}{\exp\{(\lambda_f - \varepsilon) r^{p'}\}}. \quad (13)$$

As $p' > \rho_g$ so we can choose $\varepsilon(> 0)$ such that

$$p' > \rho_g + \varepsilon. \quad (14)$$

Thus the theorem follows from (13) and (14).

Theorem: 6 Let f be a transcendental meromorphic function and g be a transcendental entire function such that $0 < \lambda_f \leq \rho_f < \infty$ and for $n \geq 1$, $F = f^n Q[f]$. Then for every $A > 0$,

$$\lim_{r \rightarrow \infty} \frac{\log T(r, f_0 g)}{\log T(r^A, F)} = \infty.$$

If further $\rho_g < \infty$ and for $n \geq 1$, $G = g^n Q[g]$, then

$$\lim_{r \rightarrow \infty} \frac{\log T(r, f_0 g)}{\log T(r^A, G)} = \infty.$$

Proof: Since $\lambda_f > 0$, $\lambda_{f_0 g} = \infty$ {cf. [2]}. So it follows that for arbitrary large N and for all sufficiently large values of r ,

$$\log T(r, f_0 g) > AN \log r. \quad (15)$$

Again since $\rho_f < \infty$, for all sufficiently large values of r we get by Lemma 4 that

$$\log T(r^A, F) < A(\rho_f + 1) \log r. \quad (16)$$

Now from (15) and (16), it follows for all sufficiently large values of r that

$$\frac{\log T(r, f_0 g)}{\log T(r^A, F)} > \frac{AN \log r}{A(\rho_f + 1) \log r}$$

$$\text{and so } \lim_{r \rightarrow \infty} \frac{\log T(r, f_0 g)}{\log T(r^A, F)} = \infty.$$

Again since $\rho_g < \infty$, then for all sufficiently large values of r we obtain by Lemma 5 that

$$\log T(r^A, G) < A(\rho_g + 1) \log r. \quad (17)$$

Now from (15) and (17), it follows for all sufficiently large values of r that

$$\frac{\log T(r, f_0 g)}{\log T(r^A, G)} > \frac{AN \log r}{A(\rho_g + 1) \log r}. \quad (18)$$

Thus the theorem follows from (18).

Theorem: 7 Let f be a transcendental meromorphic function with $0 < \lambda_f \leq \rho_f < \infty$ and for $n \geq 1$, $F = f^n Q[f]$ and g be entire. Then

$$\lim_{r \rightarrow \infty} \sup \frac{\log^{[2]} T(\exp(r^{\rho_g}), f_o g)}{\log T(\exp(r^\mu), F)} = \infty, \text{ where } 0 < \mu < \rho_g.$$

Proof: Let $0 < \mu' < \rho_g$. Then in view of Lemma 3, we get for a sequence of values of r tending to infinity that

$$\log T(r, f_o g) \geq \log T(\exp(r^{\mu'}), f)$$

$$\text{i.e., } \log T(r, f_o g) \geq (\lambda_f - \varepsilon) \log\{\exp(r^{\mu'})\}$$

$$\text{i.e., } \log T(r, f_o g) \geq (\lambda_f - \varepsilon) r^{\mu'}$$

$$\text{i.e., } \log^{[2]} T(r, f_o g) \geq O(1) + \mu' \log r.$$

So for a sequence of values of r tending to infinity,

$$\log^{[2]} T(\exp(r^{\rho_g}), f_o g) \geq O(1) + \mu' \log\{\exp(r^{\rho_g})\}$$

$$\text{i.e., } \log^{[2]} T(\exp(r^{\rho_g}), f_o g) \geq O(1) + \mu' r^{\rho_g}. \quad (19)$$

Again in view of Lemma 4, we obtain for all sufficiently large values of r that

$$\log T(\exp(r^\mu), F) \leq (\rho_f + \varepsilon) \log\{\exp(r^\mu)\}$$

$$\text{i.e., } \log T(\exp(r^\mu), F) \leq (\rho_f + \varepsilon) r^\mu. \quad (20)$$

Combining (19) and (20), it follows for a sequence of values of r tending to infinity that

$$\frac{\log^{[2]} T(\exp(r^{\rho_g}), f_o g)}{\log T(\exp(r^\mu), F)} \geq \frac{O(1) + \mu' r^{\rho_g}}{(\rho_f + \varepsilon) r^\mu}. \quad (21)$$

Since $\mu < \rho_g$, we get from (21) that

$$\lim_{r \rightarrow \infty} \sup \frac{\log^{[2]} T(\exp(r^{\rho_g}), f_o g)}{\log T(\exp(r^\mu), F)} = \infty.$$

This proves the theorem.

Remark: 7 The condition $\mu < \rho_g$ in Theorem 7 is essential as we see in the following example.

Example: 7 Let $f = \exp z$, $g = z$ and $\mu = 1$. Then $\lambda_f = 1 = \rho_f$, $\rho_g = 0$ and let for $n \geq 1$, $F = f^n Q[f]$. Taking $n = 1$, $A_j = 1$, $n_{0j} = 1$ and $n_{1j} = \dots = n_{kj} = 0$;

we see that $F = \exp(2z)$. Also

$$T(r, \exp z) = \frac{r}{\pi}.$$

So

$$\log^{[2]} T(\exp(r^{\rho_g}), f_o g) = \log^{[2]} T(e, \exp z) = \log^{[2]} \left(\frac{e}{\pi} \right) = O(1)$$

$$\text{and } \log T(\exp(r^\mu), F) = \log T(\exp r, \exp 2z) = \log \left\{ \frac{2 \exp r}{\pi} \right\} = r + O(1).$$

Therefore

$$\lim_{r \rightarrow \infty} \sup \frac{\log^{[2]} T(\exp(r^{\rho_g}), f_o g)}{\log T(\exp(r^\mu), F)} = \lim_{r \rightarrow \infty} \sup \frac{O(1)}{r + O(1)} = 0,$$

which is contrary to Theorem 7.

Theorem: 8 Let f be rational and g be transcendental meromorphic satisfying

$$(i) \ 0 < \bar{\lambda}_{f_o g} \leq \bar{\rho}_{f_o g} < \infty,$$

$$(ii) \ 0 < \bar{\lambda}_g \leq \bar{\rho}_g < \infty \text{ and}$$

(iii) for $n \geq 1$, $G = g^n Q[g]$. Then for any positive number A ,

$$\frac{\bar{\lambda}_{f_0g}}{A\bar{\rho}_g} \leq \liminf_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{\bar{\lambda}_{f_0g}}{A\bar{\lambda}_g} \leq \limsup_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{\bar{\rho}_{f_0g}}{A\bar{\lambda}_g}.$$

Proof: From the definition of hyper order and hyper lower order and by Lemma 6, we get for arbitrary positive ε and for all sufficiently large values of r that

$$\log^{[2]}T(r, f_0g) \geq (\bar{\lambda}_{f_0g} - \varepsilon) \log r \quad (22)$$

$$\text{and } \log^{[2]}T(r^A, G) \leq (\bar{\rho}_G + \varepsilon) \log r^A$$

$$\text{i.e., } \log^{[2]}T(r^A, G) \leq A(\bar{\rho}_g + \varepsilon) \log r. \quad (23)$$

Combining (22) and (23), we obtain for all sufficiently large values of r that

$$\frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \geq \frac{(\bar{\lambda}_{f_0g} - \varepsilon) \log r}{A(\bar{\rho}_g + \varepsilon) \log r}.$$

Since $\varepsilon(> 0)$ is arbitrary, it follows from above that

$$\liminf_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \geq \frac{\bar{\lambda}_{f_0g}}{A\bar{\rho}_g}. \quad (24)$$

Again for a sequence of values of r tending to infinity,

$$\log^{[2]}T(r, f_0g) \leq (\bar{\lambda}_{f_0g} + \varepsilon) \log r. \quad (25)$$

Also in view of Lemma 6, we have for all sufficiently large values of r ,

$$\log^{[2]}T(r^A, G) \geq (\bar{\lambda}_G - \varepsilon) \log r^A$$

$$\text{i.e., } \log^{[2]}T(r^A, G) \geq A(\bar{\lambda}_g - \varepsilon) \log r. \quad (26)$$

Combining (25) and (26), we get for a sequence of values of r tending to infinity that

$$\frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{(\bar{\lambda}_{f_0g} + \varepsilon) \log r}{A(\bar{\lambda}_g - \varepsilon) \log r}.$$

As $\varepsilon(> 0)$ is arbitrary, it follows from above that

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{\bar{\lambda}_{f_0g}}{A\bar{\lambda}_g}. \quad (27)$$

Also for a sequence of values of r tending to infinity and by Lemma 6,

$$\log^{[2]}T(r^A, G) \leq A(\bar{\lambda}_G + \varepsilon) \log r$$

$$\text{i.e., } \log^{[2]}T(r^A, G) \leq A(\bar{\lambda}_g + \varepsilon) \log r. \quad (28)$$

Combining (22) and (28), we get for a sequence of values of r tending to infinity that

$$\frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \geq \frac{\bar{\lambda}_{f_0g}}{A\bar{\lambda}_g}.$$

Since $\varepsilon(> 0)$ is arbitrary, it follows from above that

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \geq \frac{\bar{\lambda}_{f_0g}}{A\bar{\lambda}_g}. \quad (29)$$

Also for all sufficiently large values of r ,

$$\log^{[2]}T(r, f_0g) \leq (\bar{\rho}_{f_0g} + \varepsilon) \log r. \quad (30)$$

From (26) and (30), we obtain for all sufficiently large values of r that

$$\frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{(\bar{\rho}_{f_0g} + \varepsilon) \log r}{A(\bar{\lambda}_g - \varepsilon) \log r}.$$

Since $\varepsilon(> 0)$ is arbitrary, it follows from above that

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]}T(r, f_0g)}{\log^{[2]}T(r^A, G)} \leq \frac{\bar{\rho}_{f_0g}}{A\bar{\lambda}_g}. \quad (31)$$

Thus the theorem follows from (24), (27), (29) and (31).

Theorem: 9 Let f be meromorphic and g be transcendental entire such that

- (i) $0 < \rho_g < \infty$,
- (ii) $\sigma_g > 0$,
- (iii) $0 < \rho_{f_0g} < \infty$,
- (iv) $\sigma_{f_0g} < \infty$,
- (v) $\rho_f^* < 1$ and
- (vi) for $n \geq 1$, $G = g^n Q[g]$. Then

$$\liminf_{r \rightarrow \infty} \frac{T(r, f_0g)}{T(r, G)} = 0.$$

Proof: From the definition of type, we have for arbitrary positive ε and for all sufficiently large values of r ,

$$T(r, f_0g) \leq (\sigma_{f_0g} + \varepsilon)r^{\rho_{f_0g}}. \quad (32)$$

Again in view of Lemma 4, we get for a sequence of values of r tending to infinity that

$$T(r, G) \geq (\sigma_G - \varepsilon)r^{\rho_g}$$

$$\text{i.e., } T(r, G) \geq (\sigma_g - \varepsilon)r^{\rho_g}. \quad (33)$$

Since $\rho_{f_0g} < \infty$, it follows that $\rho_f = 0$ {cf. [6]}. So in view of Lemma 7, from (32) and (33), we obtain for a sequence of values of r tending to infinity that

$$\frac{T(r, f_0g)}{T(r, G)} \leq \frac{(\sigma_{f_0g} + \varepsilon)r^{\rho_f^* \rho_g}}{(\sigma_g - \varepsilon)r^{\rho_g}}$$

$$\text{i.e., } \frac{T(r, f_0g)}{T(r, G)} \leq \frac{(\sigma_{f_0g} + \varepsilon)r^{(\rho_f^* - 1)\rho_g}}{(\sigma_g - \varepsilon)}.$$

Since $\varepsilon(> 0)$ is arbitrary, in view of condition (v), it follows that

$$\liminf_{r \rightarrow \infty} \frac{T(r, f_0g)}{T(r, G)} = 0.$$

This proves the theorem.

Remark: 8 The condition $\rho_f^* < 1$ in Theorem 9 is essential which is evident from the following example.

Example: 8 Let $f = z$ and $g = \exp z$. Then $\rho_g = 1 = \sigma_g$, $\rho_{f_0g} = 1 = \sigma_{f_0g}$, $\rho_f = 0$ and let $G = g^n Q[g]$ for $n \geq 1$. Taking $n = 1$, $A_j = 1$, $n_{0j} = 1$ and

$n_{1j} = \dots = n_{kj} = 0$; we see that $G = \exp(2z)$. Also we have

$$\rho_f^* = \limsup_{r \rightarrow \infty} \frac{\log^{[2]}M(r, f)}{\log^{[2]}r} = \limsup_{r \rightarrow \infty} \frac{\log^{[2]}r}{\log^{[2]}r} = 1.$$

Again

$$T(r, f_0g) = \frac{r}{\pi} \text{ and } T(r, G) = \frac{2r}{\pi}.$$

$$\text{Therefore } \liminf_{r \rightarrow \infty} \frac{T(rf_0g)}{T(r,G)} = \liminf_{r \rightarrow \infty} \frac{(\frac{r}{\pi})}{(\frac{2r}{\pi})} = \frac{1}{2},$$

which contradicts Theorem 9.

REFERENCES

- [1] Bergweiler, W.: *On the Nevanlinna Characteristic of a composite function*, Complex Variables, Vol. 10(1988), pp. 225-236.
- [2] Bergweiler, W.: *Order and lower order of composite meromorphic functions*, Michigan Math. J., Vol. 36(1989), pp. 135-146.
- [3] Bergweiler, W.: *On the growth rate of composite meromorphic functions*, Complex Variables, Vol. 14 (1990), pp. 187-196.
- [4] Bhooshnurmath, S. S. and Prasad, K. S. L. N.: *The value distribution of some differential polynomials*, Bull. Cal. Math. Soc., Vol. 101, No. 1 (2009), pp. 55-62.
- [5] Clunie, J.: *The composition of entire and meromorphic functions*, Mathematical essays dedicated to A. J. Macintyre, Ohio University Press (1970), pp. 75-92.
- [6] Gross, F.: *Factorization of meromorphic functions*, U. S. Government Printing Office, Washington, D. C., 1972.
- [7] Hayman, W. K.: *Meromorphic functions*, The Clarendon Press, Oxford, 1964.
- [8] Lahiri, I.: *Growth of composite integral functions*, Indian J. Pure Appl. Math., Vol. 20, No. 9 (September 1989), pp. 899-907.
- [9] Liao, L. and Yang, C. C.: *On the growth of composite entire functions*, Yokohama Math. J. Vol. 46 (1999), pp. 97-107.
- [10] Singh, A. P.: *Growth of composite entire functions*, Kodai Math. J., Vol. 8 (1985), pp. 99-102.
- [11] Valiron, G.: *Lectures on the general theory of integral functions*, Chelsea Publishing Company, 1949.
- [12] Yi, H. X.: *On characteristic function of a meromorphic function and its derivative*, Indian J. Math., Vol. 33, No. 2 (1991), pp. 119-133.

Source of support: Nil, Conflict of interest: None Declared