

## GENERALIZED FUZZY SOFT SETS: A NEW APPROACH

**Manash Jyoti Borah\***

*Department of Mathematics, Bahona College, Jorhat, Assam, India.*

*(Received On: 04-07-14; Revised & Accepted On: 22-07-14)*

### ABSTRACT

*In this paper, we redefined null, absolute, union, and intersection of generalized fuzzy soft sets and study some of their properties. We also defined disjunctive sum, difference and product of two generalized fuzzy soft sets and their basic properties.*

**Key Words:** *Fuzzy soft sets, Generalized fuzzy soft set, Disjunctive sum, Difference and Product of fuzzy soft set.*

### 1. INTRODUCTION

Maji *et al.* [4] introduced the concept of Fuzzy Soft Set and some properties regarding fuzzy soft union, intersection, complement of a fuzzy soft set, De Morgan Law etc. In 2011, Neog and Sut [6] put forward some more propositions regarding fuzzy soft set theory. In 2009 Majumder and Samanta [5] initiated another important part, known as Generalized Fuzzy Soft Set Theory. While generalizing the concept of Fuzzy Soft Sets, Majumder and Samanta [5] considered the same set of parameter. But in 2012 Borah, Neog and Sut [3] considering generalized fuzzy soft sets different sets of parameters. In our work, we redefined null, absolute, union, and intersection of generalized fuzzy soft sets and study some of their properties. We also defined disjunctive sum, difference and product of two generalized fuzzy soft sets and their basic properties.

### 2. PRELIMINARIES

In this section, we recall some basic concepts and definitions regarding fuzzy soft sets and generalized fuzzy soft sets.

**Definition 2.1:** [4] A pair  $(F, A)$  is called a fuzzy soft set over  $U$  where  $F : A \rightarrow \tilde{P}(U)$  is a mapping from  $A$  into  $\tilde{P}(U)$ .

**Definition 2.2:** [4] A soft set  $(F, A)$  over  $U$  is said to be null fuzzy soft set denoted by  $\phi$  if  $\forall \varepsilon \in A, F(\varepsilon)$  is the null fuzzy set  $\bar{0}$  of  $U$  where  $\bar{0}(x) = 0 \forall x \in U$ .

**Definition 2.3:** [4] A soft set  $(F, A)$  over  $U$  is said to be absolute fuzzy soft set denoted by  $\tilde{A}$  if  $\forall \varepsilon \in A, F(\varepsilon)$  is the null fuzzy set  $\bar{1}$  of  $U$  where  $\bar{1}(x) = 1 \forall x \in U$

**Definition 2.4:** [4] For two fuzzy soft sets  $(F, A)$  and  $(G, B)$  in a fuzzy soft class  $(U, E)$ , we say that  $(F, A)$  is a fuzzy soft subset of  $(G, B)$ , if

- (i)  $A \subseteq B$
- (ii) For all  $\varepsilon \in A, F(\varepsilon) \subseteq G(\varepsilon)$  and is written as  $(F, A) \subseteq (G, B)$ .

**Definition 2.5:** [1] A binary operation  $*: [0,1] \times [0,1] \rightarrow [0,1]$  is continuous  $t$ -norm if  $*$  satisfies the following conditions.

- (i)  $*$  is commutative and associative
- (ii)  $*$  is continuous
- (iii)  $a * 1 = a \quad \forall a \in [0,1]$
- (iv)  $a * b \leq c * d$  Whenever  $a \leq c, b \leq d$  and  $a, b, c, d \in [0,1]$

**Corresponding author: Manash Jyoti Borah\***  
*Department of Mathematics, Bahona College, Jorhat, Assam, India.*

**Definition 2.6:** [1] A binary operation  $\diamond: [0,1] \times [0,1] \rightarrow [0,1]$  is continuous  $t$ -conorm if  $\diamond$  satisfies the following conditions:

- (i)  $\diamond$  is commutative and associative
- (ii)  $\diamond$  is continuous
- (iii)  $a \diamond 0 = a \quad \forall a \in [0,1]$
- (iv)  $a \diamond b \leq c \diamond d$  whenever  $a \leq c, b \leq d$  and  $a, b, c, d \in [0,1]$

**Definition 2.7:** [5] Let  $U = \{x_1, x_2, x_3, \dots, x_n\}$  be the universal set of elements and  $E = \{e_1, e_2, e_3, \dots, e_m\}$  be the universal set of parameters. The pair  $(U, E)$  will be called a soft universe. Let  $F: E \rightarrow I^U$  and  $\mu$  be a fuzzy subset of  $E$ , i.e.  $\mu: E \rightarrow I = [0,1]$ , where  $I^U$  is the collection of all fuzzy subsets of  $U$ . Let  $F_\mu$  be the mapping  $F_\mu: E \rightarrow I^U \times I$  be a function defined as follows:

$F_\mu(e) = (F(e), \mu(e))$ , where  $F(e) \in I^U$ . Then  $F_\mu$  is called generalized fuzzy soft sets over the soft universe  $(U, E)$ . Here for each parameter  $e_i$ ,  $F_\mu(e_i) = (F(e_i), \mu(e_i))$  indicates not only the degree of belongingness of the elements of  $U$  in  $F(e_i)$  but also the degree of possibility of such belongingness which is represented by  $\mu(e_i)$ .

**Definition 2.8:** [5] Let  $F_\mu$  and  $G_\delta$  be two GFSS over  $(U, E)$ .

Now,  $F_\mu$  is said to be a generalized fuzzy soft subset of  $G_\delta$  if

- (i)  $\mu$  is a fuzzy subset of  $\delta$
- (ii)  $F(e)$  is also a fuzzy subset of  $G(e) \quad \forall e \in E$ .

In this case, we write  $F_\mu \subseteq G_\delta$

**Definition 2.9:** [5] Let  $F_\mu$  be a GFSS over  $(U, E)$ . Then the complement of  $F_\mu$ , denoted by  $F_\mu^c$  and is defined by  $F_\mu^c = G_\delta$ , where  $\delta(e) = \mu^c(e)$  and  $G(e) = F^c(e)$ ,  $\forall e \in E$ .

**Definition 2.10:** [5] The union of two GFSS  $F_\lambda$  and  $G_\mu$  over  $(U, E)$  is denoted by  $F_\lambda \cup G_\mu$  and defined by GFSS  $H_\delta: E \rightarrow I^U \times I$  such that for each  $e \in E$   
 $H_\delta(e) = (F(e) \diamond G(e), \lambda(e) \diamond \mu(e))$

**Definition 2.11:** [5] The intersection of two GFSS  $F_\lambda$  and  $G_\mu$  over  $(U, E)$  is denoted by  $F_\lambda \cap G_\mu$  and defined by GFSS  $H_\delta: E \rightarrow I^U \times I$  such that for each  $e \in E$   
 $H_\delta(e) = (F(e) * G(e), \lambda(e) * \mu(e))$

**Definition 2.12:** [5] A GFSS is said to be a generalized null fuzzy soft set, denoted by  $\theta_\phi$ , if  $\theta_\phi: E \rightarrow I^U \times I$  such that  $\theta_\phi(e) = (F(e), \phi(e))$  Where  $F(e) = \bar{0}$ ,  $\phi(e) = 0$ ,  $\forall e \in E$

**Definition 2.13:** [5] A GFSS is said to be a generalized absolute fuzzy soft set, denoted by  $\tilde{A}_\lambda$ , if  $\tilde{A}_\lambda: E \rightarrow I^U \times I$  such that  
 $A_\phi(e) = (F(e), \lambda(e))$  Where  $F(e) = \bar{1}$ ,  $\lambda(e) = 1$ ,  $\forall e \in E$

**Definition 2.14:** [3] Let  $X = \{x_1, x_2, x_3, \dots, x_n\}$  be the universal set of elements and  $E = \{e_1, e_2, e_3, \dots, e_m\}$  be the set of parameters. Let  $A \subseteq E$  and  $F: A \rightarrow I^U$  and  $\lambda$  be a fuzzy subset of  $A$  i.e.  $\lambda: A \rightarrow I = [0,1]$ , where  $I^U$  is the collection of all fuzzy subsets of  $U$ . Let  $F: A \rightarrow I^U \times I$  be a function defined as follows:

$F_\lambda^A(e) = (F(e), \lambda(e))$ , where  $F(e) \in I^U$ . Then  $F_\lambda^A$  is called a generalized fuzzy soft set (GFSS) over  $(U, E)$ .

Here for each parameter  $e_i$ ,  $F_\lambda^A(e_i)$  indicates not only degree of belongingness of the elements of  $U$  in  $F(e_i)$  but also degree of preference of such belongingness which is represented by  $\lambda(e_i)$ .

**Example 2.1:** Let  $U = \{S_1, S_2, S_3\}$ ,  $E = \{e_1, e_2, e_3, e_4\}$  and  $A = \{e_1, e_3, e_4\} \subseteq E$ .

We define  $F_\lambda^A$  as follows:

$$F_\lambda^A(e_1) = (\{S_1/0.3, S_2/0.5, S_3/0.7\}, 0.4),$$

$$F_\lambda^A(e_3) = (\{S_1/0.6, S_2/0.1, S_3/0.3\}, 0.5), F_\lambda^A(e_4) = (\{S_1/0.3, S_2/0.9, S_3/0.7\}, 0.8) \text{ is the generalized fuzzy soft set.}$$

**Definition 2.15:** [3] Let  $F_\lambda^A$  and  $G_\mu^B$  be two generalized fuzzy soft set over  $(U, E)$ . Now  $F_\lambda^A$  is called a generalized fuzzy soft subset of  $G_\mu^B$  if

- (i)  $A \subseteq B$ ,
- (ii)  $\lambda$  is a fuzzy subset of  $\mu$ ,
- (iii)  $\forall \lambda \in A, F(e)$  is a fuzzy subset of  $G(e)$  i.e.  $F(e) \subseteq G(e), \forall e \in E$

We write  $F_\lambda^A \subseteq G_\mu^B$

**Example 2.2:** We consider the GFSS  $F_\lambda^A$  given in **Example 2.1** and let  $B = \{e_1, e_2, e_3, e_4\} \subseteq E$

We define  $G_\mu^B$  as follows:

$$G_\mu^B(e_1) = (\{S_1/0.6, S_2/0.9, S_3/0.8\}, 0.4),$$

$$G_\mu^B(e_2) = (\{S_1/0.6, S_2/0.1, S_3/0.3\}, 0.3),$$

$$G_\mu^B(e_3) = (\{S_1/0.8, S_2/0.7, S_3/0.5\}, 0.7), G_\mu^B(e_4) = (\{S_1/0.6, S_2/0.9, S_3/0.8\}, 0.9)$$

Then  $F_\lambda^A \subseteq G_\mu^B$ .

**Definition 2.16:** [3] Let  $F_\mu^A$  be a GFSS over  $(U, E)$ . Then the complement of  $F_\mu^A$ , denoted by  $F_\mu^{A^c}$  and is defined by  $F_\mu^{A^c} = G_\delta^A$ , where  $\delta(e) = \mu^c(e)$  and  $G^A(e) = F^{A^c}(e), \forall e \in A$ .

### 3. GENARALIZED FUZZY SOFT SET REDEFINED

In this section, we put forward the notion of generalized fuzzy soft sets considering different sets of parameters and accordingly redefine the notions of null, absolute, union, intersection, complement etc. of generalized fuzzy soft sets in the following manner:

**Definition 3.1:** A generalized fuzzy soft sets (GFSS) is said to be a generalized null fuzzy soft set denoted by  $\theta_\phi^A$ , if  $\theta_\phi^A : A \rightarrow I^U \times I$  such that  $\theta_\phi^A(e) = (F(e), \phi(e))$  where  $F(e) = \bar{0}, \phi(e) = 1, \forall e \in A \subseteq E$

It is clear from our definition that the generalized fuzzy soft null set is not unique in our way, it would depend upon the set of parameters under consideration.

**Definition 3.2:** A GFSS is said to be a generalized absolute fuzzy soft set, denoted by  $\tilde{A}_\lambda$ , if  $\tilde{A}_\lambda : A \rightarrow I^U \times I$  such that  $\tilde{A}_\lambda(e) = (F(e), \lambda(e))$  Where  $F(e) = \bar{1}, \lambda(e) = 0, \forall e \in A \subseteq E$

It is clear from our definition that the generalized fuzzy soft absolute set is also not unique in our way, it would depend upon the set of parameters under consideration.

**Definition 3.3:** The union of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A \cup G_\mu^B$  and defined by GFSS  $H_\delta^{A \cup B} : A \cup B \rightarrow I^U \times I$  such that for each  $e \in A \cup B$  and  $A, B \subseteq E$

$$\text{where } H_\delta^{A \cup B}(e) = \begin{cases} (F(e), \lambda(e)), & \text{if } e \in A - B \\ (G(e), \mu(e)), & \text{if } e \in B - A \\ (F(e) \diamond G(e), \lambda(e) * \mu(e)), & \text{if } e \in A \cap B \end{cases}$$

**Example 3.1:** From Example 2.1 and 2.2

$$H_{\delta}^{A \cup B}(e_1) = (\{S_1 / 0.72, S_2 / 0.95, S_3 / 0.94\}, 0.16), H_{\delta}^{A \cup B}(e_2) = (\{S_1 / 0.60, S_2 / 0.10, S_3 / 0.30\}, 0.30), \\ H_{\delta}^{A \cup B}(e_3) = (\{S_1 / 0.92, S_2 / 0.73, S_3 / 0.65\}, 0.35), H_{\delta}^{A \cup B}(e_4) = (\{S_1 / 0.72, S_2 / 0.99, S_3 / 0.94\}, 0.72)$$

**Remark 3.1:** If we consider  $F(e) \diamond G(e) = \max\{F(e), G(e)\}$ ,  $\lambda(e) * \mu(e) = \min\{\lambda(e), \mu(e)\}$ .

Then

$$H_{\delta}^{A \cup B}(e_1) = (\{S_1 / 0.6, S_2 / 0.9, S_3 / 0.8\}, 0.4), H_{\delta}^{A \cup B}(e_2) = (\{S_1 / 0.6, S_2 / 0.1, S_3 / 0.3\}, 0.3), \\ H_{\delta}^{A \cup B}(e_3) = (\{S_1 / 0.8, S_2 / 0.7, S_3 / 0.5\}, 0.5), H_{\delta}^{A \cup B}(e_4) = (\{S_1 / 0.6, S_2 / 0.9, S_3 / 0.8\}, 0.8)$$

**Definition 3.4:** The intersection of two GFSS  $F_{\lambda}^A$  and  $G_{\mu}^B$  over  $(U, E)$  is denoted by  $F_{\lambda}^A \tilde{\cap} G_{\mu}^B$  and defined by GFSS  $K_{\delta}^{A \cap B} : A \cap B \rightarrow I^U \times I$  such that for each  $e \in A \cap B$  and  $A, B \subseteq E$   $K_{\delta}^{A \cap B}(e) = (K(e), \delta(e))$ ,

Where  $K(e) = F(e) * G(e)$ ,  $\delta(e) = \lambda(e) \diamond \mu(e)$ . In order to avoid degenerate case, we assume here that  $A \cap B \neq \varphi$ .

**Example 3.2:** From Example 2.1 and 2.2

$$K_{\delta}^{A \cap B}(e_1) = (\{S_1 / 0.18, S_2 / 0.45, S_3 / 0.56\}, 0.64), K_{\delta}^{A \cap B}(e_3) = (\{S_1 / 0.48, S_2 / 0.07, S_3 / 0.15\}, 0.85), \\ K_{\delta}^{A \cap B}(e_4) = (\{S_1 / 0.18, S_2 / 0.81, S_3 / 0.56\}, 0.98)$$

**Remark 3.2:** Let us define  $K_{\delta}^{A \cap B}(e) = (K(e), \delta(e))$ ,

where  $K(e) = \min\{F(e), G(e)\}$ ,

$$\delta(e) = \max\{\lambda(e), \mu(e)\}.$$

Then

$$K_{\delta}^{A \cap B}(e_1) = (\{S_1 / 0.3, S_2 / 0.5, S_3 / 0.7\}, 0.4), \\ K_{\delta}^{A \cap B}(e_3) = (\{S_1 / 0.6, S_2 / 0.1, S_3 / 0.3\}, 0.5), K_{\delta}^{A \cap B}(e_4) = (\{S_1 / 0.3, S_2 / 0.9, S_3 / 0.7\}, 0.8)$$

**Proposition 3.1:** If  $F_{\lambda}^A$  be a GFSS over  $(U, E)$ , then

- (i)  $F_{\lambda}^A \tilde{\cup} \theta_{\phi}^A = F_{\lambda}^A$
- (ii)  $F_{\lambda}^A \tilde{\cap} \theta_{\phi}^A = \theta_{\phi}^A$
- (iii)  $F_{\lambda}^A \tilde{\cup} \tilde{A}_{\alpha} = \tilde{A}_{\alpha}$
- (iv)  $F_{\lambda}^A \tilde{\cap} \tilde{A}_{\alpha} = F_{\lambda}^A$

**Proposition 3.2:** If  $F_{\lambda}^A, G_{\mu}^B, H_{\delta}^C$  be a GFSS over  $(U, E)$ , then

- (i)  $F_{\lambda}^A \tilde{\cap} G_{\mu}^B = G_{\mu}^B \tilde{\cap} F_{\lambda}^A$
- (ii)  $F_{\lambda}^A \tilde{\cup} G_{\mu}^B = G_{\mu}^B \tilde{\cup} F_{\lambda}^A$
- (iii)  $F_{\lambda}^A \tilde{\cap} (G_{\mu}^B \tilde{\cap} H_{\delta}^C) = (F_{\lambda}^A \tilde{\cap} G_{\mu}^B) \tilde{\cap} H_{\delta}^C$   
 $F_{\lambda}^A \tilde{\cup} (G_{\mu}^B \tilde{\cup} H_{\delta}^C) = (F_{\lambda}^A \tilde{\cup} G_{\mu}^B) \tilde{\cup} H_{\delta}^C$

**Proof:** Since the  $t$  - norm function and  $t$  - co norm functions are commutative and associative, therefore the theorem follows.

**Proposition 3.3:** If  $F_{\lambda}^A, G_{\mu}^A$  be a GFSS over  $(U, E)$ , then

- (i)  $(F_{\lambda}^A \tilde{\cap} G_{\mu}^A)^C = (G_{\mu}^A)^C \tilde{\cup} (F_{\lambda}^A)^C$
- (ii)  $(F_{\lambda}^A \tilde{\cup} G_{\mu}^A)^C = (G_{\mu}^A)^C \tilde{\cap} (F_{\lambda}^A)^C$

**Proof:** The proof is straight forward and follows from definition.

**Proposition 3.4:** The following results are valid if we take *max* and *min* operations. If  $F_\lambda^A, G_\mu^B, H_\delta^C$  be a GFSS over  $(U, E)$ , then

$$(i) F_\lambda^A \tilde{\cap} (G_\mu^B \tilde{\cup} H_\delta^C) = (F_\lambda^A \tilde{\cap} G_\mu^B) \tilde{\cup} (F_\lambda^A \tilde{\cap} H_\delta^C)$$

$$(ii) F_\lambda^A \tilde{\cup} (G_\mu^B \tilde{\cap} H_\delta^C) = (F_\lambda^A \tilde{\cup} G_\mu^B) \tilde{\cap} (F_\lambda^A \tilde{\cup} H_\delta^C)$$

**Definition 3.5:** The equal of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A = G_\mu^B$ ,

Where  $F(e) = G(e), \lambda(e) = \mu(e)$

**Proposition 3.5:** If  $F_\lambda^A, G_\mu^B, H_\delta^C$  be a GFSS over  $(U, E)$ , then  $F_\lambda^A = G_\mu^B, G_\mu^B = H_\delta^C \Leftrightarrow F_\lambda^A = H_\delta^C$

**Proof:** Since

$$F_\lambda^A = G_\mu^B, G_\mu^B = H_\delta^C$$

Therefore

$$\{F(e) = G(e), \lambda(e) = \mu(e), \{G(e) = H(e), \mu(e) = \delta(e)\}$$

Therefore

$$\{F(e) = H(e), \lambda(e) = \delta(e)\}$$

Hence

$$F_\lambda^A = H_\delta^C$$

#### 4. SOME NEW OPERATIONS OF GENERALIZED FUZZY SOFT SETS

**Definition 4.1:** Let  $F_\lambda^A$  and  $G_\mu^B$  be two GFSS over  $(U, E)$ . We define the disjunctive sum of  $F_\lambda^A$  and  $G_\mu^B$  be as the generalized fuzzy soft set  $K_\delta^C$  over  $(U, E)$ . Written as ' $F_\lambda^A \tilde{\oplus} G_\mu^B = K_\delta^C$ ',

where  $C = A \cap B \neq \emptyset, \forall e \in C$  and  $A, B \subseteq E$

$$K_\delta^C(e) = (K(e), \delta(e)),$$

Where  $K(e) = \max(F(e) * \mu(e), G(e) * \lambda(e))$

$$\delta(e) = \min(\min(G(e) \diamond \lambda(e), F(e) \diamond \mu(e)))$$

**Example 4.1:** Let  $U = \{S_1, S_2, S_3\}$  and  $E = \{e_1, e_2, e_3, e_4\}$  be the set of parameters and  $A = \{e_1, e_3, e_4\} \subseteq E$ ,  $B = \{e_1, e_2, e_3\} \subseteq E$

We define  $F_\lambda^A$  and  $G_\mu^B$  as follows:

$$F_\lambda^A(e_1) = (\{S_1 / 0.3, S_2 / 0.5, S_3 / 0.7\}, 0.4),$$

$$F_\lambda^A(e_3) = (\{S_1 / 0.6, S_2 / 0.1, S_3 / 0.3\}, 0.5), F_\lambda^A(e_4) = (\{S_1 / 0.3, S_2 / 0.9, S_3 / 0.7\}, 0.8)$$

$$G_\mu^B(e_1) = (\{S_1 / 0.6, S_2 / 0.9, S_3 / 0.8\}, 0.4), G_\mu^B(e_2) = (\{S_1 / 0.6, S_2 / 0.1, S_3 / 0.3\}, 0.3),$$

$$G_\mu^B(e_3) = (\{S_1 / 0.8, S_2 / 0.7, S_3 / 0.5\}, 0.7),$$

Then ' $F_\lambda^A \tilde{\oplus} G_\mu^B = K_\delta^C$ ' Where  $C = A \cap B = \{e_1, e_3\}$

$$K_\delta^C = \{$$

$$K_\delta^C(e_1) = (\{S_1 / \max\{0.12, 0.24\}, S_2 / \max\{0.20, 0.36\}, S_3 / \max\{0.28, 0.32\};$$

$$\min\{\min(0.76, 0.58), \min(0.94, 0.70), \min(0.88, 0.82)\})$$

$$K_{\delta}^C(e_3) = (\{S_1 / \max\{0.42, 0.40\}, S_2 / \max\{0.07, 0.35\}, S_3 / \max\{0.21, 0.25\}; \\ \min\{\min(0.90, 0.88), \min(0.85, 0.73), \min(0.75, 0.79)\})$$

i.e

$$K_{\delta}^C(e_1) = (\{S_1 / 0.24, S_2 / 0.36, S_3 / 0.32\}, 0.58)$$

$$K_{\delta}^C(e_3) = (\{S_1 / 0.42, S_2 / 0.35, S_3 / 0.25\}, 0.73)$$

**Proposition 4.1:**

Let  $F_{\lambda}^A$  and  $G_{\mu}^B$  be two GFSS over  $(U, E)$ . Then the following results hold.

- (i)  $F_{\lambda}^A \tilde{\oplus} G_{\mu}^B = G_{\mu}^B \tilde{\oplus} F_{\lambda}^A$
- (ii)  $F_{\lambda}^A \tilde{\oplus} (G_{\mu}^B \tilde{\oplus} H_{\delta}^C) = (F_{\lambda}^A \tilde{\oplus} G_{\mu}^B) \tilde{\oplus} H_{\delta}^C$

**Proof:** Since

$$F_{\lambda}^A(e) = (F(e), \lambda(e)), \text{ where } e \in E, A \subseteq E, G_{\mu}^B(e) = (G(e), \mu(e)), \text{ where } e \in E, B \subseteq E$$

$$\text{Now } (F_{\lambda}^A \tilde{\oplus} G_{\mu}^B)(e) = K_{\delta}^C(e) \text{ where } C = A \cap B, K_{\delta}^C(e) = (K(e), \delta(e))$$

$$\text{Where } K(e) = \max(F(e) * \mu(e), G(e) * \lambda(e)), \delta(e) = \min(\min(G(e) \diamond \lambda(e), F(e) \diamond \mu(e)))$$

$$\text{Let } (G_{\mu}^B \tilde{\oplus} F_{\lambda}^A)(e) = H_{\zeta}^C(e) \text{ where } C = A \cap B, H_{\zeta}^C(e) = (H(e), \zeta(e))$$

$$\text{Where } H(e) = \max(G(e) * \lambda(e), F(e) * \mu(e)) \\ = \max(F(e) * \mu(e), G(e) * \lambda(e)) = K(e)$$

$$\zeta(e) = \min(\min(F(e) \diamond \mu(e), G(e) \diamond \lambda(e))) \\ = \min(\min(G(e) \diamond \lambda(e), F(e) \diamond \mu(e))) = \delta(e)$$

Therefore

$$K_{\delta}^C(e) = H_{\zeta}^C(e)$$

Hence

$$F_{\lambda}^A \tilde{\oplus} G_{\mu}^B = G_{\mu}^B \tilde{\oplus} F_{\lambda}^A$$

Proof of (ii) can be done in a similar way.

**Proposition 4.2:**

$$F_{\lambda}^A \tilde{\oplus} \theta_{\phi}^A = F_{\lambda}^A$$

**Proof:**

$$F_{\lambda}^A(e) = (F(e), \lambda(e)), \text{ where } e \in E, A \subseteq E, \theta_{\phi}^A(e) = (\bar{0}, 1), \text{ where } e \in E, B \subseteq E$$

$$\text{Now } (F_{\lambda}^A \tilde{\oplus} \theta_{\phi}^A)(e) = K_{\delta}^A(e)$$

$$K_{\delta}^A(e) = (K(e), \delta(e))$$

$$\text{Where } K(e) = \max(F(e) * 1, 0 * \lambda(e)) = \max(F(e), 0) = F(e)$$

$$\delta(e) = \min(\min(0 \diamond \lambda(e), F(e) \diamond 1) = \min\{\min(\lambda(e), 1)\} = \lambda(e)$$

$$K_{\delta}^A(e) = (K(e), \delta(e)) = (F(e), \lambda(e)) = F_{\lambda}^A(e)$$

Hence

$$F_{\lambda}^A \tilde{\Theta} \theta_{\phi}^A = F_{\lambda}^A$$

**Definition 4.2:** Let  $F_{\lambda}^A$  and  $G_{\mu}^B$  be two GFSS over  $(U, E)$ . We define the difference of  $F_{\lambda}^A$  and  $G_{\mu}^B$  be as the generalized fuzzy soft set  $H_{\delta}^C$  over  $(U, E)$ . Written as ' $F_{\lambda}^A \tilde{\Theta} G_{\mu}^B = H_{\delta}^C$ '

where  $C = A \cap B \neq \phi, \forall e \in C$  and  $A, B \subseteq E$

$$H_{\delta}^C(e) = (H(e), \delta(e)),$$

Where  $H(e) = F(e) * \mu(e)$ ,  $\delta(e) = \max(G(e) \diamond \lambda(e))$

**Example 4.2:** From Example 4.1

$$K_{\delta}^C(e_1) = (\{S_1 / 0.12, S_2 / 0.20, S_3 / 0.28\}, \max\{0.76, 0.94, 0.88\})$$

$$K_{\delta}^C(e_3) = (\{S_1 / 0.42, S_2 / 0.07, S_3 / 0.21\}, \max\{0.90, 0.85, 0.75\})$$

i.e.

$$K_{\delta}^C(e_1) = (\{S_1 / 0.12, S_2 / 0.20, S_3 / 0.28\}, 0.94)$$

$$K_{\delta}^C(e_3) = (\{S_1 / 0.42, S_2 / 0.07, S_3 / 0.21\}, 0.90)$$

**Proposition 4.3**

$$F_{\lambda}^A \tilde{\Theta} \theta_{\phi}^A = F_{\lambda}^A$$

**Proof:**

$$F_{\lambda}^A(e) = (F(e), \lambda(e)), \text{ where } e \in E, A \subseteq E, \theta_{\phi}^A(e) = (\bar{0}, 1), \text{ where } e \in E, B \subseteq E$$

$$\text{Now } (F_{\lambda}^A \tilde{\Theta} \theta_{\phi}^A)(e) = K_{\delta}^A(e)$$

$$K_{\delta}^A(e) = (K(e), \delta(e))$$

$$\text{Where } K(e) = F(e) * 1 = F(e), \delta(e) = \max(0 \diamond \lambda(e)) = \lambda(e)$$

Therefore

$$K_{\delta}^A(e) = (K(e), \delta(e)) = (F(e), \lambda(e)) = F_{\lambda}^A(e)$$

Hence

$$F_{\lambda}^A \tilde{\Theta} \theta_{\phi}^A = F_{\lambda}^A$$

**Proposition 4.4:**

$$F_{\lambda}^A \tilde{\Theta} \tilde{A}_{\mu} = \theta_{\phi}^A$$

**Proof:**

$$F_{\lambda}^A(e) = (F(e), \lambda(e)), \text{ where } e \in E, A \subseteq E, \tilde{A}_{\mu}(e) = (1, 0)$$

$$\text{Now } (F_{\lambda}^A \tilde{\Theta} \tilde{A}_{\mu})(e) = K_{\delta}^A(e)$$

$$K_{\delta}^A(e) = (K(e), \delta(e))$$

$$\text{Where } K(e) = F(e) * \mu(e) = F(e) * 0 = 0, \delta(e) = \max(1 \diamond \lambda(e)) = 1$$

Therefore

$$K_{\delta}^A(e) = (K(e), \delta(e)) = (\bar{0}, 1) = \theta_{\phi}^A(e)$$

Hence

$$F_{\lambda}^A \tilde{\Theta} \tilde{A}_{\mu} = \theta_{\phi}^A$$

## 5. PROUCT OF GENERALIZED FUZZY SOFT SETS

**Definition 5.1:** The  $\wedge$ -Product of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A \wedge G_\mu^B$  and defined by GFSS  $H_\delta^{A \wedge B} : A \times B \rightarrow I^U \times I$  such that for each  $(\alpha, \beta) \in A \times B$  and  $A, B \subseteq E$

$$H_\delta^{A \wedge B}(\alpha, \beta) = (H(\alpha, \beta), \delta(\alpha, \beta)),$$

$$\text{Where } H(\alpha, \beta) = F(\alpha) * G(\beta), \delta(\alpha, \beta) = \lambda(\alpha) * \mu(\beta)$$

**Example 5.1:**

We define  $F_\lambda^A$  and  $G_\mu^B$  as follows:

$$F_\lambda^A(e_1) = (\{S_1/0.4, S_2/0.6, S_3/0.5\}, 0.3), F_\lambda^A(e_2) = (\{S_1/0.2, S_2/0.9, S_3/1.0\}, 0.7)$$

and

$$G_\mu^B(e_2) = (\{S_1/0.8, S_2/0.2, S_3/1.0\}, 0.4), G_\mu^B(e_3) = (\{S_1/0.2, S_2/0.3, S_3/0.6\}, 0.2)$$

Then

$$H_\delta^{A \wedge B}(e_1, e_2) = (\{S_1/0.32, S_2/0.12, S_3/0.50\}, 0.12)$$

$$H_\delta^{A \wedge B}(e_1, e_3) = (\{S_1/0.08, S_2/0.18, S_3/0.30\}, 0.06)$$

$$H_\delta^{A \wedge B}(e_2, e_2) = (\{S_1/0.16, S_2/0.18, S_3/1.00\}, 0.28)$$

$$H_\delta^{A \wedge B}(e_2, e_3) = (\{S_1/0.04, S_2/0.27, S_3/0.06\}, 0.14)$$

**Definition 5.2:** The  $\vee$ -product of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A \vee G_\mu^B$  and defined by GFSS

$$K_\delta^{A \vee B} : A \times B \rightarrow I^U \times I \text{ such that for each } (\alpha, \beta) \in A \times B \text{ and } A, B \subseteq E$$

$$K_\delta^{A \vee B}(\alpha, \beta) = (K(\alpha, \beta), \delta(\alpha, \beta)),$$

$$\text{Where } K(\alpha, \beta) = F(\alpha) \diamond G(\beta), \delta(\alpha, \beta) = \lambda(\alpha) \diamond \mu(\beta)$$

**Example 5.2:** From Example 5.1

$$K_\delta^{A \vee B}(e_1, e_2) = (\{S_1/0.88, S_2/0.68, S_3/1.00\}, 0.58)$$

$$K_\delta^{A \vee B}(e_1, e_3) = (\{S_1/0.52, S_2/0.72, S_3/0.80\}, 0.44)$$

$$K_\delta^{A \vee B}(e_2, e_2) = (\{S_1/0.84, S_2/0.92, S_3/1.00\}, 0.82)$$

**Definition 5.3:** The  $\overline{\wedge}$ -product of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A \overline{\wedge} G_\mu^B$  and defined by

GFSS  $H_\delta^{A \overline{\wedge} B} : A \times B \rightarrow I^U \times I$  such that for each  $(\alpha, \beta) \in A \times B$  and  $A, B \subseteq E$

$$H_\delta^{A \overline{\wedge} B}(\alpha, \beta) = (H(\alpha, \beta), \delta(\alpha, \beta)), \text{ Where } H(\alpha, \beta) = F(\alpha) * G^C(\beta), \delta(\alpha, \beta) = \lambda(\alpha) * \mu^C(\beta)$$

**Example 5.3:** From Example 5.1

$$H_\delta^{A \overline{\wedge} B}(e_1, e_2) = (\{S_1/0.08, S_2/0.48, S_3/0.00\}, 0.18)$$

$$H_\delta^{A \overline{\wedge} B}(e_1, e_3) = (\{S_1/0.32, S_2/0.42, S_3/0.20\}, 0.24)$$

$$H_\delta^{A \overline{\wedge} B}(e_2, e_2) = (\{S_1/0.04, S_2/0.72, S_3/0.00\}, 0.42)$$

$$H_\delta^{A \overline{\wedge} B}(e_2, e_3) = (\{S_1/0.16, S_2/0.63, S_3/0.40\}, 0.56)$$

**Definition 5.4:** The  $\overline{\vee}$ -product of two GFSS  $F_\lambda^A$  and  $G_\mu^B$  over  $(U, E)$  is denoted by  $F_\lambda^A \overline{\vee} G_\mu^B$  and defined by

GFSS  $K_\delta^{A \overline{\vee} B} : A \times B \rightarrow I^U \times I$  such that for each  $(\alpha, \beta) \in A \times B$  and  $A, B \subseteq E$

$$K_\delta^{A \overline{\vee} B}(\alpha, \beta) = (K(\alpha, \beta), \delta(\alpha, \beta)), \text{ Where } K(\alpha, \beta) = F(\alpha) \diamond G^C(\beta), \delta(\alpha, \beta) = \lambda(\alpha) \diamond \mu^C(\beta)$$



**Example 5.4:** From **Example 5.1**

$$K_{\delta}^{A \vee B}(e_1, e_2) = (\{S_1/0.52, S_2/0.92, S_3/0.50\}, 0.72)$$

$$K_{\delta}^{A \vee B}(e_1, e_3) = (\{S_1/0.88, S_2/0.88, S_3/0.70\}, 0.86)$$

$$K_{\delta}^{A \vee B}(e_2, e_2) = (\{S_1/0.36, S_2/0.98, S_3/1.00\}, 0.88)$$

$$K_{\delta}^{A \vee B}(e_2, e_3) = (\{S_1/0.84, S_2/0.97, S_3/1.00\}, 0.94)$$

**Proposition 5.1:**

If  $F_{\lambda}^A, G_{\mu}^A$  be a GFSS over  $(U, E)$ , then

$$(i) F_{\lambda}^A \wedge F_{\lambda}^A \subseteq F_{\lambda}^A$$

$$(ii) F_{\lambda}^A \vee F_{\lambda}^A \supseteq F_{\lambda}^A$$

$$(iii) F_{\lambda}^A \bar{\wedge} F_{\lambda}^A \subseteq F_{\lambda}^A$$

$$(iv) F_{\lambda}^A \bar{\vee} F_{\lambda}^A \supseteq F_{\lambda}^A$$

$$(v) (F_{\lambda}^A \vee G_{\mu}^A)^C = (G_{\mu}^A)^C \wedge (F_{\lambda}^A)^C$$

$$(vi) (F_{\lambda}^A \wedge G_{\mu}^A)^C = (G_{\mu}^A)^C \vee (F_{\lambda}^A)^C$$

$$(vii) (F_{\lambda}^A \bar{\vee} G_{\mu}^A)^C = (G_{\mu}^A)^C \bar{\wedge} (F_{\lambda}^A)^C$$

$$(viii) (F_{\lambda}^A \bar{\wedge} G_{\mu}^A)^C = (G_{\mu}^A)^C \bar{\vee} (F_{\lambda}^A)^C$$

**Proof:** The proof is straight forward and follows from definition.

## 6. CONCLUSION

In this paper, we redefined null, absolute, union, and intersection of generalized fuzzy soft sets and study some of their properties. Also we have put forward some new idea such as disjunctive sum, difference and product of two generalized fuzzy soft sets and their basic properties. It is hoped that our findings will help enhancing this study on generalized fuzzy soft sets.

## ACKNOWLEDGEMENT

The author would like to thank the fund for university Grant Commission (UGC), India, for funding the research elaborated on in this paper. [Grant No. F.5-60/2013-14/MRP/NERO/17424]

## 7. REFERENCES

- [1] B. Schweirer, A. Sklar, "Statistical metric space", Pacific Journal of Mathematics 10(1960), 314-334.
- [2] H. L. Yang, "Notes On Generalized Fuzzy Soft Sets", Journal of Mathematical Research and Exposition, Vol. 31, No. 3, May - 2011, pp.567-570.
- [3] M.J. Borah, T.J. Neog and D.K. Sut, "Some Results on Generalized Fuzzy Soft Sets", International Journal of Computer Technology and Application, Vol. 3(2), 583-591.
- [4] P. K. Maji, R. Biswas and A.R. Roy, "Fuzzy Soft Sets", Journal of Fuzzy Mathematics, Vol. 9, no.3, pp.-589-602, 2001.
- [5] P. Majumdar, S. K. Samanta, "Generalized Fuzzy Soft Sets", Computers and Mathematics with Applications, 59(2010), pp.1425-1432.
- [6] T. J. Neog, D. K. Sut, "On Union and Intersection of Fuzzy Soft Sets", Int.J. Comp. Tech. Appl., Vol. 2 (5), 1160-11.
- [7] T. J. Neog, D. K. Sut, "On Fuzzy Soft Complement and Related Properties", Accepted for publication in International Journal of Energy, Information and communications (IJEIC), Japan.

**Source of support: University Grant Commission (UGC), India, Conflict of interest: None Declared**

**[Copy right © 2014. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**