# **COMPACT STANDARD FUZZY METRIC SPACE**

# Jehad R. Kider\*

Department of Applied Sciences, University of Technology, Iraq.

(Received On: 25-06-14; Revised & Accepted On: 18-07-14)

# ABSTRACT

In this paper we recall the definition of standard fuzzy metric space then we define a compact standard fuzzy metric space and F-totally bounded after that we prove that F-totally bounded complete standard fuzzy metric space is compact. Moreover we recall the definition of continuous and uniform continuous function to prove that continuous function and uniform continuous functions are equivalent on a compact standard fuzzy metric spaces.

*Key Words:* Standard fuzzy metric space, compact standard fuzzy metric space, F-bounded, F-totally bounded, continuous function.

# INTRODUCTION

Theory of fuzzy sets was introduced by Zadeh in 1965 [1]. Many authors have introduced the concept of fuzzy metric in different ways [2,3, 4 and 5]. Kramosil and Michalek in 1975 [6] introduced the definition of fuzzy metric space which is called later KM-fuzzy metric space .George and Veeramani in 1994[3] introduced the definition of continuous \* t-norm to modify the concept of KM-fuzzy metric space which was introduced by Kramosil and Michalek which is called later GV-fuzzy metric space.

In section one of this paper we recall the definition of standard fuzzy metric space [9] which is a modification of the definition GV-fuzzy metric space after that we introduce basic definitions, basic concepts and properties of standard fuzzy metric space.

In section two the notion of compact standard fuzzy metric space is introduced, we try to prove results similar to that in the ordinary case.

The aim of studying a continuous function on compact spaces in section three is to prove that continuous function and uniform continuous functions are equivalent on compact standard fuzzy metric space.

# 1. STANDARD FUZZY METRIC SPACE

**Definition 1.1:** [3] A binary operation  $*: [0,1] \times [0,1] \rightarrow [0,1]$  is a continuous t-norm if \* satisfies the following conditions:

- 1. \* is associative and commutative.
- 2. \* is continuous.
- 3. a\*1 = a for all  $a \in [0,1]$ .
- 4.  $a*b \le c*d$  whenever  $a \le c$  and  $b \le d$  where  $a,b, c,d \in [0,1]$ .

**Remark 1.2:** [3] For any  $r_1 > r_2$  we can find  $r_3$  such that  $r_1 * r_3 \ge r_2$  and for any  $r_4$  we can find an  $r_5$  such that  $r_5 * r_5 \ge r_4$  where  $r_1, r_2, r_3, r_4, r_5 \in (0, 1)$ .

Corresponding author: Jehad R. Kider\* Department of Applied Sciences, University of Technology, Iraq. We introduce the following definition.

**Definition 1.3:[9]** A triple (X, M,\*) is said to be standard fuzzy metric space if X is an arbitrary set, \* is a continuous t- norm and M is a fuzzy set on X<sup>2</sup> satisfying the following conditions:

 $\begin{array}{ll} (FM_1) \ M(x,\,y) > 0 \ for \ all \ x,\,y \in X \\ (FM_2) \ M(x,\,y) = 1 \ if \ and \ only \ if \ x = y \\ (FM_3) \ M(x,\,y) = M(y,\,x) \ for \ all \ x,\,y \in X \\ (FM_4) \ M(x,z) \geq M(x,y) \ *M(y,z) \ for \ all \ x,\,y \ and \ z \in X \\ (FM_5) \ M(x,\,y) \ is \ a \ continuous \ fuzzy \ set \end{array}$ 

**Example 1.4:** [9] Let  $X = \mathbb{N}$ , and let a \* b = a.b for all  $a, b \in [0, 1]$ .

Define  $M(x, y) = \begin{cases} \frac{x}{y} & \text{if } x \le y \\ y & \text{for all } x, y \in \mathbb{N}. \end{cases}$  $\frac{y}{x} & \text{if } y \le x \end{cases}$ 

Then  $(\mathbb{N}, \mathbb{M}, .)$  is a standard fuzzy metric space.

**Example 1.5:** [9] Let  $X=\mathbb{R}$  and let a\*b = a.b for all  $a, b \in [0, 1]$ .

Define  $M(x, y) = \frac{1}{e^{|x-y|}}$  for all  $x, y \in \mathbb{R}$ .

Then  $(\mathbb{R}, M, .)$  is a standard fuzzy metric space.

**Definition 1.6:** [9] Let (X, M, \*) be a standard fuzzy metric space then M is continuous if whenever  $x_n \rightarrow x$  and  $y_n \rightarrow y$  in X then  $M(x_n, y_n) \rightarrow M(x, y)$  that is  $\lim_{n \rightarrow \infty} M(x_n, y_n) = M(x, y)$ .

**Definition 1.7:** [9] Let (X, M, \*) be a standard fuzzy metric space .Then  $B(x, r) = \{y \in X: M(x, y) > 1 - r\}$  is an open ball with center  $x \in X$  and radius r, 0 < r < 1.

**Proposition 1.8:** [9] Let  $B(x,r_1)$  and  $B(x,r_2)$  be two open balls with same center x in a standard fuzzy metric space (X,M,\*). Then either  $B(x,r_1) \subseteq B(x,r_2)$  or  $B(x,r_2) \subseteq B(x,r_1)$  where  $r_1,r_2 \in (0,1)$ .

**Definition 1.9:[9]** A subset A of a standard fuzzy metric space (X, M,\*) is said to be open if given any point a in A there exists r, 0 < r < 1 such that B(a, r)  $\subseteq$  A. A subset B is said to be closed if B<sup>c</sup> is open.

Theorem 1.10: [9] Every open ball in a standard fuzzy metric space (X, M,\*) is an open set.

**Theorem 1.11:** [9] Let (X, M,\*) is a standard fuzzy metric space. Define  $\tau_M = \{A \subset X : x \in A \text{ if and only if there exists } 0 < r < 1 \text{ such that } B(x, r) \subset A\}$  Then  $\tau_M$  is a topology on X.

Theorem 1.12: [9] Every standard fuzzy metric space is a Hausdorff space.

**Definition 1.13:[9]** A sequence  $(x_n)$  in a standard fuzzy metric space (X,M,\*) is said to be converge to a point x in X if for each r, 0 < r < 1 there exists a positive number N such that  $M(x_n, x) > (1-r)$ , for each  $n \ge N$ .

**Theorem 1.14:** [9] Let (X, M, \*) be a standard fuzzy metric space then for a sequence  $(x_n)$  in X converge to x if and only if  $\lim_{n\to\infty} M(x_n, x) = 1$ .

**Definition 1.15:** [9] A sequence  $(x_n)$  in a standard fuzzy metric space (X,M,\*) is Cauchy if for each r, 0 < r < 1, there exists a positive number N such that  $M(x_n,x_m) > (1-r)$ , for each m,  $n \ge N$ .

**Proposition 1.16:** [9] Let (X, d) be an ordinary metric space and let a\*b = a.b for all  $a, b \in [0,1]$ . Define  $M_d(x, y) = \frac{1}{1+d(x,y)}$ , then  $(X, M_d, *)$  is a standard fuzzy metric space and it is called the standard fuzzy metric induced by the metric d.

**Proposition 1.17:** [9] Let (X, d) be a metric space and let  $(X,M_d,*)$  be the standard fuzzy metric space induced by d. Let  $(x_n)$  be a sequence in X. Then  $(x_n)$  converges to  $x \in X$  in (X, d) if and only if  $(x_n)$  converges to x in  $(X,M_d,*)$ . **Proposition 1.18:** [9] Let (X, d) be a metric space and let  $M_d(x, y) = \frac{1}{1+d(x,y)}$ . Then  $(x_n)$  is a Cauchy sequence in (X, d) if and only if  $(x_n)$  is a Cauchy sequence in (X,  $M_d$ ,\*).

**Definition 1.19:** [9] Let (X, M, \*) be a standard fuzzy metric space. A subset A of X is said to be F-bounded if there exists 0 < r < 1 such that, M(x, y) > 1- r, for all x,  $y \in A$ .

**Proposition 1.20:** [9] Let (X, d) be a metric space and let  $M_d(x, y) = \frac{1}{1+d(x,y)}$  then a subset A of X is F-bounded if and only if it is bounded.

**Definition1.21:** [9] A standard fuzzy metric space (X, M,\*) is complete if every Cauchy sequence in X converges to a point in X.

**Definition 1.21:** [9] Let  $(X, M_X, *)$  and  $(Y, M_Y, *)$  be standard fuzzy metric spaces and  $A \subseteq X$ . A function f:  $A \rightarrow Y$  is said to be continuous at  $a \in A$ , if for every  $0 < \varepsilon < 1$ , there exist some  $0 < \delta < 1$ , such that  $M_Y(f(x), f(a)) > (1 - \varepsilon)$  whenever  $x \in A$  and  $M_X(x, a) > (1 - \delta)$ . If f is continuous at every point of A, then it is said to be continuous on A.

**Theorem 1.22:** [9] Let  $(X, M_X, *)$  and  $(Y, M_Y, *)$  be standard fuzzy metric spaces and A $\subseteq X$ . A function f: A $\rightarrow Y$  is continuous at  $a \in A$  if and only if whenever a sequence  $(x_n)$  in A converge to a, the sequence  $(f(x_n))$  converges to f(a).

**Theorem 1.23:[9]** A function  $f: X \rightarrow Y$  is continuous on X if and only if  $f^{-1}(G)$  is open in X for all open subset G of Y.

**Theorem 1.24:** [9] A mapping f:  $X \rightarrow Y$  is continuous on X if and only if  $f^{-1}(F)$  is closed in X for all closed subset F of Y.

### 2. COMPACTNESS

**Definition 2.1:** Let (X, M, \*) be a standard fuzzy metric space and  $Y \subseteq X$ . Let  $\widehat{G}$  be a collection of open sets in X with the property that  $Y \subseteq \bigcup_{G \in \widehat{G}} G$ . Equivalently, for each  $x \in Y$  there is  $G \in \widehat{G}$  such that  $x \in G$ . Then  $\widehat{G}$  is called an open cover or an open covering of Y. A finite subcollection of  $\widehat{G}$ . which itself a cover is called a finite subcover or a finite subcovering of Y.

**Definition 2.2:** A standard fuzzy metric space (X,M,\*) is said to be compact if every open covering  $\widehat{G}$  of X has a finite subcovering that is there is a finite subcollection  $\{G_1, G_2, G_3, ..., G_n\} \subseteq \widehat{G}$  such that  $X = \bigcup_{i=1}^n G_i$ .

**Definition 2.3:** A nonempty subset Y of X is said to be compact if it is compact with the standard fuzzy metric induced on it by M.

**Example 2.4:** The interval (0, 1) in the standard fuzzy metric space ( $\mathbb{R}$ , M,\*) where M(x, y) =  $\frac{1}{1+|x-y|}$ , a\*b = a.b for all a, b  $\in$  [0,1], is not compact. In order to prove the assertion suffices to exhibit an open covering from which no finite subcovering can be selected. The open covering {( $\frac{1}{n}$ , 1): n = 2, 3...} is one such covering of (0, 1), from which no finite subcovering can be selected.

Remark 2.5: Let Y be a finite subset of a standard fuzzy metric space (X, M,\*). Then Y is compact.

**Definition 2.6:** A collection F of sets in X is said to have the finite intersection property if every finite subcollection of F has a nonempty intersection.

**Definition 2.7:** Let S be a subset of X where (X,M,\*) is a standard fuzzy metric space then S is called F-totally bounded if for each  $0 < \varepsilon < 1$ , there is a finite set of points  $\{y_1, y_2, y_3, \dots, y_n\} \subset S$  such that whenever x in X,  $M(x, y_i) > (1 - \varepsilon)$  for some  $y_i \in \{y_1, y_2, y_3, \dots, y_n\}$ . This set of points  $\{y_1, y_2, y_3, \dots, y_n\}$  is called F- $\varepsilon$  - net.

Proposition 2.8: An F-totally bounded standard fuzzy metric space is F-bounded.

**Proof:** Let (X, M, \*) be F-totally bounded and suppose  $0 < \varepsilon < 1$  is given. Then there exists a finite F- $\varepsilon$ - net for X, say A. Since A is a finite set of points 0 < M(A) < 1, where  $M(A) = \sup\{M(y,z): y, z \in A\}$ . Now let  $x_1$  and  $x_2$  be any two points of X. There exists points y and z in A such that  $M(x_1, y) > (1 - \varepsilon)$  and  $M(x_2, z) > (1 - \varepsilon)$ . Now for M(A) and  $\varepsilon$  there is 1- r, where 0 < r < 1 such that  $M(A) * (1 - \varepsilon) * (1 - \varepsilon) \ge (1 - r)$ . It follows that  $M(x_2, y) > M(y, y) * M(y, z) * M(y, z)$ 

 $M(x_1, x_2) \ge M(x_2, y) * M(y, z) * M(z, x_2)$  $\ge (1 - \varepsilon) * M(A) * (1 - \varepsilon) \ge (1 - r)$ 

So,  $M(X) = \sup\{M(x_1, x_2): x_1, x_2 \in X\} \ge (1 - r)$ . Hence, X is F-bounded.

© 2014, IJMA. All Rights Reserved

**Theorem 2.9:** Let Y be a subset of a standard fuzzy metric space (X, M,\*). Then Y is F-totally bounded if and only if every sequence in Y contains a Cauchy subsequence.

**Proof:** Suppose Y is F- totally bounded. Let  $(y_n)$  be a sequence in Y whose range may be assumed to be infinite. Choose a finite F- $\frac{1}{2}$ -net in Y. Then one of the balls of radius  $\frac{1}{2}$  with center in the F- $\frac{1}{2}$ -net contains infinitely many elements of the range of the sequence. We shall denote the subsequence formed by these elements by  $(y_n^{(1)})$ . Choose a finite F- $\frac{1}{4}$ -net in Y. Then one of the balls of radius  $\frac{1}{4}$  with center in the finite F- $\frac{1}{4}$ -net contains infinitely many elements of the range of  $(y_n^{(1)})$ .

We shall denote the subsequence formed as  $(y_n^{(2)})$ . Proceeding in this way, we obtain a sequence of sequences, each a subsequence of the preceding one, so that at the kth stage, the terms  $(y_n^{(k)})$  lie in the ball of radius  $\frac{1}{2^k}$  with center in the F- $\frac{1}{2^k}$ -net.

Now  $(y_n^{(n)})$  is a subsequence of  $(y_n)$ . Let  $0 < \epsilon < 1$  be given. Choose N so large that  $(1 - \frac{1}{2^n}) * (1 - \frac{1}{2^{n+1}}) * \dots * (1 - \frac{1}{2^{m-1}}) > (1 - \epsilon)$ .

Then for m > n > N, we have  $M(y_n^{(n)}, y_m^{(m)}) \ge M(y_n^{(n)}, y_{n+1}^{(n+1)}) * M(y_{n+1}^{(n+1)}, y_{n+2}^{(n+2)}) * \dots * M(y_{m-1}^{(m-1)}, y_m^{(m)})$   $\ge (1 - \frac{1}{2^n}) * (1 - \frac{1}{2^{n+1}}) * \dots (1 - \frac{1}{2^{m-1}})$   $> (1 - \varepsilon)$ 

So that the sequence  $(y_n^{(n)})$  is a Cauchy sequence.

Conversely, suppose that every sequence in Y has a Cauchy subsequence

We shall show that Y is F-totally bounded. Let  $0 < \varepsilon < 1$  and let  $y_1 \in Y$ . If  $Y-B(y_1,\varepsilon) = \emptyset$ , we have found an F- $\varepsilon$ -net, namely, the set  $\{y_1\}$ . Otherwise choose  $y_2 \in Y$ -B $(y_1,\varepsilon)$ . If  $Y-[B(y_1,\varepsilon) \cup B(y_2,\varepsilon)] = \emptyset$ . We have found an F- $\varepsilon$ -net, namely the set  $\{y_1,y_2\}$ . It is enough to show that this process terminates, after a finite number of steps. If it dose not terminate, we shall obtain an infinite sequence  $(y_n)$  with property that  $M(y_n, y_m) \le (1-\varepsilon)$ ,  $n \ne m$ . Consequently, the sequence  $(y_n)$  would have no Cauchy subsequence, contrary to hypothesis.

We give below a characterization of compact standard fuzzy metric space.

Proposition 2.10: Let (X, M,\*) be a compact standard fuzzy metric space. Then X is F-totally bounded.

**Proof:** For any given  $0 < \varepsilon < 1$ , the collection of all balls  $B(x,\varepsilon)$  for  $x \in X$  is an open cover of X. The compactness of X implies that this open cover contains a finite sub cover. Hence for  $0 < \varepsilon < 1$ , X is covered by a finite number of open balls of radius  $\varepsilon$  i.e the centers of the balls in the finite subcover form a finite F- $\varepsilon$ -net for X. So, X is F-totally bounded.

**Proposition 2.11:** Let (X, M,\*) be a compact standard fuzzy metric space. Then (X, M,\*) is complete.

**Proof:** Suppose, if possible, that (X, M, \*) is a compact standard fuzzy metric space is not complete. Then there exists a Cauchy sequence  $(x_n)$  in (X,M, \*) not having a limit in X. Let  $y \in X$ , since  $(x_n)$  dose not converge to y there exists 0 < r < 1. Such that  $M(x_n, y) \le (1 - r)$  for infinitely many values of n, since  $(x_n)$  is Cauchy, there exists an integer N such that n,  $m \ge N$ . That implies  $M(x_n, x_m) > (1 - \varepsilon)$ . Choose  $m \ge N$  for which  $M(x_m, y) > (1 - \varepsilon)$ . So, the open ball  $B(y,\varepsilon)$  contains  $x_n$  for only finitely many values of n. In this manner, we can associate with each  $y \in X$  a ball  $B(y,\varepsilon(y))$ , where  $0 < \varepsilon(y) < 1$  depends on y, and the ball  $B(y, \varepsilon(y))$  contains  $x_n$  for only finitely many values of n. Observe that

 $X = \bigcup_{y \in X} B(y, \varepsilon(y))$  which means that  $\{B(y, \varepsilon(y)): y \in X\}$  is a covering of X. Since X is compact there exists a finite subcovering  $B(y_i, \varepsilon(y_i)), i = 1, 2, ..., n$ .

Since each ball contains  $x_n$  for only a finite number of values of n, therefore the balls are in the finite subcovering and, hence, also X, must contains  $x_n$  for only a finite number of values of n. This however, is impossible. Hence (X, M,\*) must be complete

**Theorem 2.12:** Let (X, M,\*) be F-totally bounded and complete standard fuzzy metric space. Then (X, M,\*) is compact.

**Proof:** Suppose, if possible, that (X, M, \*) is F-totally bounded and complete but not compact. Then there exists an open covering  $\{G_{\lambda}: \lambda \in \Lambda\}$  of X that dose not admit a finite subcovering. Since (X, M, \*) is F-totally bounded, it is F-bounded, hence for some 0 < r < 1 and some  $x \in X$ , we have  $X \subseteq B(x, r)$ . Observe that  $X \subseteq B(x, r)$  implies X = B(x, r).

Let  $\varepsilon_n = \frac{r}{2^n}$ . We know that X being F-totally bounded can be covered by finite many balls of radius  $\varepsilon_1$ . By our hypothesis at least one of these balls, say B(x<sub>1</sub>, $\varepsilon_1$ ), cannot be covered by a finite number of sets G<sub> $\lambda$ </sub>.

[for if each had a finite subcovering, the same would be true for X]

Because  $B(x_1, \varepsilon_1)$  is itself F-totally bounded [any nonempty subset of F-totally bounded set is F-totally bounded], we can find an  $x_2 \in B(x_1, \varepsilon_1)$  such that  $B(x_2, \varepsilon_2)$  cannot be covered by a finite number of sets  $G_{\lambda}$ .

In this way, a sequence  $(x_n)$  may be defined with the property that for each n,  $B(x_n, \varepsilon_n)$  cannot be covered by a finite number of sets  $G_{\lambda}$  and  $x_{n+1} \in B(x_n, \varepsilon_n)$ . We next show that the sequence  $(x_n)$  is convergent.

Since  $x_{n+1} \in B(x_n, \varepsilon_n)$  it follows that  $M(x_n, x_{n+1}) > (1 - \varepsilon_n)$ .

Let  $0 < \varepsilon < 1$  such that  $(1 - \varepsilon_n) * (1 - \varepsilon_{n+1}) * \dots * (1 - \varepsilon_m) > (1 - \varepsilon)$ 

Hence

$$\begin{split} M(x_n, x_m) &\geq M(x_n, x_{n+1}) * \dots * M(x_{m-1}, x_m) \\ &\geq (1 \text{-} \epsilon_n) * (1 \text{-} \epsilon_{n+1}) * \dots * (1 \text{-} \epsilon_m) \\ &> (1 \text{-} \epsilon) \end{split}$$

So  $(x_n)$  is a Cauchy sequence in X and since X is complete, it converges to  $y \in X$ , say. Since  $y \in X$  there exists  $\lambda_0 \in \Lambda$  such that  $y \in G_{\lambda_0}$ . Because  $G_{\lambda_0}$  is open it contains  $B(y,\delta)$  for some  $0 < \delta < 1$ . Choose N so large that,  $M(x_n,y) > (1-\delta)$  and  $(1-\varepsilon_n) > (1-\delta)$ . Then, for any  $x \in X$  such that

 $M(x, x_n) > (1 - \varepsilon_n)$ . It follows that

 $M(x, y) \ge M(x, x_n) * M(x_n, y)$  $\ge (1 - \delta) * (1 - \delta) > (1 - r),$ 

for some 0 < r < 1. So that  $B(x_n, \varepsilon_n) \subseteq B(y, r)$ . Therefore  $B(x_n, \varepsilon_n)$  admits a finite subcovering, namely by the set  $G_{\lambda_0}$ .

Since this contradicts  $B(x_n, \varepsilon_n)$  cannot be covered by a finite number of sets  $G_{\lambda}$ , the proof is complete.

Theorem 2.13: A standard fuzzy metric space is compact if and only if it is complete and F-totally bounded.

**Theorem 2.14:** (X, d) is a compact metric space if and only if  $(X, M_d, *)$  is a compact standard fuzzy metric space where  $M_d(x,y) = \frac{1}{1+d(x,y)}$ .

**Proof:** Suppose that (X, d) is compact. Let  $(x_n)$  be a sequence in  $(X,M_d,*)$  then  $(x_n)$  is a sequence in (X,d). But (X,d) is compact hence  $(x_n)$  has a convergent subsequence. Then  $(x_n)$  has a convergent sequence in  $(X,M_d,*)$  by Proposition 1.17. Hence  $(X,M_d,*)$  is compact. In similar way we can prove that if  $(X,M_d,*)$  is compact then (X, d) is compact by using Proposition 1.17.

**Proposition 2.15:** Let (X, M,\*) be a standard fuzzy metric space. Then the following tatements are equivalent:

- (i) Every infinite set in (X, M,\*) has at least one limit point in X.
- (ii) Every infinite sequence in (X, M,\*) contains a convergent subsequence.

**Proof:** (i) $\Rightarrow$ (ii): Let  $(x_n)$  be a sequence in X. If the set  $\{x_1, x_2, ...,\}$  is finite, then one of the points, say  $x_{i_0}$ , satisfies  $x_{i_0} = x_j$ , for infinitely many  $j \in \mathbb{N}$ . Hence the constant sequence  $(x_{i_0})$  is a subsequence of  $(x_n)$ , which converges to the point  $x_{i_0}$ . Suppose that the set  $\{x_1, x_2, ...,\}$  is infinite.

In view of (i), the infinite set  $\{x_1, x_2, ...,\}$  has at least one limit point  $x \in X$ . A subsequence of  $(x_n)$  that converges to x may be obtained as follows: Let  $n_1$  be any integer such that  $M(x_{n_1}, x) > 0$ .

Having defined  $n_k$ , let  $n_{k+1}$  be the smallest integer such that  $n_{k+1} > n_k$  and  $M(x_{n_{k+1}},x) > 1 - \frac{1}{k+1}$ . Then the sequence  $(x_{n_k})$  converges to x.

**Proof (ii)** $\Rightarrow$ (i): Let Y be an infinite subset of X. Then there exists a sequence  $(y_n)$  in X of distinct terms. In view of (ii)  $(y_n)$  contains a subsequence  $(y_{n_i})$  of distinct terms that converges to  $y \in X$ . Hence every open ball with center y contains an infinite number of terms of the convergent subsequence  $(y_{n_i})$ . But the terms are distinct; hence every open ball centered at y contains an infinite number of points of Y.  $y \in X$  is a limit point of Y.

**Theorem 2.16:** The standard fuzzy metric space (X, M,\*) is compact if and only if every sequence of points in X has a subsequence converging to a point in X.

**Proof:** Suppose first that X is compact (equivalently, F-totally bounded and complete see Theorem 2.13) and that  $(x_n)$  is any sequence of points in X. Since X is F-totally bounded, it follows, using Theorem 2.13, that  $(x_n)$  contains a Cauchy subsequence  $(x_{n_i})$ . But  $(x_{n_i})$  converges to a point  $x \in X$  because X is complete. Thus, if X is complete, then every sequence in X contains a convergent subsequence.

Conversely, suppose every sequence in X has a convergent subsequence, it follows in view of the fact that every convergent sequence is Cauchy and Theorem 2.13 that X is F-totally bounded. It remains to show that X is complete. To this end let  $(x_n)$  be a Cauchy sequence in X. By assumption  $(x_n)$  has a subsequence  $(x_{n_i})$  that converges to a point  $x \in X$ . We shall show that  $x_n \rightarrow x$ . Let  $0 < \varepsilon < 1$  be given by Remark 1.2, there is 0 < r < 1 such that  $(1-r) *(1-r) > (1-\varepsilon)$ .

Now  $x_{n_i} \rightarrow x$ , there exist  $N_1$  such that  $M(x_{n_i}, x) > (1 - r)$  for all  $n_i \ge N_1$ .

Since the sequence  $(x_n)$  is Cauchy, there exists  $N_2$  such that

 $M(x_n,x_m) > (1-r) \text{ for all } m, n \ge N_2.$ 

Let N = minN<sub>1</sub>, N<sub>2</sub>} then M(x<sub>n</sub>,x)  $\ge$  M(x<sub>n</sub>,x<sub>n<sub>i</sub></sub>) \* M(x<sub>n<sub>i</sub></sub>,x) > (1-r) \* (1-r) > (1-ε)

For all  $n \ge N$ . This complete the proof.

The results of this section can be summed up as follows:

**Theorem 2.17:** Let (X, M,\*) be a standard fuzzy metric space. The following statements are equivalent:

- a. (X,M,\*) is compact
- b. (X,M,\*) is complete and F-totally bounded
- c. Every infinite set in X has at least one limit point
- d. Every sequence in X contains a convergent subsequence.

Corollary 2.18: Let Y be a closed subset of the compact standard fuzzy metric space (X, M,\*). Then Y is compact.

**Proof:** Let  $(y_n)$  be a sequence of points in Y. Then  $(y_n)$  considered as sequence of points in X, has a subsequence converging to a point  $x \in X$ . But then  $x \in Y$  since Y is closed. Thus any sequence in Y has a subsequence converging to a point in Y. By Theorem 2.13 Y is compact.

**Theorem 2.19:** Let Y be a subset of a standard fuzzy metric space (X, M,\*). If Y is compact then Y is a closed subset of X.

**Proof:** Let  $x \in X$  be a limit point of Y. Then there is a sequence  $(y_n)$  in Y converging to x. But then  $(y_n)$  is a Cauchy sequence in Y. Since Y is complete, $(y_n)$  converges to a point y in Y. This point y must be x and so  $x \in Y$ . Thus Y contains all its limit points and is therefore closed.

### 3. CONTINUOUS FUNCTION ON COMPACT SPACE

**Theorem 3.1:** Let f be a continuous function from a compact standard fuzzy metric space  $(X, M_X, *)$  into a standard metric space  $(Y, M_Y, *)$ . Then the range f(X) of f is also compact.

**Proof:** Let  $\{G_{\lambda}: \lambda \in \Lambda\}$  be an open covering of f(X). Since f is continuous  $f^{-1}(G_{\lambda})$  is open in X. Moreover  $\{f^{-1}(G_{\lambda}):\lambda \in \Lambda\}$  is an open covering of X. Since X is compact, there exists  $\lambda_1, \lambda_2, \lambda_3, ..., \lambda_n$  in  $\Lambda$  such that  $X = \bigcup_{i=1}^{n} f^{-1}(G_{\lambda_i})$ .

Now

 $f(X) = f(\bigcup_{i=1}^{n} f^{-1}(G_{\lambda_i})) = \bigcup_{i=1}^{n} f(f^{-1}(G_{\lambda_i})) \subseteq \bigcup_{i=1}^{n} G_{\lambda_i}$ 

So  $\{G_{\lambda_i}, i = 1, 2, ..., n\}$  is a finite sub covering of f(X). Consequently, f(X) is compact

**Corollary 3.2:** Let f be a homeomorphism of a standard fuzzy metric space  $(X,M_X,*)$  onto a standard fuzzy metric space  $(Y,M_Y,*)$ . Then X is compact if and only if Y is compact.

**Corollary 3.3:** Let f be a continuous function from a compact standard fuzzy metric space  $(X, M_X, *)$  onto a standard fuzzy metric space  $(Y, M_Y, *)$ . Then f(X) is F-bounded and closed subset of Y.

**Theorem 3.4:** If f is a one-to-one continuous mapping of a compact standard fuzzy metric space  $(X,M_X,*)$  onto a standard fuzzy metric space  $(Y,M_Y,*)$  then  $f^{-1}$  is continuous on Y and, hence, f is homeomorphism of  $(X,M_X,*)$  onto  $(Y,M_Y,*)$ .

**Proof:** Suppose f: X→Y is one-to-one and onto. Its inverse  $f^{-1}$ :Y→X is well defined. Let F be a closed subset of X. By Theorem 2.19, F is compact. By Theorem 3.1 f(F) is compact and hence, a closed subset of Y by Theorem 2.19. But  $f(F) = (f^{-1})^{-1}(F)$  and so  $(f^{-1})^{-1}(F)$  is closed in Y. Hence by Theorem 1.24  $f^{-1}$  is continuous.

**Definition 3.5:** Let  $\{G_{\lambda}: \lambda \in \Lambda\}$  be an open covering of the standard fuzzy metric space (X,M,\*). Any number  $0 < \delta < 1$  such that for each  $x \in X$  there exists  $\lambda \in \Lambda$  (dependent on x) for which  $B(x,\delta) \subseteq G_{\lambda}$  is called a Lebesgue number of the covering  $\{G_{\lambda}: \lambda \in \Lambda\}$ .

**Theorem 3.6:** Let  $(X, M_X, *)$  be a compact standard fuzzy metric space,  $(Y, M_Y, *)$  be an arbitrary standard fuzzy metric space and f:  $X \rightarrow Y$  be continuous.

Then for each  $0 < \varepsilon < 1$ , there exists a  $\delta$ ,  $0 < \delta < 1$  ( $\delta$  depending on  $\varepsilon$  only) such that  $f(B(x,\delta)) \subseteq B(f(x),\varepsilon)$  for every  $x \in X$ . That is f is uniformly continuous on X.

**Proof:** Let 0 < r < 1 such that  $(1 - r) * (1 - \epsilon)$ . The collection of ball {B(y, r):  $y \in Y$ } constitutes an open cover of Y. The set { $f^{-1}(B(y, r))$ :  $y \in Y$ } therefore form an open cover of the compact standard fuzzy metric space X. Let  $\delta$  be a Lebesgue number of this open cover of X.

Since each open ball  $B(x,\delta)$  lies in one of these sets,

 $f(B(x,\delta)) \subseteq B(y, r)$  for some  $y \in Y$ .

Because  $f(x) \in B(y, r)$ , we find for any  $z \in B(x, \delta)$  that

$$\begin{split} M_{Y}(f(z), f(x)) &\geq M_{Y}(f(z), y) * M_{Y}(y, f(x)) \\ &> (1 - r) * (1 - r) > (1 - \epsilon) \end{split}$$

i.e,  $f(B(x,\delta)) \subseteq B(f(x),\varepsilon)$  this complete the proof.

### REFERENCES

[1] Zadeh L.A, Fuzzy Sets, Information and Control, Vol. 8(1965) 338 - 353.

[2] Sisodia, K. S., Rathore, M. S. Singh, D. and Khichi, S. S., A Common Fixed Point Theorem in Fuzzy Metric Spaces, Int. J. Math Analysis, Vol. 5, No.17 (2011)810 – 826.

[3] George, A. and Veeramani, P. On some results in fuzzy metric Spaces, Fuzzy Systems, Vol. 64(1994)395-399.

[4] George, A. and Veeramani, P. On some results of analysis for fuzzy metric spaces, Fuzzy Sets and Systems, Vol. 90(1997)365 – 368.

[5] Lopez, R. J and Romaguera, S., and Alvarez, J. M., The Hausdorff Fuzzy Quasi Metric, Fuzzy Sets and Systems, Vol. 161(2010)1078-1096.

[6] Kramosil, O. and Michalek, J. Fuzzy metrics and statistical metric Spaces, Kybernetika, Vol. 11(1975)326 - 334.

[7] Mihet, D. A Banach contraction theorem in fuzzy metric spaces, Fuzzy Sets and Systems, Vol. 144(2004)431 – 439.

#### © 2014, IJMA. All Rights Reserved

[8] Miheţ, D. On fuzzy contractive mappings in fuzzy metric spaces, Fuzzy Sets and Systems, Vol. 158(2007) 915-921.

[9] Hussain,Z. A., On Some Results of Analysis in a Standard fuzzy Metric Spaces, M.Sc. Thesis, University of Technology, 2013.

# Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2014. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]