GEOMETRIC MEAN LABELING ON DOUBLE TRIANGULAR SNAKES

¹S. S. Sandhya^{*} and ²S. Somasundaram

¹Department of Mathematics, Sree Ayyappa College for Women, Chunkankadai-629003, India.

²Department Mathematics, M.S. University, Tirunelveli – 627 012, India.

(Received on: 20-06-14; Revised & Accepted on: 16-07-14)

ABSTRACT

A Graph G = (V, E) with p vertices and q edges is called a Geometric mean graph if it is possible to label the vertices $x \in V$ with distinct labels f(x) from 1, 2.....q+1 in such a way that when each edge e=uv is labeled with $f(e = vu) = \left[\sqrt{f(u)f(v)}\right]$ or $\left[\sqrt{f(u)f(v)}\right]$ then the edge labels are distinct. In this case, f is called Geometric mean labeling of G. In this paper we prove that Double Triangle snake and Alternate Double Triangular snake graphs are Geometric mean graphs.

Keywords: Graph, Geometric mean graph, Double Triangular snake, Alternate Double Triangular snake.

1. INTRODUCTION

All graph in this paper are finite and undirected graph G = (V, E) with p vertices and q edges. For all detailed survey of graph labeling we refer to Gallian [1]. For all other standard terminology and notations we follow Harary [2]. We will provide brief summary of definitions and other informations which are required for the present investigation.

Definition 1.1: A graph G = (V,E) with p vertices and q edges is called a Geometric mean graph if it is possible to label the vertices $x \in V$ with distinct labels f(x) from 1,2.....q+1 in such a way that when each edge e=uv is labeled with $f(e=uv) = \left[\sqrt{f(u)f(v)}\right]$ or $\left[\sqrt{f(u)f(v)}\right]$ then the edge labels are distinct.

In this case f is called Geometric mean labeling of G.

Definition 1.2: A Triangular snake T_n is obtained from a path v_1, v_2, \ldots, v_n by joining v_i to v_{i+1} to a new vertex w_i for $1 \le i \le n-1$.

That is, every edge of path is replaced by a Triangle C_3 .

Definition 1.3: An Alternate Triangular snake $A(T_n)$ is obtained from a path u_1, u_2, \ldots, u_n by joining u_i and u_{i+1} (alternatively) to new vertex v_i . That is every alternate edge of a path is replaced by C_3 .

Definition 1.4: A Double triangular snake $D(T_n)$ is the graph obtained from the path u_{1,u_2,\ldots,u_n} by joining $u_i u_{i+1}$ to two new vertices v_i , w_i , $1 \le i \le n - 1$.

Definition 1.5: Alternate Double triangular snake $A(D(T_n))$ is the graph obtained from the path $u_1, u_2...u_n$ by joining u_i, u_{i+1} (Alternatively) with two new vertices v_i and $w_i, 1 \le i \le n-1$.

S.Somasundaram and S.S.Sandhya introduced Harmonic mean labeling of a Graphs [3] and studied their behaviour in [4]. S. Somasundaram, P. Vidhyarani and S.S.Sandhya introduced the concept of Geometric mean labeling of Graphs.

In this paper we prove that Double Triangular snakes and Alternate Double Triangular snakes are Geometric mean graphs.

Corresponding author: 1S. S. Sandhya*

¹Department of Mathematics, Sree Ayyappa College for Women, Chunkankadai-629003, India.

2. MAIN RESULTS

Theorem 2.1: A Double Triangular Snake $D(T_n)$ is a Geometric mean graph

Proof: Consider a path u_1, u_2, \dots, u_n Join u_i, u_{i+1} with two new vertices $v_i, w_i, 1 \le i \le n-1$.

Define a function f: $V(D(T_n)) \rightarrow \{1, 2, \dots, q+1\}$ by

$$\begin{split} f(u_1) =& 2 \\ f(u_i) = 5i\text{-}4, \ 2 \leq i \leq n \\ f(v_1) =& 1 \\ f(v_i) =& 5i\text{-}3, \ 2 \leq i \leq n\text{-}1 \\ f(w_i) =& 5i\text{-}1, \ 1 \leq i \leq n\text{-}1. \end{split}$$

The edges are labeled with

 $f(u_1u_2) = 4$

 $f(u_i u_{i+1}) = 5i-2, 2 \le i \le n-1.$

 $f(u_i v_i) = 5i-4, 1 \le i \le n-1$

 $f(u_1v_1) = 2$

 $f(u_{i+1}v_{i}) = 5i-1, 2 \le i \le n-1$

 $f(u_1w_1) = 3$

 $f(u_i w_{i}) = 5i-3, 2 \le i \le n-1.$

 $f(u_{i+1}w_i) = 5i, 1 \le i \le n-1.$

This makes D(T_n) a Geometric mean graph

Example 2.2: The Geometric mean labeling of $D(T_4)$ is given below



Figure: 1

Next we have the following

Theorem 2.3: Alternate Double Triangular snake A(D(T_n)) is a Geometric mean graph.

Proof: Let G be the graph $A(D(T_n))$. Let P_n be the path $u_1 u_2 \dots u_n$. To construct G, join u_i , u_{i+1} (alternatively) with two new vertices v_i , $w_i \ 1 \le i \le n-1$.

Here we consider two different cases

¹S. S. Sandhya^{*} and ²S. Somasundaram / Geometric Mean Labeling On Double Triangular Snakes / IJMA- 5(7), July-2014.

Case (1): If the Double Triangular Snake $A(D(T_n))$ starts from u_1 then we need to considered two subcases.

Sub case (1) (a): If n is odd, then



Figure: 2

In this case, f provides a Geometric mean labeling

Sub case (1) (b): If n is even, then define a function $f:V(G) \rightarrow \{1,2...,q+\}$ by $f(u_1) = 2$

 $f(u_i) = 3i-1, 2 \le i \le \frac{n}{2}$

 $f(v_1) = 1$

$$f(v_i) = 6(i-1), 2 \le i \le \frac{n-1}{2}$$

$$f(w_i) = 6i-2, \ 1 \le i \le \frac{n-1}{2}$$

The edges are labeled with

 $f(u_{i}u_{i+1}) = 3i+1, \forall i = 1,3...,n-1$ $f(u_{i}u_{i+1}) = 3i, \forall i = 2,4,6...,n-2$ $f(u_{2i-1} v_{i}) = 6i-5, \forall i = 1,2...,\frac{n}{2}$ $f(u_{2i} v_{i}) = 6i-4, \forall i = 1,2...,\frac{n}{2}$

 $f(u_{2i-1}w_i) = 6i-3, \forall i = 1, 2, \dots, \frac{n}{2}$

 $f(u_{2i} w_i) = 6i-1, \forall i = 1, 2....\frac{n}{2}$

The labeling pattern is shown below



In the labeling pattern f is a Geometric mean labeling of G.

Case (ii): If the Double Triangular snake $A(D(T_n))$ starts from u_2 , then we have to consider two subscases.

Sub case (ii) (a): If n is odd then define a function f: $V(G) \rightarrow \{1, 2, \dots, q+1\}$ by

$$f(u_{1}) = 1$$

$$f(u_{i}) = 3i \cdot 3, \ 2 \le i \le n$$

$$f(v_{1}) = 2$$

$$f(v_{i}) = 6i \cdot 5, \ 2 \le i \le \frac{n-2}{2}$$

$$f(w_{i}) = 6i \cdot 1, \ 1 \le i \le \frac{n-2}{2}$$

Then the edges are labeled with

$$f(u_{i}u_{i+1}) = 3i-2, \forall i = 1,3,5,....n-1$$

$$f(u_{i}u_{i+1}) = 3i-1, \forall i=2,4,6...n$$

$$f(u_{2i}v_{i}) = 6i-4, \forall i=1,2...n\frac{n-2}{2}$$

$$f(u_{2i+1}, v_{i}) = 6i-3, \forall i = 1,2...,\frac{n-2}{2}$$

$$f(u_{2i} w_{i}) = 6i-2, \forall i = 1,2...,\frac{n-2}{2}$$

$$f(u_{2i+1}w_{i}) = 6i, \forall i = 1,2...,\frac{n-2}{2}$$

The labeling pattern is



Figure: 4

In this case, f provides a Geometric mean labeling for G.

Sub case (ii) (b): If n is even then define a function f: $V(G) \rightarrow \{1, 2, \dots, q+1\}$ by $f(u_1) = 1$

$$f(u_i) = 3i - 3, \ 2 \le i \le n$$

$$f(v_1) = 2$$

$$f(v_i) = 6i - 5, \ 2 \le i \le \frac{n - 2}{2}$$

$$f(w_i) = 6i - 1, \ 1 \le i \le \frac{n - 2}{2}$$
Then the edges are labeled with
$$f(u_i u_{i+1}) = 3i - 2, \ \forall i = 1, 3, 5 \dots n - 1$$

$$f(u_{i}u_{i+1}) = 3i - 1, \ \forall i = 2, 4 \dots n - 2$$

$$f(u_{2i}v_1) = 6i - 4, \ \forall i = 1, 2, \dots n - 2$$

$$f(u_{2i+1}v_i) = 6i - 3, \ \forall i = 1, 2 \dots n - \frac{n - 2}{2}$$

$$f(u_{2i+1}w_i) = 6i - 2, \ \forall i = 1, 2 \dots n - \frac{n - 2}{2}$$

$$f(u_{2i+1}w_i) = 6i, \ \forall i = 1, 2 \dots n - \frac{n - 2}{2}$$

The labeling pattern is

f(u_i

f(u_i





From all the above cases, we conclude that Alternate Double Triangular snakes $A(D(T_n))$ is a Geometric mean graph.

REFERENCES

[1] J.A.Gallian, 2010, A dynamic survey of graph labeling, The electronic Journal of Combinatories 17 # DS6.

[2] Harary.F 1988, Graph theory, Narosa publishing House Reading, New Delhi.

[3] Somasundaram.S and Ponraj.R 2003, Mean labeling of graphs, National Academy of Science Letters Vol.26, p.210-213.

[4] Somasundaram.S and Ponraj.R and Sandhya S.S, Harmonic mean labeling of Graphs, Communicated.

[5] C. Jayasekharan, S.S.Sandhya and C.DavidRaj, Harmonic Mean Labeling on Double Triangular snakes, "International Journal of Mathematics Research Vol.5 No.2 (2013), pp.251-256.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2014. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]