

## K AND K\*- BI NEAR SUBTRACTION SEMIGROUPS

S. FIRTHOUS FATIMA\*<sup>1</sup>, S. JAYALAKSHMI<sup>2</sup>

<sup>1</sup>Department of Mathematics,  
Sadakathullah Appa College (Autonomous), Thirunelveli- 627 011, Tamil Nadu, India.

<sup>2</sup>Department of Mathematics,  
Sadakathullah Appa College (Autonomous), Thirunelveli- 627 011, Tamil Nadu, India.

(Received On: 31-08-15; Revised & Accepted On: 15-09-15)

### ABSTRACT

In this paper we introduce the notion of F- bi-near subtraction semigroup. Also we give characterizations of F- bi-near subtraction semigroup.

**Mathematical subject classification:** 06F35.

**Key words:** K- bi-near subtraction semigroup, strong  $S_1$ - bi-near subtraction semigroup, strong  $S_2$ - bi-near subtraction semigroup,  $S_1$ - bi-near subtraction semigroup,  $S_2$ - bi-near subtraction semigroup, Nil near subtraction semigroup, idempotent, Nolpotent, Zero devisors, Mate function, Boolean.

### 1. INTRODUCTION

In 2007, Dheena [1] introduced Near Subtraction Algebra, Throughout his paper by a Near Subtraction Algebra, we mean a Right Near Subtraction Algebra. For basic definition one may refer to Pillz[5]. Zekiye Seydali Fathima *et.al* [3, 4] introduced the notation of  $S_1$ -near subtraction semigroup and  $S_2$ -near subtraction semigroup. Recently Firthous *et.al* [2] introduced the notation of F- Bi near subtraction semigroup. In this paper we shall obtained equivalent conditions for regularity in terms of K- Bi near subtraction semigroup.

### 2. PRELIMINARIES

A non-empty subset X together with two binary operations “-“ and “.” is said to be **subtraction semigroup** If (i)  $(X, -)$  is a subtraction algebra (ii)  $(X, .)$  is a semi group (iii)  $x(y-z)=xy-xz$  and  $(x-y)z= xz-yz$  for every  $x, y, z \in X$ . A non-empty subset X together with two binary operations “-“ and “.” is said to be **near subtraction semigroup** if (i)  $(X, -)$  is a subtraction algebra (ii)  $(X, .)$  is a semi group and (iii)  $(x-y)z= xz-yz$  for every  $x, y, z \in X$ . A non-empty subset  $X=X_1 \cup X_2$  together with two binary operations “-“ and “.” Is said to be **bi-near subtraction semigroup** (right). If (i)  $(X_1, -, .)$  is a near-subtraction semigroup (ii)  $(X_2, -, .)$  is a subtraction semigroup. A non-empty subset X is said to be  **$S_1$ -near subtraction semi group** if for every  $a \in X$  there exists  $x \in X^*$  such that  $axa=xa$ . A non-empty subset X is said to be  **$S_2$ -near subtraction semi group** if for every  $a \in X$  there exists  $x \in X^*$  such that  $axa=ax$ . A non-empty subset X is said to be **strong  $S_1$ -near subtraction semi group** if  $aba=ba$  for all  $a, b \in X$ . A non-empty subset X is said to be **strong  $S_2$ -near subtraction semi group** if  $aba=ab$  for all  $a, b \in X$ . If there exists a map  $f: X \rightarrow Y$  such that  $a = f(a)$  for all  $a$  in X then f is called a **mate function** for X. An element  $a \in X$  is said to be **Boolean** if  $a^2 = a$ . A **sub commutative near subtraction semigroup** is an intersection of  $S_1$ -near subtraction semigroup and  $S_2$ -near subtraction semigroup. that is,  $xa=ax$ . A non-empty subset X is said to be **nil-near subtraction semigroup** if there exists a positive integer  $k > 1$  such that  $a^k = 0$  Which implies that  $xa=0$  where  $x=a^{k-1}$ . A non-empty subset X is said to be **zero-symmetric**. if  $0-x=0$ ,  $ox=0$  and  $xo=0$  for all  $x \in X$ . A non-empty subset Y of X is closed under “-“ and XY strictly contained in Y is called an **X-system**. A non-empty subset  $X=X_1 \cup X_2$  together with two binary operations “-“ and “.” is said to be **F- bi near subtraction semigroups**. If (i) for every  $a \in X_1$  there exists  $x \in X_1^*$  such that  $axa=xa$ . (ii) for every  $a \in X_2$  there exists  $x \in X_2^*$  such that  $axa=ax$ .

**Corresponding Author: S. Firthous Fatima\*<sup>1</sup>, <sup>1</sup>Department of Mathematics, Sadakathullah Appa College (Autonomous), Thirunelveli- 627 011, Tamil Nadu, India.**

### 3. K AND K\*-BI NEAR SUBTRACTION SEMIGROUP

**Definition: 3.1** A non-empty subset  $X=X_1 \cup X_2$  together with two binary operations “-“ and “.” Is said to be **K- bi near subtraction semigroup**. If (i) if  $aba=ba$  for all  $a, b \in X_1$ . (ii) for every  $a \in X_2$  there exists  $x \in X_2^*$  such that  $axa=ax$ .

**Example: 3.2** Let  $X_1=\{0, a, b, 1\}$  in which “-“ and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	1	b
b	b	0	0	b
1	1	0	1	0

.	0	a	b	1
0	0	0	0	0
a	a	a	a	a
b	a	0	1	b
1	0	a	b	1

Thus  $X_1$  is a strong  $s_1$ -near subtraction semi group

Let  $X_2=\{0, a, b, 1\}$  in which “-“ and “.” be defined by

-	0	a	b	c
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
c	c	c	c	0

.	0	a	b
0	0	0	0
a	a	0	0
b	b	0	0
1	c	0	0

Then  $X_2$  is a  $S_2$ -near subtraction semi group.

Hence,  $X=X_1 \cup X_2$  is a K-bi near Subtraction Semigroup.

**Note: 3.3** Obviously, every K-bi near subtraction is a F- bi-near subtraction semi group. But the converse need not be true

**Example: 3.4** Let  $X_1=\{0, a, b, c\}$  in which “-“ and “.” be defined by

-	0	a	b	c
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
c	c	c	c	0

.	0	a	b	c
0	0	0	0	0
a	a	a	b	c
b	0	0	0	0
c	0	a	b	c

Thus  $X_1$  is a strong  $s_1$ -near subtraction semi group but not  $s_1$ - near subtraction semigroup.

Let  $X_2=\{0, a, b, c\}$  in which “-“ and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	1	b
b	b	0	0	b
1	1	0	1	0

.	0	a	b	1
0	0	0	0	0
a	a	a	a	a
b	a	0	1	b
1	0	a	b	1

Thus  $X_2$  is an  $S_2$ -near subtraction semigroup

Hence, every K- bi-near subtraction semi group need not be a F-bi near subtraction semi group.

**Definition: 3.5** A non-empty subset  $X=X_1 \cup X_2$  together with two binary operations “-“ and “.” Is said to be **K\*- bi near subtraction semigroup**. If (i) if for every  $a \in X_1$  there exists  $x \in X_1^*$  such that  $axa=xa$ . (ii)  $aba=ab$  for all  $a, b \in X_2$ .

**Example: 3.6** Let  $X_1=\{0, a, b, 1\}$  in which “-“ and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
1	1	b	a	0

.	0	a	b	1
0	0	0	0	0
a	0	a	0	0
b	0	0	b	b
1	0	a	b	1

Then  $X_1$  is a  $s_1$ -near-subtraction semi group

Let  $X_2 = \{0, a, b, 1\}$  in which “-“ and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	1	b
b	b	0	0	b
1	1	0	1	0

.	0	a	b	1
0	0	0	0	0
a	a	0	a	0
b	0	0	b	b
1	0	a	b	1

Thus  $X_2$  is a strong  $S_2$ -near subtraction semigroup.

Hence,  $X = X_1 \cup X_2$  is a strong  $K^*$ -bi-near subtraction semi group.

**Note: 3.7** Obviously, every  $K^*$ -bi near subtraction is a F- bi-near subtraction semi group. But the converse is not true

**Example: 3.8** Let  $X_1 = \{0, a, b, 1\}$  in which “-“ and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
1	1	b	a	0

.	0	a	b	1
0	0	0	0	0
a	0	a	0	0
b	0	0	b	b
1	0	a	b	1

Then  $X_1$  is a  $s_1$ -near-subtraction semi group

Let  $X_2 = \{0, a, b, c\}$  in which “-“ and “.” be defined by

-	0	a	b	c
0	0	0	0	0
a	a	0	a	b
b	b	b	0	b
c	c	c	c	0

.	0	a	b	c
0	0	0	0	0
a	a	a	b	a
b	0	0	0	0
c	0	0	0	c

Thus  $(X_2, -, .)$  is a strong  $S_2$ -near-subtraction semi group but not a  $S_2$ -near subtraction semigroup (since  $ba \neq ab$ ). Hence,  $X = X_1 \cup X_2$  is not a  $k^*$ -bi-near subtraction semi group.

#### 4. RESULTS ON K AND $K^*$ -BI NEAR SUBTRACTION SEMIGROUP

**Proposition: 4.1** If  $X$  is a  $K$ -bi near subtraction semigroup then  $X$  is a zero-symmetric

**Proof:** Let  $X = X_1 \cup X_2$  be a  $K$ -bi near Subtraction Semigroup where  $X_1$  is a strong  $S_1$ -near subtraction semigroup and  $X_2$  is a  $S_2$ -near subtraction semigroup. Since  $X_1$  is a strong  $S_1$ -near subtraction semigroup that is,  $axa = xa$  for all  $a \in X_1$  and  $x \in X_1^*$ . Substituting  $a=0$  we have  $0x0 = x0$  for all  $x \in X_1^*$ . Thus  $X_1$  is a zero-symmetric. Since  $X_2$  is a strong  $S_2$ -near subtraction semigroup that is,  $axa = ax$  for all  $a \in X_1$  and  $x \in X_1^*$ . Substituting  $a=0$  we have  $0x0 = x0$  for all  $x \in X_1^*$ . Thus  $X_1$  is a zero-symmetric. Hence,  $X = X_1 \cup X_2$  is a zero-symmetric.

**Remark: 4.2** The Converse of above Proposition need not be true

**Example: 4.3** Let  $X_1 = \{0, a, b, c\}$  in which “-“ and “.” be defined by

-	0	a	b	c
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
c	c	c	c	0

.	0	a	b	c
0	0	0	0	0
a	0	0	0	a
b	a	0	0	b
c	0	0	0	c

Thus  $X_1$  is a zero-symmetric but not a strong  $s_1$ -near subtraction semi group (Since  $cac \neq ac$ ).

Let  $X_2 = \{0, a, b, 1\}$  in which “-” and “.” be defined by

-	0	a	b	1
0	0	0	0	0
a	a	0	1	b
b	b	0	0	b
1	1	0	1	0

.	0	a	b	1
0	0	0	0	0
a	a	0	a	0
b	0	0	b	b
1	0	a	b	1

Thus  $X_2$  is a zero-symmetric and also a  $S_2$ -near subtraction semigroup. Hence, zero symmetric need not be a K- bi-near subtraction semi group.

**Proposition: 4.4** The intersection of strong  $S_1$ -near subtraction semigroup and  $S_2$ -near subtraction semigroup is sub commutative near subtraction semigroup.

**Proof:** Let  $X_1$  is a strong  $S_1$ -near subtraction semigroup. there exists  $x \in X_1^*$  such that  $axa=xa$ . (1)

(by [3], Every Strong  $S_1$ -near subtraction semigroup is a  $S_1$ -near subtraction semigroup). Let  $X_2$  is an  $S_2$ -near subtraction semigroup, there exists  $x \in X_2^*$  such that  $axa=ax$  (2)

From (1) and (2), we get  $xa=ax$ . Thus, X is a sub commutative near subtraction semigroup.

**Proposition: 4.5** Let X be a Sub-commutative K-bi near Subtraction Semigroup Then X has no non zero-zero divisor function if and only if X is Boolean.

**Proof:** Let  $X = X_1 \cup X_2$  be a K-bi near Subtraction Semigroup where  $X_1$  is a strong  $S_1$ -near subtraction semigroup and  $X_2$  is a  $S_2$ -near subtraction semigroup. Let  $a \in X_1$ . Since  $a \cdot a = a$ . That is,  $a^3 = a^2$ , which implies  $(a^2 - a)a = 0$ . Since  $X_1$  has no non zero-zero divisor function,  $a^2 - a = 0$ ,  $a^2 = a$  Thus  $X_1$  is Boolean. Let  $a \in X_2$ . Since  $X_2$  is a strong  $S_2$ -near subtraction semigroup, there exists  $x \in X_2^*$  such that  $axa=xa$ . Which implies  $aax=ax$ . (Since  $X_2$  be a Sub-commutative). That is,  $a^2x = ax$  that implies  $(a^2 - a)x = 0$ . Since  $X_2$  has no non zero-zero divisor function,  $a^2 - a = 0$ ,  $a^2 = a$  Thus  $X_2$  is Boolean. Therefore  $X = X_1 \cup X_2$  where  $X_1$  is Boolean and  $X_2$  is Boolean. Hence, X is Boolean.

**Proposition: 4.6** The intersection of  $S_1$ -near subtraction semigroup and strong  $S_2$ -near subtraction semigroup is sub commutative near subtraction semigroup.

**Proof:** Let  $X_1$  is a  $S_1$ -near subtraction semigroup. there exists  $x \in X_1^*$  such that  $axa=xa$ . (1)

Let  $X_2$  is an strong  $S_2$ -near subtraction semigroup, there exists  $x \in X_2^*$  such that  $axa=ax$  (2)

From (1) and (2), we get  $xa=ax$  (by [4], Every Strong  $S_2$ -near subtraction semigroup is a  $S_2$ -near subtraction semigroup). Thus, X is a sub commutative near subtraction semigroup.

## 5. ACKNOWLEDGEMENT

The first author expresses her deep sense of gratitude to the UGC-SERO, Hyderabad, No: F. MRP- 5365/14 for financial assistance.

## REFERENCES

1. Dheena.P and Sathesh kumar.G, On Strongly Regular Near-Subtraction semi groups, Commun. Korean Math. Soc.22 (2007), No.3, pp.323-330.
2. Firthous Fatima.S and Jeya Lakshmi.S, F-bi near subtraction semigroups, international research journal of pure algebra-5(8), 2015, 1-4
3. Seyadali Fathima.S and Balakrishnan.R,  $S_1$ -near subtraction semigroups, Ultra Scientist Vol.24 (2012), (3) A, 578-584.
4. Seyadali Fathima.S and Balakrishnan.R,  $S_2$ -near subtraction semigroups, Research Journal of Pure Algebra-2(2012), (12) A, 382-386.
5. Pilz Gunter, Near-Rings, North Holland, Amsterdam, 1983.
6. Zekiye Ciloglu, Yilmaz Ceven, On Fuzzy Near Subtraction semigroups, SDU Journal of Science (E-Journal), 2014, 9(1):193-202.

**Source of support: UGC-SERO, Hyderabad, India, Conflict of interest: None Declared**

[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]