International Journal of Mathematical Archive-7(6), 2016, 27-33 MA Available online through www.ijma.info ISSN 2229 - 5046

A NOTE ON COMMUTATIVITY OF PERIODIC NEAR-FIELD SPACES OVER NEAR-FIELDS

Dr N. V. NAGENDRAM*

Professor of Mathematics, Kakinada Institute of Technology & Science Tirupathi (v), Divili 533 433, East Godavari District, Andhra Pradesh. India.

(Received On: 24-05-16; Revised & Accepted On: 10-06-16)

ABSTRACT

A near-field space N satisfying a polynomial identity of the form $ab = \phi(a, b)$, where $\phi(A, B)$ is different word from that AB, must have nil Commutator sub near-field space. First major theorem extends this result to the case where $\phi(A, B)$ varies with a, b with the restriction that all $\phi(A, B)$ have length at least three and are not of the form A^nB or AB^n . Further restrictions on the $\phi(A, B)$ are then shown to yields commutativity of a near-field space N. Among these a semi simple sub near-field space and a near-field space specifically that each $\phi(A, B)$ begins with B and has at least 2 in A. The final theorem establishes commutativity of near-field spaces N satisfying ab = bas where s = s(a, b) is an element of the center of the sub near-field space generated by a and b. All near-field spaces considered are either periodic by hypothesis or turn out to be periodic near-field spaces in the course of the in depth study and investigation of the near-field spaces.

Key words: Near-ring, Near-field, periodic Near-field, sub Near-field, sub Near-field space, ideal.

2000 Mathematics Subject Classification: 43A10, 46B28, 46H25, 46H99, 46L10, 46M20.

SECTION 1: PRELIMINARY RESULT ON PERIODIC NEAR-FIELD SPACES

Let $\phi = \phi$ (A, B) be a mapping or word or monomial in the non-commuting in -determinates A and B i.e., ϕ is a polynomial of form

$$B^{11} A^{k1} B^{12} A^{k2} - -B^{1s} A^{ks}$$
(1)

Where j_i , $k_i \forall i = 1, 2, ..., s$ and $\sum_{i=1}^{s} (j_i + k_i) > 0$. By the A-length (respectively B-length) of ϕ , which we denote by

 $|\phi|_{A}$ (respectively $|\phi|_{B}$), we shall mean the non-negative integer $\sum k_{i}$ (respectively $\sum j_{i}$). The sum $|\phi|_{A} + |\phi|_{B}$ will be called the length of ϕ and denoted by $|\phi|$. It will be convenient to divide the set of all words into nine types as follows:

- (a) maps with $|\phi|_A \ge 2$ and $|\phi|_B \ge 2$.
- (b) maps of form BA^n , $n \ge 1$
- (c) maps of form B^nA , $n \ge 1$
- (d) maps with $|\phi|_{\rm B} = 0$
- (e) maps with $|\phi|_{A} = 0$
- (f) maps of form $A^n BA^m$, n, m ≥ 1
- (g) maps of form B^nAB^m , n, $m \ge 1$
- (h) maps of form A^nB , $n \ge 1$
- (i) maps of form AB^n , $n \ge 1$.

Definition 1.1: A near-field space N is called periodic near-field space if for each $x \in N$, there exist distinct positive integers n, m depending on x, for which $x^n = x^m$.

Example 1.2: Among the periodic near-field spaces in fact finite near-fields which we shall refer to frequently are the cobras (p, k, ϕ) – near-fields which we define as follows.

Corresponding Author: Dr N. V. Nagendram* E-mail: nvn220463@yahoo.co.in

Dr N. V. Nagendram*/ A note on commutativity of periodic near-field spaces over Near-fields / IJMA- 7(6), June-2016.

Definition 1.3: N⁺ is the additive direct sum $GF(p^k) \oplus GF(p^k)$, ϕ is an automorphism of $GF(p^k)$, and near-field multiplication is defined by

$$(a, b)(c, d) = (ac, ad + b\phi(c))$$
 (2)

Note 1.4: Near-fields have the property that $D^2 = 0$, where D denoted the set of all zero divisor near-fields including 0 and they have as few zero divisors as non-domain may have – specifically, $|D|^2 = |N|$. They are commutative nearfields only when ϕ is the identity automorphism.

We shall make use of repeated use of two basic theorems on periodic near-field spaces. The second is a special case of an old theorem of I N Herstein. But since deduces it as a corollary of one of his more complicated commutativity theorems, we think it worthwhile to include a simple proof.

Lemma 1.5: If N is any periodic near-field space, then N has each of the following properties:

- (1) $\forall x \in N$, some power of x is idempotent.
- (2) $\forall x \in N$, there exists an integer n(x) > 1 such that $x x^{n(x)}$ is nilpotent.
- (3) $\forall x \in N$ can be expressed in the form of y + w, where $y^n = y$ for some n = n(y) > 1 and w is nilpotent.
- (4) If J is an ideal of N and x + J is anon-zero nilpotent element of N /J, then N contains a nilpotent element $u \ni x \equiv u \pmod{J}$

Proof: To prove (1): If $x^n = x^m$ with n > m, then $x^{j+k(n-m)} = x^j$ for each positive integer k and each $j \ge m$ thus, we may assume $n - m + 1 \ge m$. It follows that $x^{n-m+1} = (x^{n-m+1})^{n-m+1}$ and hence $(x^{n-m+1})^{n-m}$ is idempotent. Proved (1).

To prove (2): Let $x^n = x^m$, n > m > 1. Then we have, $x^{m-1}(x - x^{n-m+1}) = 0 = x^{m-2} x (x - x^{n-m+1}) = x^{m-2} x^{n-m+1} (x - x^{n-m+1})$.

Therefore, $x^{m-2}(x - x^{n-m+1}) = 0$ and the result follows by the obvious induction. Proved (2).

To prove (3): If $x^n = x^m$ with $n \ge n - m + 1 > m$, the proofs of (1) and (2) show that we may take $y = x^{n-m+1}$ and $w = x - x^{n-m+1}$. Proved (3).

To prove (4): If x + I is a non-zero nilpotent element of N/I, there exists a positive integer k such that $x^q \in I$ for all $q \ge k$. By the proofs of (1) and (2), N contains a nil potent element $v = x - x^q$ with $q \ge k$. Clearly, $v \equiv x \pmod{1}$. Proved (4). This completes the proof of the lemma.

Theorem 1.6: If N is a periodic near-field space with all nilpotent elements central, then N is commutative near-field.

Proof: Let N denote the set of nilpotent elements. The usual argument for commutative near-field spaces shows that N is an ideal. Moreover, for $x \in N$ and e is an idempotent in N, both ex - exe and xe - exe are in N, hence commute with e. Thus idempotents in N are central.

By (4) of lemma 1.5, we see that homomorphic images or maps inherit the hypothesis on N. Consequently, we need consider only the case of sub-directly irreducible N. Under this assumption, (1) of lemma 1.5 shows that N is either nil and hence commutative near-field or N has a unique non-zero central idempotent, necessarily a multiplicative identity element 1.

Considering this latter possibility, we see from (1) of lemma 1.5 that each element of N is either nilpotent or invertible. Thus the set D of zero divisor near-fields s is equal to N and hence is a central ideal. Moreover, by (2) of lemma 1.5 N = N/D has the aⁿ = a property of Jacobson. Hence N is a commutative near-field and its additive sub near-field is a torsion sub near-field. Thus if a, b \in N \ D, the sub near-field space of \overline{N} generated by and $\overline{b} = b + D$ is a finite near-field, which has cyclic multiplicative sub near-field. There must therefore exist $g \in N$ and $d_1, d_2 \in D$ such that $a = g' + d_1$ and $b = g' + d_2$ for some positive integers i, j. It follows that a and b must commute and our proof is complete.

SECTION 2: A NIL COMMUTATIVE SUB NEAR-FIELD SPACE AND SOME RELATIVES

Theorem 2.1: Let N be a near-field such that for each a, $b \in N$ there exist a map ϕ (A, B) of one of the types (a) to (g) and with $|\phi| \ge 3$, for which $ab = \phi$ (a, b). Then the set N of nilpotent elements forms an ideal and the Commutator ideal C(N) is contained in N.

Proof: Taking a = b shows that for each $a \in N$, $a^2 = a^k$ for some k > 2. Hence N is periodic near-field space and each nilpotent element squares to zero. We next show that idempotents of N must be central. Let e be a non-zero idempotent. © 2016, IJMA. All Rights Reserved 28 Let $a \in N$ and suppose ϕ (A, B) is a map of the allowed types for which $e(ex - exe) = \phi(e, ex - exe)$. Clearly, ϕ can not be a type (d) since $(ex - exe)^2 = 0$. Any other types has either two adjacent B's or B preceding an A. Thus e(ex - exe) = ex - exe = 0 and similarly, xe - exe = 0.

A periodic near-field space satisfies the conclusions of the theorem if nilpotent elements commute with each other, so we may complete our proof by showing that ab = 0 for all $a, b \in N$. Accordingly, let $a, b \in N$ and ϕa map such that $ab = \phi$ (a, b). If ϕ has two adjacent A's and B's then it is immediate that ab = 0. Otherwise, we have one of the following

Cases: (i) $ab = (ab)^k$ for some k > 1. (ii) ab = abab....a (iii) ab = bab....a

In case (i), $(ab)^{k-1}$ is idempotent, hence central and we get $ab = a (xb)^{k-1} b = 0$. In case (ii) right multiplication by a yields aba = 0 = ab, and in case (iii) left multiplication by b yields bab = 0 = ab. This completed the proof of the theorem.

Note 2.2: The idempotents are central apply (i) of lemma 1.5 to show that some power of each element is central and appeal to a well known theorem of I N Herstein[7].

Note 2.3: In the hypothesis of theorem 2.1, the restriction on the type of ϕ (A, B) is essential, for without it, as the near-field space of 2 x 2 matrices over GF(2) would satisfy the hypothesis.

Note 2.4: In the hypothesis of theorem 2.1 will not yield commutativity of N. The Corbas $(2, 2, \phi)$ -near-field space is a counter example where ϕ is the non identity automorphism of GF(4) indeed in this near-field space, if $u, v \in N$ and a, $b \notin \notin N$ we have $uv = vu^2$, $au = ua^2$, $ua = aua^2$ and $ab = (ba)^3 ab$. However, restriction of ϕ (A, B) to words of fixed type (a) to (g) does yield commutativity as we now prove the following theorem.

Theorem 2.5: Let β denote a fixed one of the map-types (a) to (g). Let N be a near-field space such that for each a, b \in N, there exists a type - β map ϕ (A, B), depending on a and b and having length at least three, for which $ab = \phi$ (a, b). Then N is commutative near-field.

Proof: If β is type (a), commutativity follows from a theorem of the present author. Suppose, then that β is type (d) i.e., for each a, $b \in N$, $ab = a^n$ for some $n = n(a, b) \ge 3$. Then since nilpotent elements square to 0, they left-annihilate N. Taking $a \in N$ and x an element such that $x^k = x, k > 1$ and recalling that idempotents are central, we obtain the result that $xa = xx^{k-1}x = xax^{k-1} = 0$ and by (3) of lemma 1.5 nilpotent elements right annihilate N as well and commutativity follows from theorem 1.6. it is clear that type (e) may be treated similarly.

To complete the proof, we discuss type (f) noting that (g) is similar. Let $\forall x \in N, \forall y \in N$ and $xy = x^n yx^m$ with n, $m \ge 1$. If either of n, m is greater than 1, then xy = 0. If xy = xyx, right multiplying by x yields xyx = 0 = xy. Also $yx = y^jxy^k \forall k \ge 1$, so yx = 0 as well and again commutativity follows by theorem 1.6. This completes the proof of the theorem.

Theorem 2.6: Suppose that for each x, $y \in N$ there exists an integer n(x, y) > 1 such that $xy = x^{n(x, y)}y$. Then the Commutator or ideal C(N) is nil and the nilpotent elements form an ideal. If the idempotents of N are central, then N is commutative.

Proof: Clearly, N is periodic with nilpotent elements squaring to zero and $\forall x \in N$ and v is nilpotent we have $vx = v^n x = 0$. Thus the set N of nilpotent elements is the set of annihilator of near-field space N, hence an ideal. The near-field space N/N has the $a^n = a$ property by lemma 1.5 (2), hence is commutative near-field. Thus $C(N) \subseteq N$.

Now assume that idempotent are central. If $a^k = a \forall k > 1$ and $v \in N$, we get $av = a^n v = a^{n-1}aa^{k-1}v = a^n va^{k-1} = 0$. Hence by lemma 1.5 (3) and theorem 1.6 implies N is commutative near-field.

Note 2.7: Centrality of idempotents is not implied by the condition $xy = x^n y$.

Example 2.8: The near-field space N with additive sub near-field space equal to the multiplication given by 0x = cx = 0 and $ax = bx = x \forall x \in N$. This near-field space satisfies the identity $xy = x^2y$.

Note 2.9: The idempotents are central in theorem 2.6 we can say a bit more about near-field space N specifically it is the direct sum of a zero near-field space and a J-near-field space i.e., one with Jacobson's $a^n = a$ property. For if x, y are arbitrary sub near-field spaces of N, \exists integers $n_1, n_2 > 1 \Rightarrow xy = x^{n_1}y$ and $yx = y^{n_2}x$.

Note 2.10: A standard computation yields a single n such that $xy = x^n y$ and $yx = y^n x$ and the commutativity now shows that $x^n y = xy^n$. The direct sum decomposition of near-field spaces with the latter type of constraint has essentially been obtained.

SECTION 3: MAIN RESULTS OF TWO COMMUTATIVITY THEOREMS ON PERIODIC NEAR-FIELD SPACES

Theorem 3.1: Let N be a periodic near-field space, the multiplicative semi simple near-field space of which is a semi simple near-field space. Then N is a commutative near-field.

Proof: If $a, b \in N$ and ab = 0 then ba = 0 also. Then the nilpotent elements of N form an ideal N, which since N is periodic near-field space, must coincide with the Jacobson radical J(N).

Again Dr N V Nagendram wish to deduce result from theorem 1.6. Suppose then, that μ is a non-central nilpotent element and $b \in N$ is an element not commuting with μ . Then

$$\mu b = b_{1}^{j} \mu_{1}^{k} \dots \mu_{i}^{k} \quad \forall j_{1} \ge 1, \forall \sum k_{i} \ge 2$$
(3)

If $k_1 \ge 2$ we obtain

 $\mu b = b^{j}{}_{1}\mu \mu^{k}{}_{1}{}^{-1}.....\mu^{k}{}_{s} = \mu^{t}(b^{j}{}_{1})^{q}.....\mu^{k}{}_{1}{}^{-1}....\mu^{k}{}_{s}$ (4)

If t = 1 we make no further substitutions in (4) otherwise we write $\mu b = \mu \mu^{t-1} b_1^{j_1 q} \dots \mu_s^{k-1} \dots \mu_s^{k} = \mu b_1^{j_1 qx} (\mu^{t-1})^n \dots \mu_s^{k}$. In either case we have $\mu b = \mu by$ for some $y \in J(N)$ from which it follows that $\mu b = 0 = b\mu$ is a contradiction to our choice of μ . \otimes

If $k_1 = 1$ in equation (3) then some other k_i is positive and a similar computation again yields the same contradiction \otimes . Thus nilpotent elements of N are central and this completes the proof of the theorem.

Corollary 3.2: Let N be any near-field space having as multiplicative semi simple near-field space is a semi simple near-field. Then N is a commutative near-field.

Note 3.3: Theorem 3.1 and corollary 3.2 would not be true if the condition $|\phi|_A \ge 2$ were omitted from the definition of maps the Corbas $(2, 2, \phi)$ -near-field space is the revealing example.

Theorem 3.4: Let N be any near-field space such that for each x, $y \in N$ there exists an element s = s(x, y) in the center of the sub near-field space generated by x and y for which xy = yxs. Then N is commutative near-field.

Proof: Taking x = y shows that $x^2 = x^2p(x)$, where p(x) is a polynomial with integer coefficients and zero constant term. It follows that N is periodic near-field space. Moreover, the given constraint shows that $ab = 0 \Rightarrow ba = 0 = arb \forall r \in N$. This result together with the obvious fact that nilpotent elements square to zero shows that $uvs = 0 \forall$ nilpotent element u and v and \forall s in the sub near-field space generated by u and v. Thus the nilpotent elements form an ideal N with $N^2 = 0$. Moreover, a standard arguement applied to e, ex – exe and xe – exe shows that all

The hypothesis of the theorem persist under the taking of homeomorphic images, so we need consider only sub-directly irreducible N. Since nil near-field spaces with our condition are zero near-field spaces and since sub directly irreducible near-field spaces can have at most one non-zero central idempotent, lemma 1.5 (1) allows us to assume that N has 1 and that every non nilpotent element is invertible. Hence the set D of zero divisors is an ideal which equal to N.

Since there exist distinct n, m with $(1 + 1)^n = (1 + 1)^m$, N⁺ must be a torsion sub near-field space which in view of sub direct irreducibility is a p-sub near-field space for some prime p. Since $D^2 = 0$, we have then $(p \cdot 1) (px) = p^2 x = 0$ for all $x \in N$.

Now N is clearly a duo near-field space, so we may apply earlier results of near-field spaces on sub directly irreducible duo near-field spaces. Specifically, letting S denote the intersection of the non-zero sub near-field spaces of N and noting that $N \neq D$, we have S equal to the annihilator of D i.e., S = D. By known lemma 1.5 (2) and the a" = a theorem we know that N/D is commutative near-field and hence that commutator near-fields in a near-field space belongs to D. Suppose now that $pN \neq 0$ let $px \neq 0$ and let y be an arbitrary sub near-field space of N. Since pxN is a non-zero sub near-field space , we have $xy - yx \in D = S \subseteq pxN$ and there exists $r \in N$ such that xy - yx = pxr and hence $p(xy - yx) = p^2xr = 0$. Thus pN = D is central and commutativity of near-fields of N follows from theorem 1.6.

Now suppose that we have a sub directly irreducible counter example with pN = 0. By known lemma 1.5 (3) and the fact that $D^2 = 0$. We can then choose a non-central nilpotent element u and an element $b \in N$ such that $b^{n(b)} = b$ for some n(b) > 1 and b does not commute with u. Since bu = ubs for some s in the sub near-field space generated by u and b, and since uru = 0 for all $r \in N$, we obtain bu = ubp(b), where p(A) is some polynomial with integer coefficients and zero constant term. It follows that the sub near-field space <u, b> of N generated by u and b is finite. Since the hypothesis of the theorem are inherited by sub near-field spaces and by homomorphic images we can conclude that some homomorphic image T of <u, b> is a finite sub directly irreducible counter example with pT = 0.

We can argue that T must be a near-field space for appropriate choices and finite near-field spaces N with 1 and with $D^2 = 0 = pN$ must have additive sub near-field space which is direct sum $K \oplus D$, where K is a finite near-field and D is a left vector space over K. Since one dimensional sub near-field spaces of D are left sub near-field spaces, the fact that our T is sub directly irreducible and a duo near-field space shows that D is one dimensional and $|T| = |D|^2$. We apply an earlier result to show that T is a near-field.

Consider near-field T with ψ a non-identity automorphism of $K = GF(p^k)$. let g be a generator of the multiplicative sub near-field space of K, and let φ be given by $x \to x^{pr}$, $1 \le r < k$. If $(a, b) \in T$ commutes with both (g, 0) and (0, g) then by (2) we have b = 0 and $a = \varphi(a)$. Then imposing the condition that (g, 0)(0, g) = (0, g)(g, 0)(a, 0) yields $g = \varphi(g)a$. Since $\varphi(g) = g^{pr}$ and $g = g^{pk}$ we have $g^{pk} = g^{pr}a$, so that $a = g^{pk - pr} = g^{pr (pk - r)}$. Now using fact that $\varphi(a) = a$, we get $g^{pr(pk - r - 1)(pr - 1)} = e$, where e denotes the identity element of K. Since g has order p^k -1, which is relatively prime to p', we conclude that $p^k - 1/(p^{k - r} - 1)(p^r - 1)$, which is absurd. The possibility of a counter example is thus demolished. This completes the proof of the theorem.

Note: It is tempting to conjecture that N must be commutative near-field space if it satisfies xy = yxs, where s = s(x, y) is merely assumed to belong to the sub near-field space generated by x and y and not necessarily to its center. However, the near-field space N shows that this is not true.

REFERENCES

- 1. A. Badawi, D.F. Anderson and D.E. Dobbs, Pseudo-valuation rings, Lecture notes on Pure Appl. Math, Vol. 185 (1997), pp 57-67, Marcel Dekker, New York/Basel.
- 2. A. Badawi, On domains which have prime ideals that are linearly ordered, comm. Algebra, Vol. 23(1995), pp no.4365-4373.
- 3. A. Badawi, On divided commutative rings, comm. Algebra, vol. 27(1999), pp. 1465-1474.
- 4. A. Badawi, On φ-pseudo valuation rings, Lecture notes Pure Appl. Math. Vol. 205(1999), pp. 101 110, Marcel Dekker, New York/Basel.
- 5. A. Badawi, On ϕ -pseudo valuation rings II, to appear in Houston J Math.
- 6. A. Badawi, Algebraic On ϕ -chained rings, and ϕ -pseudo valuation rings to appear in Houston J Math.
- 7. A Badawi, Remarks on ϕ -pseudo valuation rings comm... Algebra. vol. 28(2000), pp. 2343-2358.
- 8. A. Badawi, On chained overrings of pseudo valuation rings, Commun. Algebra 28(2000), pp.2359-2366.
- 9. G Pilz, near-rings, Amsterdam.
- N V Nagendram, T V Pradeep Kumar and Y V Reddy On "Semi Noetherian Regular Matrix δ–Near Rings and their extensions", International Journal of Advances in Algebra (IJAA), Jordan, ISSN 0973-6964, Vol.4, No.1, (2011), pp.51-55.
- 11. N V Nagendram, T V Pradeep Kumar and Y V Reddy "A Note on Bounded Matrices over a Noetherian Regular Delta Near Rings", (BMNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, Copyright@MindReaderPublications, ISSNN0:0973-6298, pp.13-19.
- N V Nagendram,T V Pradeep Kumar and Y V Reddy "A Note on Boolean Regular Near-Rings and Boolean Regular δ–Near Rings", (BR-delta-NR) published in International Journal of Contemporary Mathematics,IJCM Int. J. of Contemporary Mathematics, Vol. 2, No. 1, June 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp. 29 - 34.
- N V Nagendram,T V Pradeep Kumar and Y V Reddy "on p-Regular δ–Near-Rings and their extensions", (PR-delta-NR) accepted and to be published in int. J. Contemporary Mathematics (IJCM),0973-6298,vol.1, no.2, pp.81-85, June 2011.
- N V Nagendram, T V Pradeep Kumar and Y V Reddy "On Strongly Semi –Prime over Noetherian Regular δ– Near Rings and their extensions", (SSPNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, pp.83-90.
- 15. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Structure Theory and Planar of Noetherian Regular δ-Near–Rings (STPLNR-delta-NR)",International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.79-83, Dec, 2011.
- 16. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Matrix's Maps over Planar of Noetherian Regular δ-Near–Rings (MMPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.203-211, Dec, 2011.

- N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On IFP Ideals on Noetherian Regular-δ- Near Rings(IFPINR-delta-NR)", Int. J. of Contemporary Mathematics, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, Vol. 2, No. 1, pp.53-58, June 2011.
- 18. N V Nagendram, B Ramesh paper "A Note on Asymptotic value of the Maximal size of a Graph with rainbow connection number 2*(AVM-SGR-CN2*)" published in an International Journal of Advances in Algebra (IJAA) Jordan @ Research India Publications, Rohini, New Delhi, ISSN 0973-6964 Volume 5, Number 2 (2012), pp. 103-112.
- 19. N V Nagendram and B Ramesh on "Polynomials over Euclidean Domain in Noetherian Regular Delta Near Ring Some Problems related to Near Fields of Mappings(PED-NR-Delta-NR & SPR-NF)" Accepted and published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046,vol.3,no.8, pp no. 2998-3002,2012.
- 20. N V Nagendram research paper on "Near Left Almost Near-Fields (N-LA-NF)" communicated to for 2nd intenational conference by International Journal of Mathematical Sciences and Applications, IJMSA @ mindreader publications, New Delhi on 23-04-2012 also for publication.
- 21. N V Nagendram, T Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized Near Fields and (m, n) Bi-Ideals over Noetherian regular Delta-near rings (GNF-(m, n) BI-NR-delta-NR)", published in an International Journal of Theoretical Mathematics and Applications (TMA), Greece, Athens, dated 08-04-2012.
- 22. N V Nagendram, Smt.T.Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "Applications of Linear Programming on optimization of cool freezers (ALP-on-OCF)" Published in International Journal of Pure and Applied Mathematics, IJPAM-2012-17-670 ISSN-1314-0744 Vol-75 No-3(2011).
- 23. N V Nagendram "A Note on Algebra to spatial objects and Data Models(ASO-DM)" Published in international Journal American Journal of Mathematics and Mathematical Sciences, AJMMS,USA, Copyright @ Mind Reader Publications, Rohini, New Delhi, ISSN. 2250-3102, Vol.1, No.2 (Dec. 2012), pp. 233 247.
- 24. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "A Note on Pi-Regularity and Pi-S-Unitality over Noetherian Regular Delta Near Rings (PI-R-PI-S-U-NR-Delta-NR)" Published in International Electronic Journal of Pure and Applied Mathematics, IeJPAM-2012-17-669 ISSN-1314-0744 Vol-75 No-4(2011).
- 25. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "Ideal Comparability over Noetherian Regular Delta Near Rings (IC-NR-Delta-NR)" Published in International Journal of Advances in Algebra (IJAA, Jordan),ISSN 0973-6964 Vol:5,NO:1(2012), pp.43-53@ Research India publications, Rohini, New Delhi.
- 26. N. V. Nagendram, S. Venu Madava Sarma and T. V. Pradeep Kumar, "A Note On Sufficient Condition Of Hamiltonian Path To Complete Grphs (SC-HPCG)", IJMA-2(11), 2011, pp.1-6.
- 27. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Noetherian Regular Delta Near Rings and their Extensions(NR-delta-NR)", IJCMS, Bulgaria, IJCMS-5-8-2011,Vol.6,2011,No.6, 255-262.
- 28. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Semi Noehterian Regular Matrix Delta Near Rings and their Extensions(SNRM-delta-NR)", Jordan, @ResearchIndiaPublications, Advances inAlgebraISSN 0973-6964 Volume 4, Number 1 (2011), pp.51-55© Research India Publicationspp.51-55
- 29. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Boolean Noetherian Regular Delta Near Ring(BNR-delta-NR)s", International Journal of Contemporary Mathematics, IJCM Int. J. of Contemporary Mathematics , Vol. 2, No. 1-2, Jan-Dec 2011, Mind Reader Publications, ISSN No: 0973-6298, pp. 23-27.
- N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Bounded Matrix over a Noetherian Regular Delta Near Rings(BMNR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1-2, Jan-Dec 2011 , Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.11-16
- N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Strongly Semi Prime over Noetherian Regular Delta Near Rings and their Extensions(SSPNR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1, Jan-Dec 2011 ,Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.69-74.
- 32. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On IFP Ideals on Noetherian Regular Delta Near Rings(IFPINR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1-2, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.43-46.
- 33. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Structure Thoery and Planar of Noetherian Regular delta-Near–Rings (STPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM ,accepted for international conference conducted by IJSMA, New Delhi December 18,2011, pp:79-83, Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
- 34. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Matrix's Maps over Planar of Noetherian Regular delta-Near–Rings (MMPLNR-delta-NR)", International Journal of Contemporary Mathematics ,IJCM, accepted for international conference conducted by IJSMA, New Delhi December 18,2011,pp:203-211,Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
- 35. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Some Fundamental Results on P- Regular delta-Near–Rings and their extensions (PNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM,Jan-December'2011,Copyright@MindReader Publications,ISSN:0973-6298, vol.2,No.1-2,PP.81-85.

Dr N. V. Nagendram*/ A note on commutativity of periodic near-field spaces over Near-fields / IJMA- 7(6), June-2016.

- 36. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized ideal based-zero divisor graphs of Noetherian regular Delta-near rings (GIBDNR- d-NR)", International Journal of Theoretical Mathematics and Applications (TMA)accepted and published by TMA, Greece, Athens, ISSN:1792 - 9687 (print),vol.1, no.1, 2011, 59-71, 1792-9709 (online),International Scientific Press, 2011.
- 37. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Inversive Localization of Noetherian regular Delta-near rings (ILNR- Delta-NR)", International Journal of Pure And Applied Mathematics published by IJPAM-2012-17-668, ISSN.1314-0744 vol-75 No-3,SOFIA, Bulgaria.
- 38. N V Nagendram, S V M Sarma, Dr T V Pradeep Kumar "A note on Relations between Barnette and Sparse Graphs" published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals, 2(12),2011, pg no.2538-2542,ISSN 2229 – 5046.
- 39. N V Nagendram "On Semi Modules over Artinian Regular Delta Near Rings(S Modules-AR-Delta-NR) Accepted and published in an International Journal of Mathematical Archive (IJMA)", An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046, IJMA-3-474, 2012.
- 40. N VNagendram1, N Chandra Sekhara Rao2 "Optical Near field Mapping of Plasmonic Nano Prisms over Noetherian Regular Delta Near Fiedls (ONFMPN-NR-Delta-NR)" accepted for 2nd international Conference by International Journal of Mathematical Sciences and Applications, IJMSA @ mind reader publications, New Delhi going to conduct on 15 – 16 th December 2012 also for publication.
- 41. N V Nagendram, K V S K Murthy (Yoga), "A Note on Present Trends on Yoga Apart From Medicine Usage and Its Applications(PTYAFMUIA)" published by the International Association of Journal of Yoga Therapy, IAYT 18 th August, 2012.
- 42. N V Nagendram, B Ramesh, Ch Padma, T Radha Rani and S V M Sarma research article "A Note on Finite Pseudo Artinian Regular Delta Near Fields(FP AR-Delta-NF)" communicated to International Journal of Advances in Algebra, IJAA, Jordan on 22 nd August 2012.
- 43. Dr N V Nagendram, Professor, Kakinada Institute of Technology(KITS), Divili, Dr T V Pradeep Kumar "Characterisation of near-field spaces over Baer-ideals" IV International Conference at Thailand 19 – 20 th Dec 2015 by IJMSA.
- 44. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology (KITS), Divili "A note on TL-Ideal of near-field spaces over regular delta near-rings" published by IJMA, Vol.No.6, No.8, pp. 51 65, 2015.
- 45. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology (KITS), Divili "A note on B₁ -near-field spaces over regular delta near-rings" published by IJMA, Vol. 6, No.8, pp. No.144 151, 2015.
- 46. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology(KITS), Divili, Dr T V Pradeep Kumar "Amenability for dual concrete complete near-field spaces over regular delta near -rings" published by IJMSA, Vol. No. pp. No. 2014.
- 47. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology(KITS), Divili, Dr T V Pradeep Kumar, Dr D Venkateswarlu "Completeness of near-field spaces over near-fields" published by IJMA, 2014, Vol. No.5, ISBN. 2229 5046, No. 2, pp. 65 74.
- 48. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology(KITS), Divili, "A Note on Divided near-field spaces and phi- pseudo valuation near-field spaces over regular delta near rings" published by IJMA, 2015, Vol. No.6, ISBN. 2229 – 5046, No. 4, pp. 31 - 38.
- 49. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology (KITS), Divili, A Text Book "Algebraic topology over near-fields & Semi-simple near-fields and its Applications" published by Research India Publications, New Delhi ISBN: 978-93-84443-62-7, 2015.
- 50. Dr N V Nagendram, Professor of Mathematics, Kakinada Institute of Technology (KITS), Divili, Advanced research article "Sum of annihilator near-field spaces over near-ring is Annihilator near-field space (SA NFS ONR A NFS) "ISSN No. 2229 5046, Vol. No.7, No.1, 2016, pp No.125 136.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]