

## SUPER FIBONACCI GRACEFUL LABELING FOR (a, m)- SHELLGRAPH

A. SOLAIRAJU\*<sup>1</sup>, D. SENTHIL KUMAR<sup>2</sup> AND T. LENIN<sup>3</sup>

<sup>1</sup>Associate Professor of Mathematics, Jamal Mohamed College, Trichy, India.

<sup>2</sup>Lecturer in Mathematics, BWDA Arts & Science College, Tindivanam, India.

<sup>3</sup>Associate Professor of Mathematics, Khadhar Mohideen college, Athiramapattaim, India.

(Received On: 25-05-16; Revised & Accepted On: 10-06-16)

### ABSTRACT

A  $(p, q)$ -graph  $G = (V, E)$  has super Fibonacci graceful labeling if there exists an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  where  $F_k$  is the  $k^{\text{th}}$  Fibonacci number of the Fibonacci series such that its induced map  $f^+: E(G) \rightarrow \{F_1, F_2, \dots, F_q\}$  defined by  $f^+(xy) = |f(x) - f(y)|$  for all edge  $xy$  in  $G$ , is injective. In this paper, the existence of super Fibonacci graceful labeling for the various types of  $(a, m)$  – shell graph and all the above  $(a, m)$  – shell graph merged with a fan.

**Keywords:** Graceful graph, Fibonacci graceful graph, super Fibonacci graceful graph, shell graph,  $(a, m)$ –shellgraph.

### SECTION 1 –INTRODUCTION

The notions of Fibonacci graceful labeling and super Fibonacci graceful labeling were introduced by Kathiraesan and Amutha [2006] and they proved [2010] that Path  $(P_n)$ , caterpillar and bi-star  $(B_{m,n})$  are Fibonacci graceful. Also they proved every fan  $(F_n)$  graph is super Fibonacci graceful. Jeba Jesintha and Ezhilarasi Hilda [2015] obtain the graceful labeling for Shell – Butterfly graph. Meena, Renugha and Sivasakthi [2015] found cordial Labeling for different types of shell graph, shell graph merging with path, shell graph merging with star, multiple shell graph. These results on super Fibonacci graceful labeling and other results are mentioned in Gallian survey [2011].

### SECTION 2 - BASIC DEFINITIONS

**Definition 2.1: Graceful graph:** A  $(p, q)$ -graph  $G = (V, E)$  has graceful labeling if there exists an injection map  $f: V(G) \rightarrow \{0, 1, 2, \dots, q\}$  such that its induced map  $f^+: E(G) \rightarrow \{1, 2, 3, \dots, q\}$  defined by  $f^+(xy) = |f(x) - f(y)|$  for each edge  $xy$  in  $G$ , is injective. Then the graph  $G$  is graceful.

**Definition 2.2: Fibonacci graceful graph:** A  $(p, q)$ -graph  $G = (V, E)$  has Fibonacci graceful labeling if there exists an injective map  $f: V(G) \rightarrow \{0, 1, 2, \dots, F_q\}$  where  $F_k$  is the  $k^{\text{th}}$  Fibonacci number in the Fibonacci series such that its induced map  $f^+: E(G) \rightarrow \{F_1, F_2, \dots, F_q\}$  defined by  $f^+(xy) = |f(x) - f(y)|$  for every edge  $xy$  in  $G$ , is injective.

**Definition 2.3: Super Fibonacci graceful graph:** A  $(p, q)$ -graph  $G = (V, E)$  has super Fibonacci graceful labeling if there exists an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  where  $F_k$  is the  $k^{\text{th}}$  Fibonacci number in Fibonacci series such that its induced map  $f^+: E(G) \rightarrow \{F_1, F_2, \dots, F_q\}$  defined by  $f^+(xy) = |f(x) - f(y)|$  for every edge  $xy$  in  $G$ , is injective.

**Definition 2.4: Shell graph:** A shell  $S_{n,n-3}$  of width  $n$  is a graph obtained by taking  $(n-3)$  concurrent chords in a cycle  $C_n$  on  $n$  vertices. The vertex at which all the chords are concurrent is called apex. The two vertices adjacent to the apex have degree 2, apex has degree  $(n-1)$  and all the other vertices have degree 3.

**Definition 2.5:  $(a, m)$  – Shell graph  $(SH_{a,m})$ :** Let  $a \geq 2$  and  $m \geq 2$  be integer and  $n = am + (a+2)$ . An  $(a, m)$ –shell is a cycle  $C_n$  with vertices  $v_1, v_2, \dots, v_n$  together with  $m$  concurrent chords (edges) at  $v_1$  (Apex) and other ends of chords are adjacent to the vertex  $v_{ai+2}$ , where  $i = 1, 2, 3, \dots, m$ . Here the degree of  $v_1$  is  $(m+2)$ , and the degree of  $v_{ai+2}$ , (where  $i = 1, 2, \dots, m$ ) is 3. All the remaining vertices are of degree 2.

Corresponding Author: A. Solairaju\*<sup>1</sup>

<sup>1</sup>Associate Professor of Mathematics, Jamal Mohamed College, Trichy, India.

### SECTION 3: SUPER FIBONACCI GRACEFUL LABELING FOR VARIOUS TYPES OF SHELL MODEL GRAPHS

**Theorem 3.1:** A (4, m)- shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let G be the graph (a, m)-shell graph whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{4i+2}\}_{i=1}^{m+1}\}$  where  $n = 4m + 6$  and  $m \geq 2$  is any integers.

Construct an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let i vary 1 to n.

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{5k+(r-1)}; \text{ where } i = 4k + r; k = 0, 1, 2, \dots, (m+1); r = 2; \\ f(v_i) &= F_{5k+r}; \text{ where } i = 4k + r; k = 0, 1, 2, \dots, m; r = 3; \\ f(v_{4k}) &= F_{5k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{4k+1}) &= F_{(5k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

The induced map  $f^+$  from f defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges xy in G, is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps f and  $f^+$ . Hence the graph (4, m)-shell graph is super Fibonacci graceful.

**Theorem 3.2:** A(7, m) – shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let G be the graph (7,m)-shell graph whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{7i+2}\}_{i=1}^{m+1}\}$  where  $n = 7m + 6$  and  $m \geq 2$  is any integers.

Define an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let i be in  $\{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{8k+(r-1)}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5; \\ f(v_i) &= F_{8k+r}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, m; r = 3, 6; \\ f(v_i) &= F_{8k+(r+1)}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, m; r = 4; \\ f(v_{7k}) &= F_{8k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{7k+1}) &= F_{(8k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

The induced map  $f^+$  of f defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges xy in G, is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps f and  $f^+$ . Hence the graph (7, m)-shell graph is super Fibonacci graceful.

**Theorem 3.3:** A(10, m) – shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let G be the graph (10, m)-shell graph whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{10i+2}\}_{i=1}^{m+1}\}$  where  $n = 10m + 12$  and  $m \geq 2$  is any integers.

An injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is defined as follows:

Let  $i \in \{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{11k+(r-1)}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5, 8; \\ f(v_i) &= F_{11k+r}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, m; r = 3, 6, 9; \\ f(v_i) &= F_{11k+(r+1)}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, m; r = 4, 7; \\ f(v_{10k}) &= F_{11k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{10k+1}) &= F_{(11k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

Its induced map  $f^+$  defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges xy in G, is an injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps f and  $f^+$ . Hence the graph (10, m)-shell graph is super Fibonacci graceful.

**Theorem 3.4:** A (13, m) – shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let G be the graph (13, m)-shell graph whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{13i+2}\}_{i=1}^{m+1}\}$  where  $n = 13m + 15$  and  $m \geq 2$  is any integers.

An injective map  $f: V(SH_{13,m}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is defined as follows:

Let  $i$  be in  $\{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{14k+(r-1)}; \text{ where } i = 13k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5, 8, 11; \\ f(v_i) &= F_{14k+r}; \text{ where } i = 13k + r; k = 0, 1, 2, \dots, m; r = 3, 6, 9, 12; \\ f(v_i) &= F_{14k+(r+1)}; \text{ where } i = 13k + r; k = 0, 1, 2, \dots, m; r = 4, 7, 10; \\ f(v_{13k}) &= F_{14k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{13k+1}) &= F_{14k-1}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

Its induced map  $f^+$  defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $(13, m)$ -shell graph is super Fibonacci graceful.

**Theorem 3.5:** A  $(16, m)$  – shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let  $G$  be the graph  $(16, m)$ -shell graph whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{16i+2}\}_{i=1}^{m+1}\}$  where  $n = 16m + 18$  and  $m \geq 2$  is any integers.

Define an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let  $i$  vary from 1 to  $n$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{17k+(r-1)}; \text{ where } i = 16k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5, 8, 11, 14; \\ f(v_i) &= F_{17k+r}; \text{ where } i = 16k + r; k = 0, 1, 2, \dots, m; r = 3, 6, 9, 12, 15; \\ f(v_i) &= F_{17k+(r+1)}; \text{ where } i = 16k + r; k = 0, 1, 2, \dots, m; r = 4, 7, 10, 13; \\ f(v_{16k}) &= F_{17k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{16k+1}) &= F_{17k-1}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

Its induced map  $f^+$  defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$ , is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $(16, m)$ -shell graph is super Fibonacci graceful.

**Theorem 3.6:** A  $(19, m)$  – shell graph is super Fibonacci graceful, for all integers  $m \geq 2$ .

**Proof:** Let  $G$  be the  $(19, m)$ -shellgraph, whose vertex set is  $\{v_1, \dots, v_n\}$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{19i+2}\}_{i=1}^{m+1}\}$  where  $n = 19m + 21$  and  $m \geq 2$  is any integers.

Construct an injective map  $f: V(G) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let  $i$  be in  $\{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{20k+(r-1)}; \text{ where } i = 19k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5, 8, 11, 14, 17; \\ f(v_i) &= F_{20k+r}; \text{ where } i = 19k + r; k = 0, 1, 2, \dots, m; r = 3, 6, 9, 12, 15, 18; \\ f(v_i) &= F_{20k+(r+1)}; \text{ where } i = 19k + r; k = 0, 1, 2, \dots, m; r = 4, 7, 10, 13, 16; \\ f(v_{19k}) &= F_{20k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{19k+1}) &= F_{20k-1}; \text{ where } k = 1, 2, \dots, (m+1); \end{aligned}$$

Its induced map  $f^+$  defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$ , is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $(19, m)$ -shell graph is super Fibonacci graceful.

#### SECTION 4: (a, m) - SHELL GRAPH MERGING WITH A FAN GRAPH

The following definitions are given:

**Definition 4.1:** The join  $G_1 + G_2$  of two graphs  $G_1$  and  $G_2$  with disjoint vertex sets  $V_1$  and  $V_2$  and edge sets  $X_1$  and  $X_2$  is the graph union  $G_1 \cup G_2$  together with all edges joining each vertex of  $V_1$  to every vertex of  $V_2$ . A Fan graph  $F_n$  is  $K_1 + P_n$

**Definition 4.2:**  $SH_{a,m}^{F_t}$  is a graph having one copy of the  $(a, m)$  – shell graph  $(SH_{a,m})$  and one copy of fan graph  $(F_t)$  such that the apex of the graph  $SH_{a,m}$  is identified with the vertex of maximum degree in fan graph  $(F_t)$ .

**Theorem 4.3:** The graph  $G = SH_{4,m}^{F_t}$  is super Fibonacci graceful, for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph  $F_t$  and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of graph  $SH_{4,m}$ . Then the graph  $SH_{4,m}^{F_t}$  is obtained by the apex of the  $SH_{4,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{4i+2}\}_{i=1}^{m+1} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}\}$  where  $n = (4m + 6)$  for all integers  $m, t \geq 2$ .

An injective map  $f: V(SH_{4,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is defined as follows:

Let  $i \in \{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{5k+(r-1)}; \text{ where } i = 4k + r; k = 0, 1, 2, \dots, (m+1); r = 2; \\ f(v_i) &= F_{5k+r}; \text{ where } i = 4k + r; k = 0, 1, 2, \dots, m; r = 3; \\ f(v_{4k}) &= F_{5k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{4k+1}) &= F_{(5k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(u_j) &= F_{(5m+2j+5)}; \text{ where } j = 1, 2, \dots, t; \end{aligned}$$

Its induced map  $f^+$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{4,m}^{F_t}$  is super Fibonacci graceful.

**Theorem 4.4:** The graph  $G = SH_{7,m}^{F_t}$  is super Fibonacci graceful for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph ( $F_t$ ) and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of the graph  $SH_{7,m}$ . Then the graph  $SH_{7,m}^{F_t}$  is obtained by the apex of the  $S_{7,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{7i+2}\}_{i=1}^{m+1} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}\}$  where  $n = 7m + 9$  for all integers  $m, t \geq 2$ .

An injective map  $f: V(SH_{7,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is defined as follows:

Let  $i$  be an element in  $\{1, 2, \dots, n\}$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{8k+(r-1)}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5; \\ f(v_i) &= F_{8k+r}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, m; r = 3, 6; \\ f(v_i) &= F_{8k+(r+1)}; \text{ where } i = 7k + r; k = 0, 1, 2, \dots, m; r = 4; \\ f(v_{7k}) &= F_{8k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{7k+1}) &= F_{(8k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(u_j) &= F_{(8m+2j+8)}; \text{ for } j = 1, 2, 3, \dots, t. \end{aligned}$$

Its induced map  $f^+$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{7,m}^{F_t}$  is super Fibonacci graceful.

**Theorem 4.5:** The graph  $G = SH_{10,m}^{F_t}$  is super Fibonacci graceful, for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph ( $F_t$ ) and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of graph  $SH_{10,m}$ . Then the graph  $SH_{10,m}^{F_t}$  is obtained by the apex of the  $SH_{10,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{10i+2}\}_{i=1}^{m+1} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}\}$  where  $n = 10m + 12$  for all integers  $m, t \geq 2$ .

Define an injective map  $f: V(SH_{10,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let  $i$  vary from 1 to  $n$ .

$$\begin{aligned} f(v_1) &= F_0; \\ f(v_i) &= F_{11k+(r-1)}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, (m+1); r = 2, 5, 8; \\ f(v_i) &= F_{11k+r}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, m; r = 3, 6, 9; \\ f(v_i) &= F_{11k+(r+1)}; \text{ where } i = 10k + r; k = 0, 1, 2, \dots, m; r = 4, 7; \\ f(v_{10k}) &= F_{11k}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(v_{10k+1}) &= F_{(11k-1)}; \text{ where } k = 1, 2, \dots, (m+1); \\ f(u_j) &= F_{(11m+2j+11)}; \text{ for } j = 1, 2, 3, \dots, t. \end{aligned}$$

Its induced map  $f^+$  from  $f$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{10,m}^{F_t}$  is super Fibonacci graceful.

**Theorem 4.6:** The graph  $G = SH_{13,m}^{F_t}$  is super Fibonacci graceful, for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph  $(F_t)$  and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of graph  $SH_{13,m}$ . Then the graph  $SH_{13,m}^{F_t}$  is obtained by the apex of the  $SH_{13,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{13i+2}\}_{i=1}^{m+1}\} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}$  where  $n = 13m + 15$  for all integers  $m, t \geq 2$ .

An injective map  $f: V(SH_{13,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is found as follows:

Let  $i \in \{1, 2, \dots, n\}$ .

$f(v_1) = F_0$ ;

$f(v_i) = F_{14k+(r-1)}$ ; where  $i = 13k + r$ ;  $k = 0, 1, 2, \dots, (m+1)$ ;  $r = 2, 5, 8, 11$ ;

$f(v_i) = F_{14k+r}$ ; where  $i = 13k + r$ ;  $k = 0, 1, 2, \dots, m$ ;  $r = 3, 6, 9, 12$ ;

$f(v_i) = F_{14k+(r+1)}$ ; where  $i = 13k + r$ ;  $k = 0, 1, 2, \dots, m$ ;  $r = 4, 7, 10$ ;

$f(v_{13k}) = F_{14k}$ ; where  $k = 1, 2, \dots, (m+1)$ ;

$f(v_{13k+1}) = F_{14k-1}$ ; where  $k = 1, 2, \dots, (m+1)$ ;

$f(u_j) = F_{(14m+2j+14)}$ ; for  $j = 1, 2, 3, \dots, t$ .

Its induced map  $f^+$  from  $f$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{13,m}^{F_t}$  is super Fibonacci graceful.

**Theorem 4.7:** The graph  $G = SH_{16,m}^{F_t}$  is super Fibonacci graceful, for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph  $(F_t)$  and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of the graph  $S_{16,m}$ . Then the graph  $SH_{16,m}^{F_t}$  is obtained by the apex of the  $S_{16,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{16i+2}\}_{i=1}^{m+1}\} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}$  where  $n = 16m + 18$  for all integers  $m, t \geq 2$ .

Define an injective map  $f: V(SH_{16,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  as follows:

Let  $i$  vary from 1 to  $n$ .

$f(v_1) = F_0$ ;

$f(v_i) = F_{17k+(r-1)}$ ; where  $i = 16k + r$ ;  $k = 0, 1, 2, \dots, (m+1)$ ;  $r = 2, 5, 8, 11, 14$ ;

$f(v_i) = F_{17k+r}$ ; where  $i = 16k + r$ ;  $k = 0, 1, 2, \dots, m$ ;  $r = 3, 6, 9, 12, 15$ ;

$f(v_i) = F_{17k+(r+1)}$ ; where  $i = 16k + r$ ;  $k = 0, 1, 2, \dots, m$ ;  $r = 4, 7, 10, 13$ ;

$f(v_{16k}) = F_{17k}$ ; where  $k = 1, 2, \dots, (m+1)$ ;

$f(v_{16k+1}) = F_{17k-1}$ ; where  $k = 1, 2, \dots, (m+1)$ ;

$f(u_j) = F_{(17m+2j+17)}$ ; for  $j = 1, 2, 3, \dots, t$ .

Its induced map  $f^+$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{16,m}^{F_t}$  is super Fibonacci graceful.

**Theorem 4.8:** The graph  $G = SH_{19,m}^{F_t}$  is super Fibonacci graceful, for all integers  $m, t \geq 2$ .

**Proof:** Let  $\{u_0, u_1, u_2, \dots, u_t\}$  be the vertex set of fan graph  $(F_t)$  and  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of the graph  $SH_{19,m}$ . Then the graph  $SH_{19,m}^{F_t}$  is obtained by the apex of the  $SH_{19,m}$  graph  $v_1$  is merged with the maximum degree of the fan graph  $u_0$ .  $|V(G)| = n + t$  and edge set is  $\{\{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_{19i+2}\}_{i=1}^{m+1}\} \cup \{v_1 u_j\}_{j=1}^t \cup \{u_j u_{j+1}\}_{j=1}^{t-1}$  where  $n = 19m + 21$  for all integers  $m, t \geq 2$ .

An injective map  $f: V(SH_{19,m}^{F_t}) \rightarrow \{F_0, F_1, F_2, \dots, F_q\}$  is constructed as follows:

Let  $i$  be in  $\{1, 2, \dots, n\}$ .

$f(v_1) = F_0$ ;

$$\begin{aligned} f(v_i) &= F_{20k+(r-1)}; \text{ where } i = 19k+r; k = 0,1,\dots,(m+1); r = 2,5,8,11,14,17; \\ f(v_i) &= F_{20k+r}; \text{ where } i = 19k+r; k = 0,1,2,\dots,m; r = 3,6,9,12,15,18; \\ f(v_i) &= F_{20k+(r+1)}; \text{ where } i = 19k+r; k = 0,1,2,\dots,m; r = 4,7,10,13,16; \\ f(v_{19k}) &= F_{20k}; \text{ where } k = 1,2,\dots,(m+1); \\ f(v_{19k+1}) &= F_{20k-1}; \text{ where } k = 1,2,\dots,(m+1); \\ f(u_j) &= F_{(20m+2j+20)}; \text{ for } j = 1,2,3,\dots,t. \end{aligned}$$

Its induced map  $f^+$  from  $f$  is defined by  $f^+(xy) = |f(x) - f(y)|$  for each edges  $xy$  in  $G$  is injective. Thus the given graph has super Fibonacci graceful labeling from the injective maps  $f$  and  $f^+$ . Hence the graph  $SH_{19,m}^{F_t}$  is super Fibonacci graceful.

## CONCLUSIONS

The problem still remains open for various types of trees merged with the apex of the new types of shell graphs as super Fibonacci graceful.

## REFERENCES

1. Jeba Jesintha.J and Ezhilarasi Hilda.K, Shell - Butterfly graphs are graceful, International Journal of Pure & Applied Mathematics, Vol.101, No.6, (2015), 949-956.
2. Joseph A.Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics 17, (2010), #DS6, Thirteenth edition, November 17, (2011).
3. Kathiresan.K.M and Amutha.S, Fibonacci graceful graphs, Ph.D Thesis, Madurai Kamaraj University (October , 2006).
4. Kathiresan.K.M and Amutha.S, Fibonacci graceful graphs, Art. Combinatorics, 97(2010), 41-50.
5. Meena.S, Renugha.M and Sivasakthi.M, Cordial Labeling for different types of shell graph, International Journal of Scientific & Engineering Research, Volume 6, Issue 9,(September, 2015), 1282-1288.

**Source of support: Nil, Conflict of interest: None Declared**

**[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**