

SYMMETRIC BI- f -DERIVATIONS IN ALMOST DISTRIBUTIVE LATTICES

G. C. RAO*¹, K. RAVI BABU ²

¹ Department of Mathematics, Andhra University, India.

² Department of Mathematics, Govt. Degree College, Sabbavaram,
Visakhapatnam, Andhra Pradesh, India - 530003.

(Received On: 20-06-16; Revised & Accepted On: 25-07-16)

ABSTRACT

In this paper, we introduce the concept of symmetric bi- f -derivation in an Almost Distributive Lattice (ADL) and derive some important properties of symmetric bi- f -derivations in ADLs.

AMS 2000 subject Classification: 06D99, 06D20.

keywords: Almost Distributive Lattice (ADL), symmetric bi-derivations, symmetric bi- f -derivations, isotone symmetric bi- f -derivations and weak ideal.

1. INTRODUCTION

The concept of derivation in an ADL was introduced in our earlier paper [11]. The notion of derivation in Lattices was first given in G.Szasz [15] in 1974. Earlier Posner[9] introduced derivations in ring theory and later several authors worked on it ([2], [5]). Several authors worked on derivations in Lattices ([1], [3], [4], [6], [7], [8], [16], [17] and [18]). We have introduced the concept of f -derivations in an ADL in our paper [12] and the concept of symmetric bi-derivations in an ADL in our paper [13]. The concept of symmetric bi- f -derivations in lattices was introduced by Kyung Ho Kim [6] in 2012.

In 1980, the concept of an Almost Distributive Lattice(ADL) was introduced by U.M.Swamy and G.C Rao[14]. In this paper, we introduce the concept of symmetric bi- f -derivation in an ADL and derive some important properties. We introduce the concept of an isotone symmetric bi- f -derivation in an ADL and establish a set of conditions which are sufficient for the trace of a symmetric bi- f -derivation on an ADL with a maximal element to become an isotone . Also, we prove $D(x, y) = fx \wedge D(x \vee z, y)$ if D is isotone and $D(x, y) = [fx \wedge D(x \vee z, y)] \vee D(x, y)$ if f is a join homomorphism or an increasing function on L . We define a set $F_a(L)$ for each $a \in L$ and prove that it is a weak ideal if D is a join preserving symmetric bi- f -derivation on an ADL L with 0 where f is a join-homomorphism.

2. PRELIMINARIES

In this section , we recollect certain basic concepts and important results on Almost Distributive Lattices.

*Corresponding Author: G. C. Rao*¹*

¹ *Department of Mathematics, Andhra University, India.*

Definition 2.1:[10] An algebra (L, \vee, \wedge) of type $(2, 2)$ is called an Almost Distributive Lattice if it satisfies the following axioms:

$$L_1 : (a \vee b) \wedge c = (a \wedge c) \vee (b \wedge c) \quad (RD \wedge)$$

$$L_2 : a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c) \quad (LD \wedge)$$

$$L_3 : (a \vee b) \wedge b = b$$

$$L_4 : (a \vee b) \wedge a = a$$

$$L_5 : a \vee (a \wedge b) = a$$

Definition 2.2:[10] Let X be any non-empty set. Define, for any $x, y \in L$, $x \vee y = x$ and $x \wedge y = y$. Then (X, \vee, \wedge) is an ADL and such an ADL, we call discrete ADL.

Through out this paper L stands for an ADL (L, \vee, \wedge) unless otherwise specified.

Lemma 2.3:[10] For any $a, b \in L$, we have

$$(i) \quad a \wedge a = a$$

$$(ii) \quad a \vee a = a.$$

$$(iii) \quad (a \wedge b) \vee b = b$$

$$(iv) \quad a \wedge (a \vee b) = a$$

$$(v) \quad a \vee (b \wedge a) = a.$$

$$(vi) \quad a \vee b = a \text{ if and only if } a \wedge b = b$$

$$(vii) \quad a \vee b = b \text{ if and only if } a \wedge b = a.$$

Definition 2.4:[10] For any $a, b \in L$, we say that a is less than or equal to b and write $a \leq b$, if $a \wedge b = a$ or, equivalently, $a \vee b = b$.

Definition 2.5:[10] For any $a, b, c \in L$, we have the following

(i) The relation \leq is a partial ordering on L .

$$(ii) \quad a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c). \quad (LD \vee)$$

$$(iii) \quad (a \vee b) \vee a = a \vee b = a \vee (b \vee a).$$

$$(iv) \quad (a \vee b) \wedge c = (b \vee a) \wedge c.$$

(v) The operation \wedge is associative in L .

$$(vi) \quad a \wedge b \wedge c = b \wedge a \wedge c.$$

Theorem 2.6:[10] For any $a, b \in L$, the following are equivalent.

$$(i) \quad (a \wedge b) \vee a = a$$

$$(ii) \quad a \wedge (b \vee a) = a$$

$$(iii) \quad (b \wedge a) \vee b = b$$

$$(iv) \quad b \wedge (a \vee b) = b$$

$$(v) \quad a \wedge b = b \wedge a$$

$$(vi) \quad a \vee b = b \vee a$$

$$(vii) \text{ the supremum of } a \text{ and } b \text{ exists in } L \text{ and equals to } a \vee b$$

$$(viii) \text{ there exists } x \in L \text{ such that } a \leq x \text{ and } b \leq x$$

$$(ix) \text{ the infimum of } a \text{ and } b \text{ exists in } L \text{ and equals to } a \wedge b.$$

Definition 2.7:[10] L is said to be associative, if the operation \vee in L is associative.

Theorem 2.8:[10] The following are equivalent.

- (i) L is a distributive lattice.
- (ii) the poset (L, \leq) is directed above.
- (iii) $a \wedge (b \vee a) = a$, for all $a, b \in L$.
- (iv) the operation \vee is commutative in L .
- (v) the operation \wedge is commutative in L .
- (vi) the relation $\theta := \{(a, b) \in L \times L \mid a \wedge b = b\}$ is anti-symmetric.
- (vii) the relation θ defined in (vi) is a partial order on L .

Lemma 2.9:[10] For any $a, b, c, d \in L$, we have the following

- (i) $a \wedge b \leq b$ and $a \leq a \vee b$
- (ii) $a \wedge b = b \wedge a$ whenever $a \leq b$.
- (iii) $[a \vee (b \vee c)] \wedge d = [(a \vee b) \vee c] \wedge d$.
- (iv) $a \leq b$ implies $a \wedge c \leq b \wedge c$, $c \wedge a \leq c \wedge b$ and $c \vee a \leq c \vee b$.

Definition 2.10:[10] An element $0 \in L$ is called zero element of L , if $0 \wedge a = 0$ for all $a \in L$.

Lemma 2.11:[10] If L has 0 , then for any $a, b \in L$, we have the following

- (i) $a \vee 0 = a$, (ii) $0 \vee a = a$ and (iii) $a \wedge 0 = 0$.
- (iv) $a \wedge b = 0$ if and only if $b \wedge a = 0$.

Definition 2.12:[14] Let L be a non-empty set and $x_0 \in L$. Define, for $x, y \in L$,

$$\begin{aligned} x \wedge y &= y \text{ if } x \neq x_0 \\ &= x \text{ if } x = x_0 \text{ and} \end{aligned}$$

$$\begin{aligned} x \vee y &= x \text{ if } x \neq x_0 \\ &= y \text{ if } x = x_0, \text{ then } (L, \vee, \wedge, x_0) \text{ is an ADL with } x_0 \text{ as zero element. This is called discrete ADL} \\ &\text{with zero.} \end{aligned}$$

An element $x \in L$ is called maximal if, for any $y \in L$, $x \leq y$ implies $x = y$.

We immediately have the following.

Lemma 2.13:[10] For any $m \in L$, the following are equivalent:

- (1) m is maximal
- (2) $m \vee x = m$ for all $x \in L$
- (3) $m \wedge x = x$ for all $x \in L$.

Definition 2.14:[10] A nonempty subset I of L is said to be an ideal if and only if it satisfies the following:

- (1) $a, b \in I \Rightarrow a \vee b \in I$
- (2) $a \in I, x \in L \Rightarrow a \wedge x \in I$.

Definition 2.15:[10] A nonempty subset I of L is said to be an initial segment of L if, $a \in L$ and $x \in L$ such that $x \leq a$ imply that $x \in I$.

Definition 2.16:[13] A nonempty subset I of L is said to be a weak ideal if and only if it satisfies the following:

- (1) $a, b \in I \Rightarrow a \vee b \in I$
- (2) I is an initial segment of L .

Observe that every ideal of L is weak ideal, but not converse.

Definition 2.17: [10] A function $f : L \rightarrow L$ is said to be an ADL homomorphism if it satisfies the following:

- (1) $f(x \wedge y) = fx \wedge fy$,
- (2) $f(x \vee y) = fx \vee fy$ for all $x, y \in L$.

Definition 2.18: A function $d : L \rightarrow L$ is called an isotone, if $dx \leq dy$ for any $x, y \in L$ with $x \leq y$.

3. SYMMETRIC bi- f -Derivations IN ADLs

We begin this section with the following definition of a symmetric map and a symmetric bi-derivation in an ADL.

Definition 3.1: [13] A mapping $D : L \times L \rightarrow L$ is called symmetric if $D(x, y) = D(y, x)$ for all $x, y \in L$.

If $D(x, z) \leq D(y, z)$ for any $x, y \in L$ with $x \leq y$, then we call D as an isotone map on L .

Definition 3.2: [13] A symmetric function $D : L \times L \rightarrow L$ is called a symmetric bi-derivation on L , if $D(x \wedge y, z) = [y \wedge D(x, z)] \vee [x \wedge D(y, z)]$ for all $x, y, z \in L$.

Observe that a symmetric bi-derivation D on L also satisfies

$$D(x, y \wedge z) = [z \wedge D(x, y)] \vee [y \wedge D(x, z)] \text{ for all } x, y, z \in L.$$

The following definition introduces the notion of an symmetric bi- f -derivation on ADLs.

Definition 3.3: A symmetric function $D : L \times L \rightarrow L$ is called a symmetric bi- f -derivation on, if there exists a function $f : L \rightarrow L$ such that

$$D(x \wedge y, z) = [fy \wedge D(x, z)] \vee [fx \wedge D(y, z)] \text{ for all } x, y, z \in L.$$

Obviously, a symmetric bi- f -derivation D on L satisfies the relation

$$D(x, y \wedge z) = [fz \wedge D(x, y)] \vee [fy \wedge D(x, z)] \text{ for all } x, y, z \in L.$$

Example 3.4: Let $f : L \rightarrow L$ be a function such that $f(x \wedge y) = fx \wedge fy$ for all $x, y \in L$. Let $a \in L$ and define a function $D : L \times L \rightarrow L$ by $D(x, y) = fx \wedge fy \wedge a$ for all $x, y \in L$. Then D is a symmetric bi- f -derivation on L .

Example 3.5: Every symmetric bi-derivation on L is a symmetric bi- f -derivation, where $f : L \rightarrow L$ is the identity map.

But, a symmetric bi- f -derivation need not be a symmetric bi-derivation. For, consider the following example.

Example 3.6.: Let L be discrete ADL with 0 and $0 \neq a \in L$. Define a function $f : L \rightarrow L$ by $fx = a$ for all $x \in L$ and $D : L \times L \rightarrow L$ by $D(x, y) = a$ for all $x, y \in L$, then D is a symmetric bi- f -derivation on L but not a symmetric bi-derivation.

Example 3.7: Let L be a discrete ADL with at least two elements. Define a function $D : L \times L \rightarrow L$ by $D(x, y) = x \wedge y$ for all $x, y \in L$, then D is not a symmetric bi- f -derivation on L . Since, it is not a symmetric map.

Lemma 3.8: Let D be a symmetric bi- f -derivation on L . Then the following hold:

1. $D(x, y) = fx \wedge D(x, y)$ for all $x, y \in L$
2. $D(x \wedge z, y) = [fx \vee fz] \wedge D(x \wedge z, y)$ for all $x, y, z \in L$
3. If L has 0, then $f0 = 0$ implies $D(0, y) = 0$ for all $y \in L$

Proof: (1) Let $x, y \in L$.

Then $D(x, y) = D(x \wedge x, y) = [fx \wedge D(x, y)] \vee [fx \wedge D(x, y)] = fx \wedge D(x, y)$.

(2) Let $x, y, z \in L$. Then

$$\begin{aligned} [fx \vee fz] \wedge D(x \wedge z, y) &= [fx \vee fz] \wedge [[fz \wedge D(x, y)] \vee [fx \wedge D(z, y)]] \\ &= [[fx \vee fz] \wedge fz \wedge D(x, y)] \vee [[fx \vee fz] \wedge fx \wedge D(z, y)] \\ &= [fz \wedge D(x, y)] \vee [fx \wedge D(z, y)] = D(x \wedge z, y). \end{aligned}$$

(3) Suppose L has 0 and $f0 = 0$. Then,

by (1) above, $D(0, y) = f0 \wedge D(0, y) = 0 \wedge D(0, y) = 0$.

Corollary 3.9: If d is the trace of a symmetric bi- f -derivation D , then $dx = fx \wedge dx$ for all $x \in L$.

Theorem 3.10: If d is the trace of a symmetric bi- f -derivation on an associative ADL L , then $d(x \wedge y) = (fy \wedge dx) \vee D(x, y) \vee (fx \wedge dy)$.

Proof: Let $x, y \in L$. Then

$$\begin{aligned} d(x \wedge y) &= D(x \wedge y, x \wedge y) \\ &= [fy \wedge D(x, x \wedge y)] \vee [fx \wedge D(y, x \wedge y)] \\ &= [fy \wedge [[fy \wedge D(x, x)] \vee [fx \wedge D(x, y)]]] \vee [fx \wedge [[fy \wedge D(y, x)] \vee [fx \wedge D(y, y)]]] \\ &= (fy \wedge dx) \vee D(x, y) \vee (fx \wedge dy). \end{aligned}$$

Corollary 3.11: If d is the trace of a symmetric bi- f -derivation on an ADL L , then $fy \wedge dx \leq d(x \wedge y)$.

Proof: Let $x, y \in L$. Then

$$\begin{aligned} d(x \wedge y) &= D(x \wedge y, x \wedge y) \\ &= [fy \wedge D(x, x \wedge y)] \vee [fx \wedge D(y, x \wedge y)] \\ &= [fy \wedge [[fy \wedge D(x, x)] \vee [fx \wedge D(x, y)]]] \vee [fx \wedge D(y, x \wedge y)] \\ &= [(fy \wedge dx) \vee D(x, y)] \vee [fx \wedge D(y, x \wedge y)]. \end{aligned}$$

Thus $fy \wedge dx \leq (fy \wedge dx) \vee D(x, y) \leq d(x \wedge y)$.

Theorem 3.12: Let m be a maximal element of L and d be the trace of a symmetric bi- f -derivation D on L such that fm is also a maximal element. Then the following are equivalent.

1. d is an isotone map on L
2. $dx = fx \wedge dm$ for all $x \in L$
3. $d(x \wedge y) = dx \wedge dy$ for all $x, y \in L$
4. $d(x \vee y) = dx \vee dy$ for all $x, y \in L$.

Proof: (1) \Rightarrow (2): Let $x \in L$. By Corollary 3.11, $fx \wedge dm \leq d(m \wedge x) = dx$.

On the other hand, since d is an isotone, $d(x \wedge m) \leq dm$. Thus $fm \wedge dx \leq d(x \wedge m) \leq dm$.

Therefore, $dx = fx \wedge dx = fm \wedge fx \wedge dx = fx \wedge fm \wedge dx \leq fx \wedge dm$. Hence $dx = fx \wedge dm$.

(2) \Rightarrow (3): Let $x, y \in L$. Then $d(x \wedge y) = x \wedge y \wedge dm = x \wedge dm \wedge y \wedge dm = dx \wedge dy$.

Then $d(x \wedge y) = f(x \wedge y) \wedge d \quad \# \quad fx \wedge fy \wedge dm = fx \wedge dm \wedge fy \wedge dm = dx \wedge dy$.

(2) \Rightarrow (4): Let $x, y \in L$. Then $d(x \vee y) = (x \vee y) \wedge dm = (x \wedge dm) \vee (y \wedge dm) = dx \vee dy$.

Then $d(x \vee y) = f(x \vee y) \wedge dm = (fx \vee fy) \wedge dm = (fx \wedge dm) \vee (fy \wedge dm) = dx \vee dy$.

(3) \Rightarrow (1) and (4) \Rightarrow (1) are trivial.

Lemma 3.13: Let D be a symmetric bi- f -derivation on L . Then the following hold:

1. If D is isotone, then $D(x, y) = fx \wedge D(x \vee z, y)$
2. If f is a join homomorphism, then $D(x, y) = [fx \wedge D(x \vee z, y)] \vee D(x, y)$
3. If f is increasing, then $D(x, y) = [fx \wedge D(x \vee z, y)] \vee D(x, y)$

Proof: Let $x, y, z \in L$.

(1) Suppose D is an isotone function on L .

Then $D(x, y) \leq D(x \vee z, y)$. Thus $D(x, y) \wedge fx \wedge D(x \vee z, y) = D(x, y)$.

Therefore $D(x, y) \leq fx \wedge D(x \vee z, y)$.

Now, $D(x, y) = D((x \vee z) \wedge x, y) = [fx \wedge D(x \vee z, y)] \vee [f(x \vee z) \wedge D(x, y)]$.

Thus $fx \wedge D(x \vee z, y) \leq D(x, y)$. Hence $D(x, y) = fx \wedge D(x \vee z, y)$.

(2) Let f be a join-homomorphism on L . Then

$$\begin{aligned} D(x, y) &= D((x \vee z) \wedge x, y) \\ &= [fx \wedge D(x \vee z, y)] \vee [f(x \vee z) \wedge D(x, y)] \\ &= [fx \wedge D(x \vee z, y)] \vee [fx \vee fz \wedge D(x, y)] \\ &= [fx \wedge D(x \vee z, y)] \vee [(fx \wedge D(x, y)) \vee (fz \wedge D(x, y))] \\ &= [fx \wedge D(x \vee z, y)] \vee [D(x, y) \vee [fz \wedge D(x, y)]] \\ &= [fx \wedge D(x \vee z, y)] \vee D(x, y). \end{aligned}$$

(3) Let f be an increasing function on L . Then $fx \leq f(x \vee z)$.

Now,

$$\begin{aligned} D(x, y) &= D((x \vee z) \wedge x, y) \\ &= [fx \wedge D(x \vee z, y)] \vee [f(x \vee z) \wedge D(x, y)] \\ &= [fx \wedge D(x \vee z, y)] \vee [f(x \vee z) \wedge fx \wedge D(x, y)] \\ &= [fx \wedge D(x \vee z, y)] \vee [fx \wedge D(x, y)] \\ &= [fx \wedge D(x \vee z, y)] \vee D(x, y). \end{aligned}$$

Definition 3.14: Let D be a symmetric bi- f -derivation on L and $a \in L$. We define $F_a(L) = \{x \in L / D(a, x) \wedge fx = fx\}$.

Lemma 3.15: Let D be a symmetric bi- f -derivation on L where f is an increasing function and $a \in L$. Then $F_a(L)$ is an initial segment in L .

Proof: Let $x, y \in L$ with $x \leq y$ and $y \in Fix_a(L)$. Since f is an increasing function, $fx \leq fy$.

Now,

$$\begin{aligned} D(x, a) \wedge fx &= D(x \wedge y, a) \wedge fx \\ &= [[fy \wedge D(x, a)] \vee [fx \wedge D(y, a)]] \wedge fx \\ &= [[fy \wedge fx \wedge D(x, a)] \vee [fx \wedge fy \wedge D(y, a)]] \wedge fx \\ &= [[fx \wedge D(x, a)] \vee [fx \wedge D(y, a) \wedge fy]] \wedge fx \\ &= [D(x, a) \vee [fx \wedge fy]] \wedge fx \\ &= [D(x, a) \vee fx] \wedge fx \\ &= fx. \end{aligned}$$

Lemma 3.16: Let D be a join preserving symmetric bi- f -derivation on L where f is a join-homomorphism and $a \in L$. Then $x \vee y \in F_a(L)$ for all $x, y \in F_a(L)$.

Proof: Let $x, y \in F_a(L)$. Then

$$\begin{aligned} D(x \vee y, a) \wedge f(x \vee y) &= [D(x, a) \vee D(y, a)] \wedge f(x \vee y) \wedge f(x \vee y) \\ &= [[D(x, a) \vee D(y, a)] \wedge [fx \vee fy]] \wedge f(x \vee y) \\ &= [[D(x, a) \wedge [fx \vee fy]] \vee [D(y, a) \wedge [fx \vee fy]]] \wedge f(x \vee y) \\ &= [[fx \vee [D(x, a) \wedge fy]] \vee [[D(y, a) \wedge fx] \vee fy]] \wedge f(x \vee y) \\ &= [[[fx \vee D(x, a)] \wedge [fx \vee fy]] \vee [[D(y, a) \vee fy] \wedge [fx \vee fy]]] \wedge f(x \vee y) \\ &= [fx \vee fy] \wedge f(x \vee y) \\ &= f(x \vee y) \wedge f(x \vee y) \\ &= f(x \vee y). \end{aligned}$$

Hence $x \vee y \in F_a(L)$.

Finally we conclude this paper with the following theorem, which is a direct consequence of Lemma 3.15 and Lemma 3.16.

Theorem 3.17: Let L be an ADL with 0 and D be a join preserving symmetric bi- f -derivation on L where f is a join-homomorphism and $a \in L$. Then $F_a(L)$ is a weak ideal of L .

REFERENCES

1. N.O.Alshehri., *Generalized Derivations of Lattices*, *International Journal of Contemp.Math.Sciences*, 5 (2010), No. 13, 629-640.
2. H.E.Bell, L.C.Kappe., *Ring in which derivations satisfy certain algebraic conditions*, *Acta Math. Hungar.*, 53(3-4) (1989), 339-346.
3. Yilmaz Ceven., *On f -derivations of lattices*, *Bull. Korean Math. Soc.*, 45 (2008), No.4, 701-707.
4. Yilmaz Ceven., *Symmetric Bi-derivations of lattices*, *Quaestiones Mathematicae*, 32 (2009), 241-245.
5. K.Kaya., *Prime rings with a derivations*, *Bull. Mater. Sci. Eng.*, 16-17 (1987), 63-71, 1988.
6. Kyung Ho Kim., *Symmetric Bi- f -derivations in lattices*, *International Journal of Mathematical Archive*, 3(10) (2012), 3676-3683.
7. Mustafa Asci and Sahin Ceran., *Generalized (f,g)-Derivations of Lattices*, *Mathematical Sciences and Applications*, E-notes (2013), Volume No.2, 56-62.
8. Mehmet Ali Ozlurk et al., *Permuting Tri-derivations in lattices*, *Quaestiones Mathematicae*, 32 (2009), 415-425.
9. E. Posner., *Derivations in prime rings*, *Proc. Amer. Math. Soc.*, 8 (1957), 1093-1100.
10. Rao, G.C., *Almost Distributive Lattices. Doctoral Thesis*, Dept. of Mathematics, Andhra University, Visakhapatnam, (1980).
11. Rao, G.C. and Ravi Babu, *The theory of Derivations in Almost Distributive Lattices*, *Comunicated to the Bulletin of International Mathematical Virtual Institute*.
12. Rao, G.C. and Ravi Babu, *f -derivations in Almost Distributive Lattices*, *Intenational Journal of Mathematical Archive*, 7(5) (2016), 134-140.
13. Rao, G.C. and Ravi Babu, *Symmetric bi-derivations in Almost Distributive Lattices*, *Comunicated to Discussiones Mathematicae - Genaral Algebra and Applications*.
14. Swamy, U.M. and Rao, G.C., *Almost Distributive Lattices*, *J. Aust. Math. Soc. (Series A)*, 31 (1981), 77-91.
15. G. Szasz., *Derivations of lattices*, *Acta Sci. Math.(Szeged)*, 37 (1975), 149-154.
16. X.L.Xin et al., *On derivations of lattices*, *Information Science*, 178 (2008), 307-316.
17. X.L.Xin., *The fixed set of derivations in lattices*, *A Springer Open Journal*, (2012).
18. Hesret Yazarli and Mehmet Ali Ozlurk., *Permuting Tri- f -derivations in lattices*, *Commun. Korean Math. Soc.*, 26 (2011), No.1, 13-21.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]