

DENSELY CO-HOPFIAN SUB NEAR-FIELD SPACES OVER NEAR-FIELD

Dr. N. V. NAGENDRAM*

Professor of Mathematics.

Kakinada Institute of Technology & Science, Tirupathi (v),
Peddapuram (M), Divili 533 433, East Godavari District, Andhra Pradesh. INDIA.

(Received On: 24-09-16; Revised & Accepted On: 21-10-16)

ABSTRACT

The sub near-field spaces of near-field of the title are introduced as a proper generalization of quasi co-Hopfian sub near-field spaces of near-field, and are characterized in several ways. The near-field space N is right non-singular if and only if the densely co-Hopfian right N -sub near-field spaces over near-field and quasi co-Hopfian right N -near-field spaces over near-field coincide. Dense co-Hopfianity is investigated for certain N -sub near-field spaces over near-field that has indecomposable decompositions complementing direct summands. For some classes of near-field spaces N , including rings, near-rings, regular δ -near-rings, near-fields with dense right socles, we determine when the injective envelope of an N -sub near-field space over near-field is densely co-Hopfian.

Key words: Goldie torsion theory, reduced rank, dense sub-near-field space, densely co-Hopfian, quasi co-Hopfian, weakly co-Hopfian, prime sub near-field space.

2000 Mathematics Subject Classification: 43A10, 46B28, 46H25, 46H99, 46L10, 46M20, 16D10, 16D70.

SECTION-1: INTRODUCTION

Throughout near-field spaces will have unity, sub near-field spaces; near-fields will be unitary. Let K denote a right sub near-field space over a near-field space N . Co-Hopfian sub near-field spaces are generalized in the following ways. A weakly co-Hopfian (wcH-sn-fs) sub near-field space is defined by the property that all injective N -endomorphisms of the sub near-field space are essential.

In depth study makes Dr N V Nagendram to investigate these concepts are clearly co-Hopfian sub near-field spaces implies weakly co-Hopfian sub near-field spaces implies quasi co-Hopfian sub near-field spaces and none of these implications can be reserved.

In this paper Dr N V Nagendram introduce and study a notion for sub near-field spaces called densely co-Hopfian sub near-field spaces (dcH-sn-fs). K_N is densely co-Hopfian sub near-field spaces (dcH-sn-fs) if for all injective N -endomorphisms f of K , $f(K)$ is a dense sub near-field space of K in the extension of Goldie torsion theory of $\text{Mod-}R$ to sub near-field space of a near-field space N , $K/f(K)$ is Z_2 – torsion. We show that Goldie torsion theory sub near-field spaces i.e. Z_2 -torsion sub near-field spaces as well as quasi co-Hopfian sub near-field spaces (dcH-sn-fs) are densely co-Hopfian sub near-field spaces (dcH-sn-fs) but not conversely.

The dcH-sn-fs property is investigated for direct sums and it can be determine that the dcH-sn-fs property for a certain sub near-field space that has an indecomposable decomposition complementing direct summands. We then discuss when the dcH-sn-fs property transfers between a sub near-field space and its injective envelope. We also consider sub near-field spaces with the property that all their sub near-field spaces are dcH-sn-fs. Such sub near-field spaces will be called completely co-Hopfian sub near-field spaces (ccH-sn-fs), and they are characterized. Finally in for three types of near-field spaces N , including near-field spaces with dense right socles, we prove that K_N is ccH-sn-fs if and only if the injective hull $E(K_N)$ of K_N is dcH-sn-fs.

Corresponding Author: Dr. N. V. Nagendram*,
Professor of Mathematics. Kakinada Institute of Technology & Science, Tirupathi (v),
Peddapuram (M), Divili 533 433, East Godavari District, Andhra Pradesh. INDIA.
E-mail: nvn220463@yahoo.co.in

We now fix our notation and state a few well known preliminary results that will be needed. As a hereditary torsion class, the class of Z_2 -torsion sub near-field spaces is closed under sub near-field spaces, factor sub near-field spaces, direct sums and extensions. These facts imply that $S \leq_d T$ and $T \leq_d U$ if and only if $S \leq_d U$. Moreover, the Goldie torsion theory is stable, that is, the Z_2 -torsion class is closed under taking injective envelopes. Let us write $Z(K/S) = S^*/S$ and $Z(K/S^*) = S^{**}/S$, and put on record several properties each of which is equivalent to S being dense in K . We omit the proofs.

Definition 1.1: Co-Hopfian sub near-field space. Let K a right sub near-field space over a near-field space N is called co-Hopfian sub near-field space if every injective N -endomorphism of K is surjective.

Definition 1.2: Quasi Co-Hopfian sub near-field space. The right (or left) sub near-field space K is called quasi co-Hopfian sub near-field space (qcH-sn-fs) if $K/f(K)$ is singular sub near-field space whenever f is an injective N -endomorphism of K .

Definition 1.3: Densely Co-Hopfian sub near-field space. The right (or left) sub near-field spaces called densely co-Hopfian sub near-field spaces (dcH-sn-fs).

Definition 1.4: Dense sub near-field space. Let K_N is densely co-Hopfian sub near-field spaces (dcH-sn-fs) if for all injective N -endomorphisms f of K , $f(K)$ is a dense sub near-field space of K in the extension of Goldie torsion theory of $\text{Mod-}R$ to sub near-field space of a near-field space N , $K/f(K)$ is Z_2 – torsion.

Definition 1.5: Essential sub near-field space. Let K be a right N -sub near-field space, S be a N -sub near-field space. Then $S \leq_e K$ will mean that S is an essential sub near-field space of K .

Definition 1.6: The singular sub near-field space of K is denoted by (K) , and $Z_2(K)$ is defined by $Z(K/Z(K)) = Z_2(K)/Z(K)$. The uniform dimension of $K/Z_2(K)$ is called the reduced rank of K , and we denote this by $r(K)$. K is called singular sub near-field space if $K = Z(K)$ and non-singular sub near-field space if $Z(K) = 0$.

Definition 1.7: Goldie torsion. The sub near-field space K is called Goldie torsion or Z_2 – torsion if $Z_2(K) = K$.

Definition 1.8: Dense near-field space. If K/S is Goldie torsion sub near-field space, then S is said to be a dense sub near-field space of K , and this fact is denoted by $S \leq_e K$.

Note 1.9: For any sub near-field space K_N , $Z_2(K) = \{x \in K : \text{ann}(x) \leq_d N_N\}$.

Definition 1.10: CS or Σ -extending. A sub near-field space of near-field space K is called CS or extending if every closed sub near-field space of K is a direct summand of K .

Note 1.11: A near-field space N for which every free sub near-field space over a near-field is CS is called Σ -extending.

Note 1.12: Over such rings, near-rings, δ -near-rings, near-fields, sub near-field spaces and near-field spaces a dcH-sn-fs is exactly a direct sum of a wch sub near-field space and a Z_2 – torsion sub near-field space.

Definition 1.13: A sub near-field space of a near-field space over a near-field K is quasi co-Hopfian sub near-field space (quasi co-Hopfian sub near-field space) if $f(K)^* = K$ for every injective endomorphism f of K .

Definition 1.14: K is densely co-Hopfian sub near-field space (dcH-sn-fs) if $f(K)^{**} = K$ for every injective endomorphism f of K .

Definition 1.15: Compressible (resp. Retractable). A sub near-field space K is called compressible respectively retractable if there exists a monomorphism (respectively non-zero homomorphism) $f: K \rightarrow N$ for any non-zero sub near-field space N of K .

Definition 1.16: Continuous. An extending sub near-field space K is called continuous if every sub near-field space V of K which is isomorphic to a direct summand of K , is a direct summand of K , every non-singular injective sub near-field space of a near-field space N over a near-field of finite reduced rank is a direct sum of indecomposable sub near-field spaces.

Definition 1.17: The endomorphism sub near-field space of an indecomposable continuous sub near-field space is a local sub near-field space, a non-singular continuous sub near-field space over a near-field space of finite reduced rank has a decomposition into indecomposable continuous sub near-field spaces such that the endomorphism near-field space over a near-field of each direct summand is local sub near-field space.

Note 1.18: A continuous sub near-field space has the finite exchange property such a decomposition complements direct summands.

Definition 1.19: Σ -quasi injective. A sub near-field space K is called Σ -quasi injective sub near-field space if every direct sum of copies of K is quasi –injective.

Note 1.20: Every Σ -CS sub near-field space, hence every Σ -quasi injective sub near-field space has an indecomposable decomposition which complements direct summands.

Proposition 1.21: Let $S \leq_d K$. The following are equivalent statements.

- (a) $S \leq_d K$
- (b) K/S^* is singular
- (c) $S^* \leq_e K$
- (d) $S^{**} = K$
- (e) $S + Z(K) \leq_e K$
- (f) $S + Z_2(K) \leq_e K$
- (g) $(S + Z_2(K))/Z_2(K) \leq_e K/Z_2(K)$
- (h) $S \oplus T \leq_e K$ for some Z_2 -torsion sub near-field space T of K
- (i) $S \cap T \neq 0$ for every non-singular sub near-field space T of K
For every sub near-field space T of K , $S \cap T \leq Z_2(K)$
- (j) For all $k \in K \setminus Z_2(K)$, there exists $n \in \mathbb{N}$ such that $kn \in S \setminus Z_2(K)$.

A notable property of dense sub near-field spaces is that their inverse under homomorphisms are again dense sub near-field spaces. We shall also make use of the following well known facts, proofs of which are given for reader's convenience.

Proposition 1.22: (a) The intersection of all dense sub near-field spaces of K is the sum $P(K)$ of all non-singular simple sub near-field spaces of K . Consequently $Z_2(K) \cap P(K) = 0$. (b) The product of two dense right sub near-field spaces is a dense sub near-field space.

Proof: To prove (a): Let $D(K)$ the intersection of all dense sub near-field spaces of K . If D is a dense sub near-field space and P is a non-singular simple sub near-field space of K , then $P \cap D \neq 0$. Hence $P \leq D$ and so $P(K) \leq D(K)$. On the other hand, if W is complement to $Z_2(K)$ then $W \leq_d K$. Thus $D(K)$ is non-singular.

However, $D(K) \leq \text{Soc}(K)$ since every essential sub near-field space is dense. Thus $D(K) \leq P(K)$.

Now if $k \in Z_2(K)$ then $\text{ann}(k) \leq_d N_N$. However, $P(N_N)$ is the intersection of all dense right sub near-field spaces of a near-field space N over a near-field, hence $P(N_N) \leq \text{ann}(k)$. Thus $Z_2(K) \cap P(K) = 0$. Proved (a).

To prove (b): Let U and V be dense right (or left) sub near-field spaces of a near-field space N over a near-field. Then N/U and U/UV are Z_2 -torsion. Therefore from the isomorphism $[N/UV]/[U/UV] \cong N/U$ we conclude that N/UV is Z_2 -torsion, hence UV is dense sub near-field space. Proved (b). This completes the proof of the proposition.

SECTION-2: DENSELY CO-HOPFIAN SUB NEAR-FIELD SPACES

In section 2, we obtain the result which describes several equivalent conditions to dcH-sn-fs property. It reduces to some parts of the base near-field space over a near-field is right non-singular sub near-field space.

Definition 2.1: K_N is called densely co-Hopfian sub near-field space (dcH-sn-fs) if the image of any injective N -endomorphism of K is a dense sub near-field space (dsn-fs)

Theorem 2.2: The following statements are equivalent for an N-sub near-field space K.

- K is dcH-sn-fs (dense co-Hopfian sub near-field space)
- K contains a dense sub near-field space V which is dcH-sn-fs(dense co-Hopfian sub near-field space) as on N-sub near-field space and $f(V) \leq_d V$ for any injective endomorphism f of K.
- There exists a dense sub near-field space V of K such that $f(V) \cap V \leq_d V$ whenever f is an injective endomorphism of K.
- If there exists an N-monomorphism $K \oplus N \rightarrow K$ then N is Z_2 -torsion.
- For every dense sub near-field space V of K and every injective endomorphism f of K, $f(V) \leq_d V$.
- For every non- Z_2 -torsion sub near-field space V of K and every injective endomorphism f of K, $f^{-1}(V)$ is non- Z_2 -torsion.
- There exists a sub near-field space V of K such that V and K/V are dcH-sn-fs (dense co-Hopfian sub near-field space) and $f^{-1}(V) = V$ for any injective endomorphism f of K.

Proof: It is obvious that (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (a).

To prove (a) \Rightarrow (d): Let $f: K \oplus V \rightarrow K$ be a monomorphism and $\tau: K \rightarrow K \oplus V$ be the canonical injection. Then $f \circ \tau$ is an injective endomorphism of K. Thus $f(K \oplus 0) \leq_d K$, hence $f(0 \oplus V)$ is Z_2 -torsion and so V is Z_2 -torsion. Proved (a) \Rightarrow (d).

To prove (d) \Rightarrow (a): Let f be an injective endomorphism of K. There exists a sub near-field space $V \leq K$ such that $f(K) \oplus V \leq_e K$. Therefore V must be Z_2 -torsion and consequently $f(K) \leq_d K$. Proved (d) \Rightarrow (a).

To prove (a) \Rightarrow (e): Let f be an injective endomorphism of K. Then we have $f(K) \leq_d f(K)$. On the other-hand, $V \leq_d K$ implies that $f(V) \leq_d f(K)$. Thus $f(V) \leq_d K$. Proved (a) \Rightarrow (e).

To prove (e) \Rightarrow (a): On applying (e) for $V = K$. Proved (e) \Rightarrow (a).

To prove (a) \Rightarrow (f): Let f be an injective endomorphism of K and V be a non- Z_2 -torsion sub near-field space of a near-field space N over a near-field. By proposition 1.21 we have that for every sub near-field space T of K, $S \cap T \leq Z_2(K)$, $f(K) \cap V$ is non- Z_2 -torsion. Thus there exists $k \in K \setminus Z_2(K)$ such that $f(k) \in K$. Consequently $k \in f^{-1}(V)$ and $k \notin Z_2(K)$. Thus $f^{-1}(V)$ is non- Z_2 -torsion. Proved (a) \Rightarrow (f).

To prove (f) \Rightarrow (a): Let g be an injective endomorphism of K such that $g(K) \not\leq_d K$. There exists a non- Z_2 -torsion sub near-field space of a near-field space N over a near-field such that $g(K) \oplus V \leq_e K$. Thus by (f) $g^{-1}(V)$ is non- Z_2 -torsion. However, $g^{-1}(V) = g^{-1}(g(K) \cap V) = g^{-1}(0) = 0$ is \otimes a contradiction. Proved (f) \Rightarrow (a).

To prove (a) \Rightarrow (g): Set $V = K$, and finally we define a mapping $\bar{f}: K/V \rightarrow K/V$ by $\bar{f}(k + V) = f(k) + V$. By hypothesis $f(K) + V \leq_d K$. On the other hand, $[f(K) + V]/f(K) \cong V/f(K) \cong V/[f(K) \cap V] = V/f(V)$. By assumption that V is dcH-sn-fs (dense co-Hopfian sub near-field space), the sub near-field space $V/f(V)$ is Z_2 -torsion and so $f(K) \leq_d f(K) + V$. It follows that $f(K) \leq_d K$. Proved (a) \Rightarrow (g). This completes the proof of the theorem.

Corollary 2.3: Let K be an N-sub near-field space.

- If K is dcH-sn-fs(dense co-Hopfian sub near-field space) then so every direct summand of K
- If $K/Z_2(K)$ is dcH-sn-fs(dense co-Hopfian sub near-field space) then so is K. In particular, every sub near-field space of finite reduced rank is dcH- sn-fs(dense co-Hopfian sub near-field space).
- Assume that U is a dense right(or left) sub near-field space of N. If KU is dcH-sn-fs(dense co-Hopfian sub near-field space) then so is K.

Proof: Obvious.

Proposition 2.4: Let K be a sub near-field space of a near-field space N over a near-field.

- If K is dcH-sn-fs(dense co-Hopfian sub near-field space) then so is every direct summand of K.
- If $K = \sum_{i \in U} K_i$ such that $f(K_i) \cap K_i \leq_d K_i$ for any injective endomorphism f of K, then K is dcH-sn-fs (dense co-Hopfian sub near-field space).

Proof: To prove (a): Let $K^{(\Lambda)}$ be dcH-sn-fs (dense co-Hopfian sub near-field space). We can assume that Λ is countable. Then $K/f(K)$ is Z_2 -torsion where $f: K^{(\Lambda)} \rightarrow K^{(\Lambda)}$ is the shift map. However K is isomorphic to $K/f(K)$ hence K is Z_2 -torsion. The converse is clear since every direct sum of Z_2 -torsion sub near-field spaces of a near-field space over a near-field is Z_2 -torsion. Proved (a).

To prove (b): Let f be an injective endomorphism of K and $V_i = f(K_i) \cap K_i$ for each $i \in I$. Define $\phi : \bigoplus_{i \in I} (K_i/V_i) \rightarrow K/f(K)$ by $(k_i + V_i)_{i \in I} \mapsto \left(\sum_{j \in J} k_j \right) + f(K)$, where J is the largest sub near-field space of I such that $k_j \notin V_j$ for any $j \in J$. Then ϕ is an epi-morphism and $\bigoplus_{i \in I} (K_i/V_i)$ is Z_2 -torsion. Thus $K/f(K)$ is Z_2 -torsion and so K is dcH-sn-fs (dense co-Hopfian sub near-field space). Proved (b). This completes the proof of the proposition.

Corollary 2.5: Let K be semi simple sub near-field space. Then K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if every non-singular homogeneous component of K is finitely generated.

Proposition 2.6: The following statements are equivalent for a sub near-field space K of a near-field space N over a near-field.

- K is wcH-sn-fs
- K is qcH-sn-fs (quasi co-Hopfian sub near-field space) and for any injective endomorphism f of K , $f(Z(K)) \leq_e Z(K)$.
- K is dcH-sn-fs (dense co-Hopfian sub near-field space) and for any injective endomorphism f of K , $f(Z(K)) \leq_e Z(K)$.
- K is qcH-sn-fs (quasi co-Hopfian sub near-field space) and for any injective endomorphism f of K , $f(Z_2(K)) \leq_e Z_2(K)$.
- K is dcH-sn-fs (dense co-Hopfian sub near-field space) and for any injective endomorphism f of K , $f(Z_2(K)) \leq_e Z_2(K)$.

Proof: Is obvious.

Corollary 2.7: Let K be a sub near-field space of a near-field space N over a near-field such that $Z(K)$ or $Z_2(K)$ is wcH-sn-fs. Then K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if K is qcH-sn-fs (quasi co-Hopfian sub near-field space) if and only if K is wcH-sn-fs (weakly co-Hopfian sub near-field space).

Proposition 2.8: Let N be a near-field space over a near-field.

- The class of dcH-sn-fs N -sub near-field spaces coincides with the class of wcH-sn-fs (weakly co-Hopfian sub near-field space) N -sub near-field spaces if and only if N is semi simple.
- The class of dcH-sn-fs N -sub near-field spaces coincides with the class of qcH-sn-fs (quasi co-Hopfian sub near-field space) N -sub near-field spaces if and only if N is right non-singular.

Proof: (a) is obvious.

To prove (b) (\Rightarrow): Let K be a N -sub near-field space of a near-field space N over a near-field. Then $Z_2(K)^{(N)}$ is Z_2 -torsion and so it is dcH-sn-fs (dense co-Hopfian sub near-field space). Thus by hypothesis $Z_2(K)^{(N)}$ is qcH-sn-fs (quasi co-Hopfian sub near-field space), hence $Z_2(K)$ is singular sub near-field space. Consequently $Z_2(K) = Z(K)$, in particular, $Z_2(E) = Z(E)$ where $E = E(N_N)$. However E is extending and so $Z(E)$ is a direct summand of E . This implies that $Z(E) = 0$, hence $Z(N_N) = 0$.

(\Leftarrow) Since the notions of Z_2 -torsion and singular sub near-field space are the same for a sub near-field space of a near-field space over a near-field is a right non-singular near-field space, the properties dcH-sn-fs (dense co-Hopfian sub near-field space) and qcH-sn-fs (quasi co-Hopfian sub near-field space) are equivalent. Proved (b). This completes the proof of the proposition.

Example 2.9: We now construct examples of dcH-sn-fs sub near-field spaces which are neither Z_2 -torsion nor qcH. Let N be a right Noetherian near-field spaces over a near-field such that $Z(N_N) \neq 0$ and $Z_2(N_N) \neq N$. then $Z_2(E) \neq Z(E)$ where $E = E(N_N)$ since $Z(N_N) \neq 0$. And also $N/Z_2(N_N) \oplus Z_2(E)^{(N)}$ is dcH-sn-fs (dense co-Hopfian sub near-field space) which is neither Z_2 -torsion nor qcH-sn-fs (quasi co-Hopfian sub near-field space).

Theorem 2.10: The following statements are equivalent for a near-field space N over a near-field.

- Every projective, free N -sub near-field space K is dcH-sn-fs (dense co-Hopfian sub near-field space)
- Every projective, free N -sub near-field space K , $K/Z(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space)
- Every projective, free N -sub near-field space K , $K/Z_2(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space)
- Every projective, free N -sub near-field space K , $K/Z_2(K)$ is qcH-sn-fs (q co-Hopfian sub near-field space)
- Every projective, free N -sub near-field space K , $K/Z(K)$ is singular
- Every projective, free N -sub near-field space K is Z_2 -torsion.
- There exists a nilpotent dense right (or left) sub near-field space in N .

- (h) $\text{Rad}(N) \leq_d N_N$ and $\text{Soc}(N_N) \leq_d N_N$.
- (i) $Z(N_N) \leq_e N_N$.
- (j) N_N is Z_2 -torsion.

Proof: This theorem we prove by cyclic method of proof.

To prove (a) \Rightarrow (f): Let K be an N -sub near-field space and Λ be an infinite set. By hypothesis K^Λ is dcH-sn-fs (dense co-Hopfian sub near-field space), hence K is Z_2 -torsion. Proved (a) \Rightarrow (f).

To prove (f) \Rightarrow (e): is obvious since $Z(K/Z(K)) = Z_2(K)/Z(K)$. Proved (f) \Rightarrow (e).

It is obvious that (e) \Rightarrow (d) \Rightarrow (b).

Now let (b) hold. Then $L = (N/Z(N_N))^{(N)} \cong N^{(N)}/Z(N_N)^{(N)}$ is dcH-sn-fs (dense co-Hopfian sub near-field space). Since $L \oplus (N/Z(N_N)) \cong L$ it implies that $N/Z(N_N)$ is Z_2 -torsion, thus $Z(N_N) \leq_d N_N$ and therefore it is evident that $Z(N_N) \leq_e N_N$. this shows that (b) \Rightarrow (i). Obvious that (i) \Rightarrow (j).

If (j) holds then every N -sub near-field space K is Z_2 -torsion as $KZ_2(N_N) \leq Z_2(K)$ hence K is dcH-sn-fs (dense co-Hopfian sub near-field space). Thus (j) \Rightarrow (a) proved.

Clearly (f) \Rightarrow (c) and (c) \Rightarrow (j) by setting $Z_2(K)$ instead of $Z(K)$ in the proof of (b) \Rightarrow (i).

If (f) holds the zero sub near-field space is nilpotent and dense right sub near-field space of a near-field space N , say $V^n = 0$. If $n = 1$ then $V = 0$ is Z_2 -torsion. If $n > 1$ then V^{n-1} is a dense right sub near-field space and so $V V^{n-1} = 0$ implies that V is Z_2 -torsion. Therefore N contains a Z_2 -torsion dense right sub near-field space, hence N_N is Z_2 -torsion. Thus every right sub near-field space of N is dense sub near-field space; hence N_N is Z_2 -torsion. Thus every right sub near-field space of N is dense and so (g) \Rightarrow (h).

Now let (h) holds and K be an N -sub near-field space. Since $\text{rad}(N_N) \leq_d N_N$ and $K \text{Rad}(N) \leq \text{Rad}(K)$ we conclude that $\text{Rad}(K) \leq_d K$. Thus $0 = \text{Rad}(P) \leq_d P$ for every simple N -sub near-field space P and so every simple N -sub near-field space is Z_2 -torsion which is singular and hence $\text{Soc}(N_N) \leq Z_2(N_N)$. By hypothesis we have $\text{Soc}(N_N) \leq_d N_N$ we conclude that $Z_2(N_N) \leq_d N_N$ and so $Z_2(N_N) = N_N$. Hence (h) \Rightarrow (j) proved. This completes the proof of the theorem.

Proposition 2.11: For a near-field space N , if $\text{Soc}(N_N) \leq_e N_N$ then each of the statements of theorem 2.11 is equivalent to any one of the following conditions.

- (a) $\text{Rad}(N) \leq_e N_N$
- (b) For every N -sub near-field space K , $K/\text{Rad}(N)$ is singular
- (c) For every N -sub near-field space K , $K/\text{Rad}(K)$ is qcH-sn-fs
- (d) Every simple N -sub near-field space is singular.
- (e) $\text{Soc}(N_N)^2 = 0$
- (f) $S(N_N)^2 = 0$.

Proof: To prove (a): Every simple N -sub near-field space is singular and so every maximal right sub near-field space is essential. Thus $\text{Soc}(N_N) \leq \text{Rad}(N)$ and then the hypothesis $\text{Soc}(N_N) \leq_e N_N$ implies that $\text{Rad}(N) \leq_e N_N$. This shows that proved (a).

It is clear that and follows (a) \Rightarrow (b) \Rightarrow (c).

Clearly, (b) \Rightarrow (d). The equality $Z(N_N) \text{Soc}(N_N) = 0$. Since $\text{Soc}(N_N) \leq_e N_N$ and $S(N_N) \leq_d \text{Soc}(N_N)$, we conclude that $S(N_N) \leq_d N_N$ and so $S(N_N)$ is a nilpotent dense right sub near-field space of near-field space N over a near-field.

Thus (f) \Rightarrow There exists a nilpotent dense right (or left) sub near-field space in N of (f) theorem 2.10. This completes the proof of the proposition.

Proposition 2.12: Let K be a homomorphic image of a CS sub near-field space. K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if K is isomorphic to a direct sum of a non-singular dcH-sn-fs (dense co-Hopfian sub near-field space) and a Z_2 -torsion sub near-field space.

Proof: Assume that $K \equiv W/V$ where U is a CS sub near-field space and $V \leq W$. First we show that $Z_2(W/V)$ is a direct summand of W/V . Assume that $Z_2(W/V) = V'/V$. Then V' is a closed (or open) sub near-field space of a near-field space over a near-field of W . In fact if $V' \leq_d N \leq W$ then N/V is Z_2 -torsion since $(N/V)/(V'/V)$ and V'/V are Z_2 -torsion. Thus $N/V \leq Z_2(W/V)$ and so $N = V'$. Therefore V' is a direct summand of W , say $W = V' \oplus V''$. Hence $W/V = V'/V \oplus (V'' + V)/V$ as desired. This completes the proof of the proposition.

Theorem 2.13: Let K be a near-field space over a near-field N such that $f(K)$ is a direct summand of K for every injective endomorphism f of K . Suppose K has an indecomposable decomposition that complements direct summands. Then K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if every non- Z_2 -torsion homogeneous component of such a decomposition of K is a finite direct sum.

Proof: (\Rightarrow) A non- Z_2 -torsion homogeneous component of an indecomposable decomposition of K is isomorphic to $N^{(\Lambda)}$ for some non- Z_2 -torsion indecomposable sub near-field space of a near-field space over a near-field V of K . Since K is dcH-sn-fs (dense co-Hopfian sub near-field space) so is $N^{(\Lambda)}$ thus Λ is finite.

(\Leftarrow) Let f be an injective endomorphism of K . By hypothesis $K = f(K) \oplus V$ for some sub near-field space V . Assume that V is non- Z_2 -torsion. By hypothesis there exists an indecomposable decomposition $K = \bigoplus_{\alpha \in S} K_\alpha$ that components direct summands and so there exists such a decomposition for V . Then $V = \bigoplus_{\beta \in T} K_\beta$. Then $K = \bigoplus_{\alpha \in S} K_\alpha$ (*)

$$\text{And } K = \bigoplus_{\alpha \in S} f(K_\alpha) \oplus \left(\bigoplus_{\beta \in T} K_\beta \right) \quad (**)$$

(*) and (**) are equivalent. Now for a $\beta_1 \in T$ such that V_{β_1} is non- Z_2 -torsion, $V_{\beta_1} \equiv K_{\alpha_1}$ for some $\alpha_1 \in S$. By hypothesis the homogeneous component of K corresponding to K_{α_1} has finitely many direct summands say $K_{\alpha_1}, K_{\alpha_2}, \dots, K_{\alpha_n}$. Then in (**) there is a homogeneous component with at least $n+1$ direct summands, i.e. $f(K_{\alpha_1}), f(K_{\alpha_2}), f(K_{\alpha_3}), \dots, f(K_{\alpha_n}), K_{\beta_1}$ which are all isomorphic to K_{α_1} . This contradicts the equivalence of (*) and (**). Therefore V is Z_2 -torsion and so $f(K) \leq_d K$. This completes the proof of the theorem.

Corollary 2.14: N be a near-field space over a near-field.

- If N is of finite reduced rank and K is a continuous N -sub near-field space, then K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if every non-singular homogeneous component of a decomposition of K into indecomposable continuous sub near-field spaces is a finite direct sum.
- If K is a Σ - quasi - injective sub near-field space, then K is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if every non-singular homogeneous component of a decomposition of K into indecomposable Σ - quasi - injective sub near-field spaces is a finite direct sum.

Note 2.15: A divisible abelian sub near-field space of a near-field space K over a near-field N is dcH-sn-fs (dense co-Hopfian sub near-field space) if and only if $n(K) < \infty$.

Note 2.16: Let N be a right Artinian local sub near-field space that is not a division sub near-field space. Then

$$\text{Rad}(N) \leq_e N_N \text{ and } \text{Soc}(N_N) \leq_e N_N \text{ and so } N \text{ satisfies all the conditions of theorem 2.10. Moreover, } S = \begin{pmatrix} N & \text{Rad}(N) \\ 0 & N \end{pmatrix}$$

$$\text{is a right Artinian sub near-field space and clearly } \text{Rad}(S) = \begin{pmatrix} \text{Rad}(N) & \text{Rad}(N) \\ 0 & \text{Rad}(N) \end{pmatrix} \leq_e S_S$$

Thus S also satisfies the conditions of theorem 2.10 although S is not local sub near-field space. Also a right Artinian sub near-field space each of the statements of theorem 2.10 holds good if and only if $\text{Rad}(N) \leq_e N_N$.

SECTION-3: COMPLETE CO-HOPFICITY AND INJECTIVE ENVELOPE NEAR-FIELD SPACES OVER NEAR-FIELD

In this chapter, we have a natural question is whether the dense co-Hopfity passes to injective envelope. As the notions of dcH-sn-fs (dense co-Hopfian sub near-field space) and qcH-sn-fs (quasi co-Hopfian sub near-field space) are the same for a non-singular sub near-field space shows that in general the answer to this question is negative. By the additive property of reduced rank if K is of finite reduced rank, then so is $E(K)$ and hence it follows that $E(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space). However, every sub near-field space of a finite reduced rank sub near-field space is dcH-sn-fs (dense co-Hopfian sub near-field space). Let us call a sub near-field space all of whose sub near-field spaces are dcH-sn-fs (dense co-Hopfian sub near-field space), a (ccH-sn-fs) completely co-Hopfian sub near-field space. Now it is natural to ask whether $E(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space) if K is ccH-sn-fs

(completely co-Hopfian sub near-field space). In the following we show that a quasi injective dcH-sn-fs (dense co-Hopfian sub near-field space) sub near-field space is ccH-sn-fs (completely co-Hopfian sub near-field space). Moreover for some classes of sub near-field spaces we show that the answer to the latter question is affirmative.

Proposition 3.1: The following statements are equivalent for a sub near-field space K of a near-field space N over a near-field.

- (a) K is ccH-sn-fs (class of co-Hopfian sub near-field space)
- (b) every dense sub near-field space of K is dcH-sn-fs (dense co-Hopfian sub near-field space)
- (c) every non-dense sub near-field space of K is dcH-sn-fs (dense co-Hopfian sub near-field space)
- (d) $X^{(N)}$ can not be embedded in K , for any non Z_2 -torsion sub near-field space X .

Proof: clearly (a) \Rightarrow (b) and (a) \Rightarrow (c)

To prove (b) \Rightarrow (a): this follows by every sub near-field space is a direct summand of an essential sub near-field space of a near-field space over a near-field. Proved (b) \Rightarrow (a).

To prove (c) \Rightarrow (a): Let V be a sub near-field space of K which is not dcH-sn-fs (dense co-Hopfian sub near-field space). There exists an injective endomorphism f of V such that $f(K) \not\leq_d K$ (not less than or equal to). Thus $f(K) \not\leq_d K$, however $f(K) \equiv K$ implies that $f(K)$ is not dcH-sn-fs (dense co-Hopfian sub near-field space) which \otimes contradicts (c). Proved (c) \Rightarrow (a). Proof of (a) \Rightarrow (d) is obvious.

To prove (d) \Rightarrow (a): Let V be a sub near-field space of K . If V is not dcH-sn-fs (dense co-Hopfian sub near-field space) then there exists a non Z_2 -torsion sub near-field space X such that $V \oplus X$ can be embedded in V which \otimes contradicts (d). Proved (d) \Rightarrow (a). This completes the proof of the proposition.

Proposition 3.2: The following statements are equivalent for a sub near-field space K of a near-field space N over a near-field.

- (a) K is dcH-sn-fs (dense co-Hopfian sub near-field space)
- (b) K is ccH-sn-fs (complete co-Hopfian sub near-field space)
- (c) $E(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space)

Proof: To prove (a) \Rightarrow (b): It is sufficient to show that every essential sub near-field space of a near-field space over a near-field K is dcH-sn-fs (dense co-Hopfian sub near-field space). Let V be an essential sub near-field space of K and g be an injective endomorphism of V . As K is quasi co-Hopfian sub near-field space injective, there exists an endomorphism f of K such that $f|_V = g$. The essentiality of V implies that f is an injective endomorphism of K , hence $f(K) \leq_d K$. Clearly, $f(V) \leq_d f(K)$, thus $f(V) \leq_d K$ and so $f(V) \leq_d V$. Proved (a) \Rightarrow (b).

To prove (b) \Rightarrow (c): obvious from theorem 2.2 of (b) for $V = K$. Proved (b) \Rightarrow (c)

To prove (c) \Rightarrow (a): by applying (a) \Rightarrow (b) to the sub near-field space $E(K)$ we conclude that $E(K)$ is ccH-sn-fs (complete co-Hopfian sub near-field space), hence K is dcH-sn-fs (dense co-Hopfian sub near-field space).

Proved (c) \Rightarrow (a). This completes the proof of the proposition.

Lemma 3.3: Let N be a near-field space for which $\text{Soc}(N_N) \leq_d N_N$ or let N be a near-field space of finite reduced rank which is either semi prime near-field space or has a. c. c. on two sided sub near-field spaces. Then every non-zero non-singular right N -sub near-field space K contains an essential sub near-field space $L = \bigoplus_{i \in I} U_i$ where each U_i is a uniform compressible right N -sub near-field space.

Theorem 3.4: Let N be a near-field space for which $\text{Soc}(N_N) \leq_d N_N$ or let N be a near-field space of finite reduced rank which is either semi prime near-field space or has a. c. c. on two sided sub near-field spaces. Then K is a ccH-sn-fs (class of co-Hopfian sub near-field space) N -sub near-field space if and only if $E(K)$ is dcH-sn-fs (dense co-Hopfian sub near-field space).

Corollary 3.5: Let N be a near-field space.

- (a) If $\text{Soc}(N_N) \leq_d N_N$, then K is ccH-sn-fs if and only if $\text{Soc}(K)$ is dcH-sn-fs
- (b) If N is right Artinian near-field space then on N -sub near-field space K is ccH-sn-fs (class of co-Hopfian sub near-field space) if and only if $r(K) < \infty$

Corollary 3.6: Let N be a near-field space for which $\text{Soc}(N_N) \leq_d N_N$ or let N be a near-field space of finite reduced rank which is either semi prime near-field space or has a. c. c. on two sided sub near-field spaces. The following statements are equivalent.

- (a) In sub near-field space- N , $\{\text{ccH-sn-fs sub near-field spaces}\} = \{\text{sub near-field spaces of finite reduced rank}\}$
- (b) Up to isomorphisms, there are only finitely many non-singular indecomposable injective N -sub near-field spaces of a near-field space over a near-field.

ACKNOWLEDGEMENT

The author being a Professor is indebted to the referee for his various valuable comments leading to the improvement of the advanced research article. This work was supported by the chairman Sri B Srinivasa Rao, Kakinada Institute of Technology & Science (K.I.T.S.), Affiliated to JNTU Kakinada, R&D education Department Science & Humanities (Mathematics), Divili 533 433. Andhra Pradesh INDIA.

REFERENCES

1. A Badawi, On 2-absorbing ideals of commutative rings, Bull. Austral. Math. Soc.75 (2007), 417-429.
2. A Badawi, U Tekir and E yetkin, On 2-absorbing primary ideals in commutative rings, Bull. Korean. Math. Soc. 51(2014), 1163-1173.
3. A Haghany and M R Vedadi Modules whose injective endomorphism are essential J Algebra 243 2001, 765-779.
4. Clark, C Lomp, N Vanaja and R Wisbauer, Lifting modules (frontiers in mathematics), Basel, 2006.
5. F W Anderson and K R Fuller Rings and Categories of modules.
6. J L Gome Pardo and P A Guil Asensio Every Σ - CS module has an indecomposable decomposition Proc. Amer. Math. Soc. 129, 2001, 947 – 954.
7. K R Goodearl and R B Warfield Jr Introduction to non-commutative Noetherian rings.
8. M L Teply Torsion free injective modules Pacific Math 28, 1969, 441-453.
9. M L Teply, Some aspects of Goldie; torsion theory Pacific J Math 29, 1969, 447-459.
10. N V Dung, D V Huynh P F Smith and W Wisbauer Extending modules, Pitman R notes 1994.
11. N V Nagendram, T V Pradeep Kumar and Y V Reddy On “Semi Noetherian Regular Matrix δ -Near Rings and their extensions”, International Journal of Advances in Algebra (IJAA), Jordan, ISSN 0973 - 6964, Vol.4, No.1, (2011), pp.51-55.
12. N V Nagendram, T V Pradeep Kumar and Y V Reddy “A Note on Bounded Matrices over a Noetherian Regular Delta Near Rings”, (BMNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, Copyright@MindReader Publications, ISSNNo:0973-6298, pp.13-19.
13. N V Nagendram, T V Pradeep Kumar and Y V Reddy “A Note on Boolean Regular Near-Rings and Boolean Regular δ -Near Rings”, (BR-delta-NR) published in International Journal of Contemporary Mathematics, IJCM Int. J. of Contemporary Mathematics, Vol. 2, No. 1, June 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp. 29 - 34.
14. N V Nagendram, T V Pradeep Kumar and Y V Reddy “on p -Regular δ -Near-Rings and their extensions”, (PR-delta-NR) accepted and to be published in int. J. Contemporary Mathematics (IJCM), 0973-6298, vol.1, no.2, pp.81-85, June 2011.
15. N V Nagendram, T V Pradeep Kumar and Y V Reddy “On Strongly Semi-Prime over Noetherian Regular δ -Near Rings and their extensions”, (SSPNR-delta-NR) published in International Journal of Contemporary Mathematics, Vol.2, No.1, June 2011, pp.83-90.
16. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy “On Structure Theory and Planar of Noetherian Regular δ -Near-Rings (STPLNR-delta-NR)”, International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.79-83, Dec, 2011.
17. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy “On Matrix's Maps over Planar of Noetherian Regular δ -Near-Rings (MMPLNR-delta-NR)”, International Journal of Contemporary Mathematics, IJCM, published by IJSMA, pp.203-211, Dec, 2011.
18. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy “On IFP Ideals on Noetherian Regular- δ - Near Rings (IFPINR-delta-NR)”, Int. J. of Contemporary Mathematics, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, Vol. 2, No. 1, pp.53-58, June 2011.
19. N V Nagendram, B Ramesh paper "A Note on Asymptotic value of the Maximal size of a Graph with rainbow connection number $2*(AVM-GR-CN2*)$ " published in an International Journal of Advances in Algebra(IJAA) Jordan @ Research India Publications, Rohini , New Delhi , ISSN 0973-6964 Volume 5, Number 2 (2012), pp. 103-112.
20. N V Nagendram research paper on "Near Left Almost Near-Fields (N-LA-NF)" communicated to for 2nd international conference by International Journal of Mathematical Sciences and Applications, IJMSA @mindreader publications, New Delhi on 23-04-2012 also for publication.

21. N V Nagendram, T Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized Near Fields and (m, n) Bi-Ideals over Noetherian regular Delta-near rings (GNF-(m, n) BI-NR-delta-NR)", published in an International Journal of Theoretical Mathematics and Applications (TMA), Greece, Athens, dated 08-04-2012.
22. N V Nagendram, Smt.T.Radha Rani, Dr T V Pradeep Kumar and Dr Y V Reddy "Applications of Linear Programming on optimization of cool freezers(ALP-on-OCF)" Published in International Journal of Pure and Applied Mathematics, IJPAM-2012-17-670 ISSN-1314-0744, Vol-75 No-3(2011).
23. N V Nagendram "A Note on Algebra to spatial objects and Data Models (ASO-DM)" Published in international Journal American Journal of Mathematics and Mathematical Sciences, AJMMS, USA, Copyright @ Mind Reader Publications, Rohini, New Delhi, ISSN. 2250-3102, Vol.1, No.2 (Dec. 2012), pp. 233 – 247.
24. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "A Note on Pi-Regularity and Pi-S-Unitality over Noetherian Regular Delta Near Rings (PI-R-PI-S-U-NR-Delta-NR)" Published in International Electronic Journal of Pure and Applied Mathematics, IeJPAM-2012-17-669 ISSN-1314-0744 Vol-75 No-4 (2011).
25. N V Nagendram, Ch Padma, Dr T V Pradeep Kumar and Dr Y V Reddy "Ideal Comparability over Noetherian Regular Delta Near Rings(IC-NR-Delta-NR)" Published in International Journal of Advances in Algebra (IJAA, Jordan), ISSN 0973-6964 Vol:5,NO:1(2012), pp.43-53@ Research India publications, Rohini, New Delhi.
26. N. V. Nagendram, S. Venu Madava Sarma and T. V. Pradeep Kumar, "A Note on Sufficient Condition of Hamiltonian Path to Complete Graphs (SC-HPCG)", IJMA-2(11), 2011, pp.1-6.
27. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Noetherian Regular Delta Near Rings and their Extensions(NR-delta-NR)", IJCMS, Bulgaria, IJCMS-5-8-2011, Vol.6, 2011, No.6, 255-262.
28. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Semi Noetherian Regular Matrix Delta Near Rings and their Extensions(SNRM-delta-NR)", Jordan, @Research India Publications, Advances in Algebra ISSN 0973-6964 Volume 4, Number 1 (2011), pp.51-55© Research India Publicationspp.51-55
29. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Boolean Noetherian Regular Delta Near Ring(BNR-delta-NR)s",International Journal of Contemporary Mathematics, IJCM Int. J. of Contemporary Mathematics, Vol. 2, No. 1-2, Jan-Dec 2011, Mind Reader Publications, ISSN No: 0973-6298, pp. 23-27.
30. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Bounded Matrix over a Noetherian Regular Delta Near Rings (BMNR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1-2, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.11-16
31. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Strongly Semi Prime over Noetherian Regular Delta Near Rings and their Extensions (SSPNR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1, Jan-Dec 2011,Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.69-74.
32. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On IFP Ideals on Noetherian Regular Delta Near Rings(IFPINR-delta-NR)", Int. J. of Contemporary Mathematics, Vol. 2, No. 1-2, Jan-Dec 2011, Copyright @ Mind Reader Publications, ISSN No: 0973-6298, pp.43-46.
33. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Structure Thoery and Planar of Noetherian Regular delta-Near-Rings (STPLNR-delta-NR)",International Journal of Contemporary Mathematics, IJCM, accepted for Ist international conference conducted by IJSMA, New Delhi December 18,2011, pp:79-83, Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
34. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "On Matrix's Maps over Planar of Noetherian Regular delta-Near-Rings (MMPLNR-delta-NR)", International Journal of Contemporary Mathematics, IJCM, accepted for Ist international conference conducted by IJSMA, New Delhi December 18,2011, pp:203-211, Copyright @ Mind Reader Publications and to be published in the month of Jan 2011.
35. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Some Fundamental Results on P- Regular delta-Near-Rings and their extensions (PNR-delta-NR)",International Journal of Contemporary Mathematics, IJCM,Jan-December'2011, Copyright @ MindReader Publications, ISSN:0973-6298, vol.2, No.1-2, PP.81-85.
36. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "A Generalized ideal based-zero divisor graphs of Noetherian regular Delta-near rings (GIBDNR- d-NR)", International Journal of Theoretical Mathematics and Applications (TMA)accepted and published by TMA, Greece, Athens, ISSN:1792- 9687 (print), vol.1, no.1, 2011, 59-71, 1792-9709 (online),International Scientific Press, 2011.
37. N V Nagendram, Dr T V Pradeep Kumar and Dr Y V Reddy "Inversive Localization of Noetherian regular Delta-near rings (ILNR- Delta-NR)", International Journal of Pure And Applied Mathematics published by IJPAM-2012-17-668, ISSN.1314-0744 vol-75 No-3, SOFIA, Bulgaria.
38. N V Nagendram¹, N Chandra Sekhara Rao² "Optical Near field Mapping of Plasmonic Nano Prisms over Noetherian Regular Delta Near Fields (ONFMPN-NR-Delta-NR)" accepted for 2nd international Conference by International Journal of Mathematical Sciences and Applications, IJMSA @ mind reader publications, New Delhi going to conduct on 15 – 16 th December 2012 also for publication.
39. N V Nagendram, K V S K Murthy (Yoga), "A Note on Present Trends on Yoga Apart From Medicine Usage and Its Applications (PTYAFMUJA)" Pubished by the International Association of Journal of Yoga Therapy, IAYT 18 th August, 2012.
40. N V Nagendram, B Ramesh, Ch Padma, T Radha Rani and S V M Sarma research article "A Note on Finite Pseudo Artinian Regular Delta Near Fields (FP AR-Delta-NF)" communicated to International Journal of Advances in Algebra, IJAA, Jordan on 22nd August 2012.

41. N V Nagendram "Amenability for dual concrete complete near-field spaces over a regular delta near-rings (ADC-NFS-R- δ -NR)" accepted for 3rd international Conference by International Journal of Mathematical Sciences and Applications, IJMSA @ mind reader publications, New Delhi going to conduct on 15 – 16th December 2014 also for publication.
42. N V Nagendram "Characterization of near-field spaces over Baer-ideals" accepted for 4th international Conference by International Journal Conference of Mathematical Sciences and Applications, IJCMSA @ mind reader publications, New Delhi going to conduct on 19 – 20 th December 2015 at Asian Institute of Technology AIT, Klaung Lange 12120, Bangkok, Thailand.
43. N V Nagendram, S V M Sarma Dr T V Pradeep Kumar "A note on sufficient condition of Hamiltonian path to Complete Graphs" published in International Journal of Mathematical archive IJMA, ISSN 2229-5046, Vol.2, No.2, Pg. 2113 – 2118, 2011.
44. N V Nagendram, S V M Sarma, Dr T V Pradeep Kumar "A note on Relations between Barnette and Sparse Graphs" published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals, 2(12), 2011, pg no.2538-2542, ISSN 2229 – 5046.
45. N V Nagendram "On Semi Modules over Artinian Regular Delta Near Rings(S Modules-AR-Delta-NR) Accepted and published in an International Journal of Mathematical Archive (IJMA)", An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046, IJMA-3-474, 2012.
46. N V Nagendram "A note on Generating Near-field efficiently Theorem from Algebraic K - Theory" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.3, No.10, Pg. 1 – 8, 2012.
47. N V Nagendram and B Ramesh on "Polynomials over Euclidean Domain in Noetherian Regular Delta Near Ring Some Problems related to Near Fields of Mappings (PED-NR-Delta-NR & SPR-NF)" Accepted and published in an International Journal of Mathematical Archive (IJMA), An International Peer Review Journal for Mathematical, Science & Computing Professionals ISSN 2229-5046, vol.3, no.8, pp no. 2998-3002,2012.
48. N V Nagendram "Semi Simple near-fields Generating efficiently Theorem from Algebraic K - Theory" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.3, No.12, Pg. 1 – 7, 2012.
49. N V Nagendram "-----" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.3, No.10, Pg. 3612 – 3619, 2012.
50. N V Nagendram, E Sudeeshna Susila, "Applications of linear infinite dimensional system in a Hilbert space and its characterizations in engg. Maths (AL FD S HS & EM)", IJMA, ISSN. 2229-5046, Vol.4, No.7, Pg. 1 – 11(19 – 29), 2013.
51. N V Nagendram, Dr T V Pradeep Kumar, "Compactness in fuzzy near-field spaces (CN-F-NS)", IJMA, ISSN. 2229 – 5046, Vol.4, No.10, Pg. 1 – 12, 2013.
52. N V Nagendram, Dr T V Pradeep Kumar and Dr Y Venkateswara Reddy, " Fuzzy Bi- Γ ideals in Γ semi near – field spaces (F Bi-Gamma I G)" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.4, No.11, Pg. 1 – 11, 2013.
53. N V Nagendram," EIFP Near-fields extension of near-rings and regular delta near-rings (EIFP-NF-E-NR) "published by International Journal of Mathematical Archive, IJMA, ISSN. 2229 - 5046, Vol.4, No.8, Pg. 1–11, 2013.
54. N V Nagendram, E Sudeeshna Susila, "Generalization of $(\in, \in Vqk)$ fuzzy sub near-fields and ideals of near-fields (GF-NF-IO-NF)", IJMA, ISSN.2229-5046, Vol.4, No.7, Pg. 1 – 11, 2013.
55. N V Nagendram, Dr T V Pradeep Kumar," A note on Levitzki radical of near-fields(LR-NF)" , Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.4, No.4, Pg.288 – 295, 2013.
56. N V Nagendram, "Amalgamated duplications of some special near-fields(AD-SP-N-F)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.4, No.2, Pg.1 – 7, 2013.
57. N V Nagendram, "Infinite sub near-fields of infinite near-fields and near-left almost near-fields (IS-NF-INF-NL-A-NF)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.4, No.2, Pg. 90 – 99, 2013.
58. N V Nagendram "-----"published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.4, No.7, Pg. 1 – 11, 2013
59. N V Nagendram, E Sudeeshna Susila, Dr T V Pradeep Kumar "Some problems and applications of ordinary differential equations to Hilbert Spaces in Engg mathematics (SP-O-DE-HS-EM)", IJMA, ISSN. 2229-5046, Vol.4, No.4,Pg. 118 – 125, 2013.
60. N V Nagendram, Dr T V Pradeep Kumar and D Venkateswarlu, "Completeness of near-field spaces over near-fields (VNFS-O-NF)" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.5, No.2, Pg. 65 – 74, 2014
61. Dr N V Nagendram "A note on Divided near-field spaces and ϕ -pseudo – valuation near-field spaces over regular δ -near-rings (DNF- ϕ -PVNFS-O- δ -NR)" published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.6, No.4, Pg. 31 – 38, 2015.
62. Dr. N V Nagendram "A Note on B_1 -Near-fields over R-delta-NR (B_1 -NFS-R- δ -NR)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.6, No.8, Pg. 144 – 151, 2015.

63. Dr. N V Nagendram "A Note on TL-ideal of Near-fields over R-delta-NR(TL-I-NFS-R- δ -NR)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.6, No.8, Pg. 51 – 65, 2015.
64. Dr. N V Nagendram "A Note on structure of periodic Near-fields and near-field spaces (ANS-P-NF-NFS)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.7, No.4, Pg. 1 – 7, 2016.
65. Dr. N V Nagendram "Certain Near-field spaces are Near-fields(C-NFS-NF)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.7, No.4, Pg. 1 – 7, 2016.
66. Dr. N V Nagendram "Sum of Annihilators Near-field spaces over Near-rings is Annihilator Near-field space (SA-NFS-O-A-NFS)", Published by International Journal of Mathematical Archive, IJMA, ISSN. 2229-5046, Vol.7, No.1, Pg. 125 – 136, 2016.
67. Dr. N V Nagendram "A note on commutativity of periodic near-field spaces", Published by IJMA, ISSN. 2229 - 5046, Vol.7, No. 6, Pg. 27 – 33, 2016.
68. Dr. N V Nagendram "Closed (or Open) sub near-field spaces of commutative near-field Space over Near-field ", Published by IJMA, ISSN. 2229 - 5046, Vol. xx, No. xx, Pg. xx – xx, 2016.
69. Sh Asgari On weakly co-Hopfian modules Bull Iran Math Soc 33, 2007, 65-72.
70. Sh Asgari, A Haghany and M R Vedadi Quasi co-hopfian modules and applications Comm. Algebra 36, 2008, 1801-1816.

Source of support: Nil, Conflict of interest: None Declared

[Copy right © 2016. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]