

TOTALLY NANO Sg CONTINUOUS FUNCTIONS
AND SLIGHTLY NANO Sg CONTINUOUS FUNCTIONS IN NANO TOPOLOGICAL SPACES

S. JEYASHRI¹, S. CHANDRASEKAR² AND G. RAMKUMAR³

¹Assistant Professor, Department of Mathematics,
Mother Teresa Women's University, Kodaikanal, Tamil Nadu, India.

²Assistant Professor, Department of Mathematics,
Arignar Anna Government Arts College, Namakkal (DT), Tamil Nadu, India

³Assistant Professor, Department of Mathematics
Arul Anandar College, Karumathur, Madurai (DT), Tamil Nadu, India.

E-mail: jeyabala1@yahoo.com¹, chandrumat@gmail.com² and ramanujam_1729@yahoo.com³

ABSTRACT

Semi generalized closed sets plays important role generalized topological spaces .In this paper we studied about Totally Nano sg continuous functions and slightly Nano sg continuous functions in Nano topological spaces

Keywords: Nano Topological spaces, nano sg-closed set, nano sg-open set, Totally Nano sg continuous functions and slightly Nano-sg continuous functions.

1. INTRODUCTION

The concept of nano topology was introduced by Lellis Thivagar [7] which was defined in terms of approximations and boundary region of a subset of an universe using an equivalence relation on it. He also established the weak forms of nano open sets namely nano open sets, nano semi open sets and nano pre open sets in a nano topological space. nano sg closed introduced by K.Bhuvaneswari [1] *et al.* .Since the advent of these notions several research papers with interesting results in different respects came to existence. In this paper we studied about Totally Nano sg continuity and Slightly Nano sg continuous functions using Nano sg closed sets in nano topological spaces

2. PRELIMINARIES

Definition 2.1: Let U be a non-empty finite set of objects called the universe and R be an equivalence relation on U named as the indiscernibility relation. Elements belonging to the same equivalence class are said to be indiscernible with one another. The pair (U, R) is said to be the approximation space. Let $X \subseteq U$.

- (i) The lower approximation of X with respect to R is the set of all objects, which can be for certain classified as X with respect to R and it is denoted by $L_R(X)$. That is,
$$L_R(X) = \bigcup_{x \in U} \{R(x) : R(x) \subseteq X\}$$
 where $R(x)$ denotes the equivalence class determined by X .
- (ii) The upper approximation of X with respect to R is the set of all objects, which can be possibly classified as X with respect to R and it is denoted by $U_R(X)$. That is,
$$U_R(X) = \bigcup_{x \in U} \{R(x) : R(x) \cap X \neq \emptyset\}$$
- (iii) The boundary region of X with respect to R is the set of all objects, which can be neither in nor as not- X with respect to R and it is denoted by $B_R(X)$.
That is, $B_R(X) = U_R(X) - L_R(X)$

Definition 2.2: If (U, R) is an approximation space and $X, Y \subseteq U$, then

- (i) $L_R(X) \subseteq X \subseteq U_R(X)$
- (ii) $L_R(\emptyset) = U_R(\emptyset) = \emptyset, L_R(U) = U_R(U) = U,$
- (iii) $U_R(X \cup Y) = U_R(X) \cup U_R(Y)$

- (iv) $U_R(X \cup Y) \subseteq U_R(X) \cap U_R(Y)$
- (v) $L_R(X \cup Y) \supseteq L_R(X) \cup L_R(Y)$
- (vi) $L_R(X \cap Y) = L_R(X) \cap L_R(Y)$
- (vii) $L_R(X) \subseteq L_R(Y)$ and $U_R(X) \subseteq U_R(Y)$ whenever $X \subseteq Y$
- (viii) $U_R(X^c) = [L_R(X)]^c$ and $L_R(X^c) = [U_R(X)]^c$
- (ix) $U_R(U_R(X)) = L_R(U_R(X)) = U_R(X)$
- (x) $L_R(L_R(X)) = U_R(L_R(X)) = L_R(X)$

Definition 2.3: Let U be an universe, R be an equivalence relation on U and $\tau_R(X) = \{U, \phi, L_R(X), U_R(X), B_R(X)\}$ where $X \subseteq U$. $\tau_R(X)$ satisfies the following axioms

- (i) $U, \phi \in \tau_R(X)$
 - (ii) The union of the elements of any sub-collection of $\tau_R(X)$ is in $\tau_R(X)$
 - (iii) The intersection of the elements of any finite sub collection of $\tau_R(X)$ is in $\tau_R(X)$.
- Then $\tau_R(X)$ is called the nano topology on U with respect to X .

The space $(U, \tau_R(X))$ is the nano topological space. The elements of $\tau_R(X)$ are called nano open sets.

Definition 2.4: A subset A of $(U, \tau_R(X))$ is called if

- (i) Nano semi-open if $A \subseteq \text{Ncl}(\text{Nint}(A))$. [6] Nano semi closed if $\text{Ncl}(\text{Nint}(A)) \subseteq A$
- (ii) Nano-regular open set if $A = \text{Ncl}(\text{Nint}(A))$. [7] Nano-regular closed set if $\text{Ncl}(\text{Nint}(A)) = A$
- (iii) Nano semi-generalized closed [1](briefly Nano-sg-closed) set if $\text{Nscl}(A) \subseteq M$ whenever $A \subseteq M$ and M is semi-open in U . The complement of nano sg-closed set is called nano sg-open

Definition 2.5: A Nano topological space U is said to be connected if U cannot be expressed as the union two disjoint non empty nano open sets in U .

Definition 2.6: A Nano topological space U is said to be Nano-sg connected if U cannot be expressed as a disjoint union of two non empty Nano-sg open sets

Definition 2.7: A space $(U, \tau_R(X))$ is called a locally indiscrete space if every nano open set of U is nano closed in U .

Definition 2.8: Every nano open set is Nano-sg open and every nano closed set is Nano-sg closed.

3. TOTALLY NANO-SG-CONTINUOUS FUNCTIONS

Definition 3.1: A function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is called totally nano continuous if the inverse image of every nano open subset of $(V, \tau_R'(Y))$ is a nano clopen subset of $(U, \tau_R(X))$.

Definition 3.2: A function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is said to be totally Nano-sg-continuous, if the inverse image of every nano open subset of $(V, \tau_R'(Y))$ is a Nano-sg clopen subset of $(U, \tau_R(X))$.

Example 3.3:

$U = \{a, b, c, d\}$, with $U/R = \{\{a\}, \{c\}, \{b, d\}\}$ and

$X = \{a, b\}$ Then the nano topology, $\tau_R(X) = \{U, \phi, \{a\}, \{a, b, d\}, \{b, d\}\}$.

Nano sg-open sets are

$\{U, \phi, \{a, c, d\}, \{a, b, d\}, \{a, b, c\}, \{b, c, d\}, \{b, d\}, \{a, d\}, \{a, b\}, \{a, c\}, \{a\}, \{b\}, \{d\}\}$

Nano sg-closed sets are

$\{U, \phi, \{b\}, \{c\}, \{d\}, \{a\}, \{a, c\}, \{b, c\}, \{c, d\}, \{b, d\}, \{b, c, d\}, \{a, c, d\}, \{a, b, c\}\}$

$V = \{a, b, c, d\}$ With $V/R = \{\{a\}, \{b, c\}, \{d\}\}$

$Y = \{a, c\}$ Then the nano topology $\tau_R'(Y) = \{V, \phi, \{a\}, \{b, c\}, \{a, b, c\}\}$.

The identity function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$

Then $f^{-1}(V) = V, f^{-1}(\phi) = \phi, f^{-1}(\{a\}) = \{a\}, f^{-1}(\{b, c\}) = \{b, c\}, f^{-1}(\{a, b, c\}) = \{a, b, c\}$

Thus the inverse image of every nano open set in V is nano sg-clopen in U .

Theorem 3.4: Every totally nano continuous functions is totally Nano-sg continuous.

Proof: Let N be an Nano open set of $(V, \tau_R'(Y))$. Since f is totally nano continuous functions, $f^{-1}(V)$ is both Nano open and Nano closed in $(U, \tau_R(X))$. Since every Nano open set is Nano-sg open and every Nano closed set is Nano-sg closed. $f^{-1}(V)$ is both Nano-sg open and Nano-sg closed in $(U, \tau_R(X))$. Therefore f is totally Nano-sg continuous.

Remark 3.5: The converse of the above theorem need not be true.

Example 3.6:

Let $U = \{1, 2, 3, 4\}$ with $U/R = \{\{1\}, \{3\}, \{2, 4\}\}$.

Let $X = \{1, 2\} \subseteq U$. Then $\tau_R(X) = \{U, \phi, \{1\}, \{1, 2, 4\}, \{2, 4\}\}$.

Nano sg-open sets are

$NSGO = \{U, \phi, \{1\}, \{2\}, \{4\}, \{2, 4\}, \{1, 4\}, \{1, 2\}, \{1, 3\}, \{1, 3, 4\}, \{1, 2, 4\}, \{1, 2, 3\}, \{2, 3, 4\}\}$.

Let $V = \{1, 2, 3, 4\}$ with $V/R = \{\{1, 2\}, \{3\}, \{4\}\}$.

Let $Y = \{1, 4\} \subseteq V$. Then $\tau_R'(Y) = \{U, \phi, \{4\}, \{1, 2, 4\}, \{1, 2\}\}$.

Then the identity function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is totally Nano-sg continuous but not totally nano continuous function

Theorem 3.7: If f is a totally Nano-sg-continuous function from a Nano-sg-connected space U onto any space V , then V is an indiscrete space.

Proof: Suppose that V is not indiscrete. Let A be a proper non-empty nano open subset of V . Then $f^{-1}(A)$ is a proper non-empty Nano-sg-clopen subset of $(U, \tau_R(X))$ which is a contradiction to the fact that U is Nano-sg-connected.

Definition 3.8: A space U is said to be Nano-sg- T_2 if for any pair of distinct points x, y of U , there exist disjoint Nano-sg-open sets M and N such that $x \in M$ and $y \in N$.

Lemma 3.9: The Nano-sg closure of every Nano-sg-open set is Nano-sg-open.

Proof: Every nano regular open set is nano open and every nano open set is Nano-sg-open. Thus, every nano regular closed set is Nano - sgclosed. Now let A be any Nano-sg-open set. There exists an nano open set M such that $M \subset A \subset \text{cl}(M)$. Hence, we have $M \subset \text{Nano-sg-cl}(M) \subset \text{Nano-sg-cl}(A) \subset \text{Nano-sg-cl}(\text{cl}(M)) = \text{cl}(M)$ since $\text{cl}(M)$ is nano regular closed. Therefore, $\text{Nano-sg-cl}(A)$ is Nano-sg-open.

Theorem 3.10: A space U is Nano-sg- T_2 if and only if for any pair of distinct points x, y of U there exist Nano-sg-open sets M and N such that $x \in M$, and $y \in N$ and $\text{Nano-sgcl}(M) \cap \text{Nano-sgcl}(N) = \phi$.

Proof: Necessity: Suppose that U is Nano-sg- T_2 . Let x and y be distinct points of U . There exist Nano-sg-open sets M and N such that $x \in M$, $y \in N$ and $M \cap N = \phi$. Hence $\text{Nano-sgcl}(M) \cap \text{Nano-sgcl}(N) = \phi$ and by Lemma 3.9, $\text{Nano-sgcl}(M)$ is Nano-sg-open. Therefore, we obtain $\text{Nano-sgcl}(U) \cap \text{Nano-sgcl}(N) = \phi$.

Sufficiency: This is obvious.

Theorem 3.11: If $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is a totally Nano-sg continuous injection and V is T_0 then U is Nano-sg- T_2 .

Proof: Let x and y be any pair of distinct points of U . Then $f(x) \neq f(y)$. Since V is T_0 , there exists an open set M containing say, $f(x)$ but not $f(y)$. Then $x \in f^{-1}(M)$ and $y \notin f^{-1}(M)$. Since f is totally Nano-sg-continuous, $f^{-1}(M)$ is a Nano-sg-clopen subset of U . Also, $x \in f^{-1}(M)$ and $y \in U - f^{-1}(M)$. By Theorem 3.10, it follows that U is Nano-sg- T_2 .

Theorem 3.12: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ be a totally Nano-sg-continuous function and V be a T_1 -space. If A is a non-empty Nano-sg-connected subset of U , then $f(A)$ is a single point.

Definition 3.13: Let $(U, \tau_R(X))$ be a Nano topological space. Then the set of all points y in U such that x and y cannot be separated by a Nano-sg-separation of U is said to be the quasi Nano-sg-component of U .

Theorem 3.14: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ be a totally Nano-sg-continuous function from a nano topological space $(U, \tau_R(X))$ into a T_1 -space V . Then f is constant on each quasi Nano-sg-component of U .

Proof: Let x and y be two points of U that lie in the same quasi-Nano-sg-component of U . Assume that $f(x) = \alpha \neq \beta = f(y)$. Since V is T_1 , $\{\alpha\}$ is nano closed in V and so $V - \{\alpha\}$ is a nano open set. Since f is totally Nano-sg-continuous, therefore $f^{-1}(\{\alpha\})$ and $f^{-1}(V - \{\alpha\})$ are disjoint Nano-sg-clopen subsets of U . Further, $x \in f^{-1}(\{\alpha\})$ and $y \in f^{-1}(V - \{\alpha\})$, which is a contradiction in view of the fact that y belongs to the quasi Nano-sg component of x and hence y must belong to every Nano-sg-open set containing x .

4. SLIGHTLY NANO-SG CONTINUOUS FUNCTIONS

Definition 4.1: A function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is called slightly nano continuous if the inverse image of every nano clopen set in $(V, \tau_R'(Y))$ is open in $(U, \tau_R(X))$.

Definition 4.2: A function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is said to be slightly Nano-sg continuous if the inverse image of every clopen set in $(V, \tau_R'(Y))$ is Nano-sg-open in $(U, \tau_R(X))$.

Example 4.3:

Let $U = V = \{1, 2, 3, 4, 5\}$ with $U/R = \{\{1\}, \{2, 4\}, \{3, 5\}\}$.

Let $X = \{1, 2\} \subseteq U$.

Then $\tau_R(X) = \{U, \phi, \{1\}, \{1, 2, 4\}, \{2, 4\}\}$.

Nano sg-open sets are

$NSGO(U) = \{U, \phi, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{1, 5\}, \{2, 3\}, \{2, 4\}, \{2, 5\}, \{3, 4\}, \{3, 5\}, \{4, 5\}, \{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 4, 5\}, \{3, 4, 5\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 4, 5\}, \{1, 3, 4, 5\}, \{2, 3, 4, 5\}\}$

$V = \{1, 2, 3, 4, 5\}$ with $V/R = \{\{1\}, \{2, 4\}, \{3, 5\}\}$.

$Y = \{1, 5\} \subseteq V$

Then $\tau_R'(Y) = \{V, \phi, \{1\}, \{1, 3, 5\}, \{3, 5\}\}$.

$f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is said to be slightly Nano-sg continuous

Then the identity function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$

Then $f^{-1}(V) = U, f^{-1}(\phi) = \phi, f^{-1}(\{1\}) = \{1\}, f^{-1}(\{3, 5\}) = \{3, 5\}, f^{-1}(\{2, 4\}) = \{2, 4\}, f^{-1}(\{1, 3, 5\}) = \{1, 3, 5\}, f^{-1}(\{1, 2, 4\}) = \{1, 2, 4\}, f^{-1}(\{2, 3, 4, 5\}) = \{2, 3, 4, 5\}$

Theorem 4.4: For a function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$, the following statements are equivalent.

- (i) f is slightly Nano-sg-continuous.
- (ii) The inverse image of every nano clopen set N of V is Nano-sg-open in U .
- (iii) The inverse image of every nano clopen set N of V is Nano-sg-closed in U .
- (iv) The inverse image of every nano clopen set N of V is Nano-sg-clopen in U .

Proof:

(i) \Rightarrow (ii): Follows from the Definition 4.4.

(ii) \Rightarrow (iii): Let N be a nano clopen set in V which implies N^c is clopen in V . By (ii), $f^{-1}(N^c) = (f^{-1}(N))^c$ is Nano-sg-open in U . Therefore, $f^{-1}(N)$ is Nano-sg-closed in U .

(iii) \Rightarrow (iv): By (ii) and (iii), $f^{-1}(N)$ is Nano-sg-clopen in U .

(iv) \Rightarrow (i): Let N be a nano clopen set in V containing $f(x)$, by (iv), $f^{-1}(N)$ is Nano-sg clopen in U . Take $U = f^{-1}(N)$, then $f(M) \subseteq N$. Hence, f is slightly Nano-sg-continuous.

Theorem 4.5: Every slightly continuous function is slightly Nano-sg-continuous.

Proof: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ be a Nano-sg continuous function. Let N be a nano clopen set in V . Then, $f^{-1}(N)$ is Nano-sg-open and Nano-sg-closed in U . Hence, f is slightly Nano-sg-continuous.

Remark 4.6: The converse of the above theorem need not be true as can be seen from the following example

Example 4.7:

Let $U = \{1, 2, 3, 4\}$ with $U/R = \{\{1\}, \{2, 3\}, \{4\}\}$.

Let $X = \{1, 3\} \subseteq U$. Then $\tau_R(X) = \{U, \phi, \{1\}, \{2, 3\}, \{1, 2, 3\}\}$.

Nano sg-open sets are

$NSGO = \{U, \phi, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{1, 4\}, \{1, 2\}, \{1, 3\}, \{2, 3, 4\}, \{1, 3, 4\}, \{1, 2, 4\}, \{1, 2, 3\}\}$.

Let $V = \{1, 2, 3, 4\}$ with $V/R = \{\{1\}, \{2, 3\}, \{4\}\}$.

Let $Y = \{1, 4\} \subseteq U$. Then $\tau_R'(Y) = \{V, \phi, \{1, 4\}\}$.

$f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ under the condition $f(1) = 2, f(2) = 1, f(3) = 4, f(4) = 3$ is slightly Nano-sg- continuous but not slightly continuous function.

Theorem 4.8: Every contra Nano-sg continuous function is slightly Nano-sg continuous.

Proof: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ be a contra Nano-sg continuous function. Let N be a clopen set in V . Then, $f^{-1}(N)$ is Nano-sg- open in U . Hence, f is slightly Nano-sg continuous.

Theorem 4.9: If the function $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is slightly Nano-sg continuous and $(U, \tau_R(X))$ is Nano-sg $T_{1/2}$ space, then f is slightly continuous.

Proof: Let N be a nano clopen set in Z . Since g is slightly Nano-sg- continuous, $f^{-1}(N)$ is Nano-sg open in U . Since U is Nano-sg $T_{1/2}$ space, $f^{-1}(N)$ is nano open in U . Hence f is slightly continuous.

Theorem 4.19: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ and $g: (V, \tau_R'(Y)) \rightarrow (W, \tau_R''(Z))$ be functions. If f is surjective and pre Nano-sg-open and $(g \circ f): (U, \tau_R(X)) \rightarrow (W, \tau_R''(Z))$ is slightly Nano-sg continuous, then g is slightly Nano-sg-continuous.

Proof: Let N be a Nano clopen set in $(W, \tau_R''(Z))$. Since $(g \circ f): (U, \tau_R(X)) \rightarrow (W, \tau_R''(Z))$ is slightly Nano-sg -continuous, $f^{-1}(g^{-1}(N))$ is Nano-sg-open in U . Since, f is surjective and pre Nano-sg -open ($f^{-1}(g^{-1}(N))) = g^{-1}(N)$ is Nano-sg -open. Hence g is slightly Nano-sg-continuous.

Theorem 4.10: Let $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ and $g: (V, \tau_R'(Y)) \rightarrow (W, \tau_R''(Z))$ be functions. If f is surjective and pre Nano-sg-open and Nano-sg -irresolute, then $(g \circ f): (U, \tau_R(X)) \rightarrow (W, \tau_R''(Z))$ is slightly Nano-sg continuous if and only if g is slightly Nano-sg continuous.

Proof: Let N be a Nano clopen set in $(W, \tau_R''(Z))$. Since $(g \circ f): (U, \tau_R(X)) \rightarrow (W, \tau_R''(Z))$ is slightly Nano-sg continuous, $f^{-1}(g^{-1}(N))$ is Nano-sg-open in U . Since, f is surjective and pre Nano-sg open ($f^{-1}(g^{-1}(N))) = g^{-1}(N)$ is Nano-sg open in V . Hence g is slightly Nano-sg-continuous.

Conversely, let g is slightly Nano-sg continuous. Let N be a Nano clopen set in $(W, \tau_R''(Z))$, then g is Nano-sg open in V . Since, f is Nano-sg irresolute, $f^{-1}(g^{-1}(N))$ is Nano-sg-open in U . $(g \circ f): (U, \tau_R(X)) \rightarrow (W, \tau_R''(Z))$ is slightly Nanosg-continuous.

Theorem 4.11: If f is slightly Nano-sg continuous from a Nano-sg- connected space $(U, \tau_R(X))$ onto a space $(V, \tau_R'(Y))$ then V is not a discrete space.

Proof: Suppose that V is a discrete space. Let N be a proper non empty nano open subset of V . Since, f is slightly Nano-sg - continuous, $f^{-1}(N)$ is a proper non empty Nano-sg clopen subset of U which is a contradiction to the fact that U is Nano-sg-connected.

Theorem 4.12: If $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is a slightly Nano-sg continuous surjection and U is Nano-sg connected, then V is connected.

Proof: Suppose V is not connected, then there exists non empty disjoint Nano open sets M and N such that $V = M \cup N$. Therefore, M and N are Nano clopen sets in V . Since, f is slightly Nano-sg continuous, $f^{-1}(M)$ and $f^{-1}(N)$ are non empty disjoint Nano-sg open in U and $U = f^{-1}(M) \cup f^{-1}(N)$. This shows that U is not Nano-sg connected. This is a contradiction and hence, V is connected.

Theorem 4.13: If $f: (U, \tau_R(X)) \rightarrow (V, \tau_R'(Y))$ is a slightly Nano- sg - continuous and $(V, \tau_R'(Y))$ is locally indiscrete space then f is Nano- sg -continuous.

Proof: Let N be an Nano-open subset of V . Since, $(V, \tau_R'(Y))$ is a locally indiscrete space, N is Nano-closed in V . Since, f is slightly Nano- sg continuous, $f^{-1}(N)$ is Nano- sg -open in U . Hence, f is Nano- sg continuous.

REFERENCES

1. Bhuvaneswari and A.Ezhilarasi, On Nano semi-generalized and Nano generalized–semi closed sets in Nano Topological Spaces, International Journal of Mathematics and Computer Applications Research, (2014), 117-124.
2. S.Chandrasekar,M.Suresh,T.Rajesh kannan Nano sg -interior and Nano sg -closure in nano topological spaces Int.J.Mathematical Archive, 8(4), 2017, 94-100.
3. S Chandrasekar, A Atkinswestly, M, Sathyabama, $g\#b$ -closed sets in Topological paces. International Journal of Advanced Research (IJAR) 5(7) , 2722-2730
4. S Chandrasekar, M Sathyabama, A Atkinswestley $g\#$ b-Interior and $g\#$ b-Closure in Topological Spaces International Journal of Mathematics and its applications, 552(B)7 (2017)
5. S.Chandrasekar, M.Suresh and T.Rajesh Kannan, On Quasi Soft sg -open and Quasi Soft sg -closed Functions in Soft Topological Spaces International Journal of Mathematics and its Applications (239-246), Volume 5, Issue 2–B (2017), 239–246.
6. S.Chandrasekar, T Rajesh Kannan, M Suresh,, $\delta\omega\alpha$ -Closed Sets in Topological Spaces, Global Journal of Mathematical Sciences: Theory and Practical.9 (2), 103-116(2017).
7. S.Chandrasekar,T Rajesh Kannan, R.Selvaraj $\delta\omega\alpha$ closed functions and $\delta\omega\alpha$ Closed Functions in Topological Spaces, International Journal of Mathematical Archive 8(11) 2017.
8. M.LellisThivagar and Carmel Richard, On Nano forms of weakly open sets, International Journal of Mathematics and Statistics Invention, 1(1)(2013), 31-37.
9. M. L.Thivagar, “Generalization of pairwise α -continuous functions,” Pure and Applied Matematika Sciences, vol. 33, no. 1-2, pp. 55–63, 1991.
10. M.LellisThivagar and Carmel Richard, On Nano continuity, Mathematical Theory and Modeling, 3(7) (2013).
11. O.Njastad, On some classes of nearly open sets, Pacific J. Math. 15 (1965), 961–970.
12. O.Ravi, S. Ganesan and S. Chandrasekar On Totally sg -Continuity, Strongly sg -Continuity and Contra sg -Continuity Gen. Math. Notes, Vol. 7, No 1, Nov (2011), pp. 13-24.
13. O.Ravi, S. Ganesan and S. Chandrasekar, Almost ugs -closed functions and separation axioms. Bulletin of Mathematical Analysis and Applications, Vol.3 (2011), 165-177.
14. O. Ravi, G.Ramkumar and S. Chandrasekar On Quasi sg -open and Quasi sg -closed Functions Advances in Applied Mathematical Analysis.Vol 4 No1 (2009), pp. 73-78.
15. O.Ravi, S.Chandrasekar, S.Ganesan.(\tilde{g},s) continuous functions between topological spaces. Kyungpook Mathematical Journal, 51 (2011), 323-338.
16. O.Ravi, S.Chandrasekar, S.Ganesan On weakly πg -closed sets in topological spaces. Italian journal of pure and applied mathematics (36), 656-666 (2016).
17. Sakkraiveeranan Chandrasekar, Velusamy Banupriya, Jeyaraman Suresh Kumar, Properties and Applications Of $\theta g^*\alpha$ -Closed Sets In Topological Spaces, Journal of New Theory (18) 1-11(2017).
18. Selvaraj Ganesan, Sakkari Veeranan Chandrasekar Another quasi μS -open and quasi μS -closed functions Journal of New Theory, 15(75-80).

Source of support: Proceedings of National Conference March 1st - 2018, On “Recent Advances in Pure and Applied Mathematics (RAPAM - 2018)”, Organized by Department of Mathematics, Arul Anandar College (Autonomous), Madurai. Tamilnadu, India.