

MULTIPLICATIVE ATOM-BOND CONNECTIVITY AND MULTIPLICATIVE GEOMETRIC-ARITHMETIC INDICES OF CHEMICAL STRUCTURES IN DRUGS

V. R. KULLI

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

(Received On: 14-04-18; Revised & Accepted On: 28-05-18)

ABSTRACT

In Medical Science, the multiplicative connectivity indices are used in the analysis of drug molecular structures which are helpful for medical scientists and pharmaceutical scientists to find out the chemical, biological characteristics and medical information of drugs. They are also used in the nanoscience engineering applications. In this paper, the first multiplicative atom bond connectivity and multiplicative geometric-arithmetic indices of some important nanostar dendrimers which appeared in nanoscience. These results have a wide application prospect in medical, nanoscience and pharmaceutical engineering.

Keywords: nanoscience, multiplicative atom bond connectivity index, multiplicative geometric-arithmetic index, dendrimer.

Mathematics Subject Classification: 05C05, 05C12, 05C35.

1. INTRODUCTION

In the field of nanoscience, many researchers found that there is a close relationship between the biological and chemical characteristics and the molecular graph structure of nanomaterial itself. Therefore it emerges as a new branch of Theoretical Chemistry and it depends on the calculation of topological index to compute the characteristics of nanomaterials. A molecular graph is a simple graph related to the structure of a nanomaterial. Each vertex of a molecular structure of drug can be expressed by a molecular graph represents an atom of the molecule and its edges to the bonds between atoms. Then chemical molecular structure of drug can be expressed by a molecular graph. Concerning the definition of the topological index on the molecular graph, and corresponding medical, chemical properties of drugs can be studied by the topological index calculation. To understand the nature of the drug, this procedure does not depend on the laboratory equipments and reagents, that it saves the cost.

In this paper, we consider only a finite, simple connected graph G with vertex set $V(G)$ and edge set $E(G)$. A single number that can be used to characterize some property of the molecular graph is called a topological index. Several degree based topological indices like Randić index, sum connectivity index, product connectivity index, sum connectivity index and so on. There are some contributions on degree based indices of certain molecular structures and they can be referred to Alikhani *et al.* [1], Ashrafi *et al.* [2], Das *et al.* [3], Estrada *et al.* [5], Gao *et al.* [6,7], Husin *et al.* [11], Kulli [17] and Zhao *et al.* [25].

In [12], Kulli introduced the first multiplicative atom bond connectivity index and multiplicative geometric- arithmetic index of a graph as follows:

The first multiplicative atom bond connectivity index of a graph G is defined as

$$ABC_1II(G) = \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}}. \quad (1)$$

The multiplicative geometric-arithmetic index of a graph G is defined as

$$GAII(G) = \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)}. \quad (2)$$

Corresponding Author: V. R. Kulli

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

Several papers contributed to different kinds of multiplicative connectivity indices of special molecular structures, and they can be referred to Gao *et al.* [8], Gao *et al.* [9], Gao *et al.* [10], Liu *et al.* [24], and Kulli [13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23].

In this paper, the first multiplicative atom bond connectivity and multiplicative geometric-arithmetic indices for certain infinite families of nanostar dendrimers are computed. For more information about nanostar dendrimers see [4].

2. RESULTS FOR $NS_1[n]$ DENDRIMER NANOSTARS

In this section, we focus on the first class of nanostar dendrimer, denoted by $NS_1[n]$, where n is the steps of growth in this type of dendrimer. The graph of $NS_1[n]$ nanostar dendrimer is presented in Figure 1.




Figure-1: The molecular graph of $NS_1[n]$

Let G be the molecular graph of $NS_1[n]$ polypropylenimine octaamine dendrimer. By calculation, we obtain that G has $32 \times 2^n - 29$ edges. Also by calculation, we obtain that G has four types of edges based on the degree of end vertices of each edge as given in Table 1.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(1, 2)	(1, 3)	(2, 2)	(2, 3)
Number of edges	2×2^n	$4 \times 2^n - 4$	$12 \times 2^n - 11$	$14 \times 2^n - 14$

Table-1: Edge partition of $NS_1[n]$

In the following theorems, we compute the first multiplicative ABC index and multiplicative GA index of $NS_1[n]$.

Theorem 1: The first multiplicative atom bond connectivity index of polypropylenimine octaamine dendrimer $NS_1[n]$ is

$$ABC_1H(NS_1[n]) = \left(\frac{1}{\sqrt{2}}\right)^{28 \times 2^n - 25} \times \left(\frac{2}{3}\right)^{2 \times 2^n - 2}.$$

Proof: From equation (1) and using Table 1, we deduce

$$\begin{aligned} ABC_1H(NS_1[n]) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\ &= \left(\sqrt{\frac{1+2-2}{1 \times 2}}\right)^{2 \times 2^n} \times \left(\sqrt{\frac{1+3-2}{1 \times 3}}\right)^{4 \times 2^n - 4} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{12 \times 2^n - 11} \times \left(\sqrt{\frac{2+3-2}{2 \times 3}}\right)^{14 \times 2^n - 14} \\ &= \left(\frac{1}{\sqrt{2}}\right)^{28 \times 2^n - 25} \times \left(\frac{2}{3}\right)^{2 \times 2^n - 2}. \end{aligned}$$

Theorem 2: The multiplicative geometric-arithmetic index of polypropylenimine octaamine $NS_1[n]$ dendrimer nanostar is

$$GAH(NS_1[n]) = \left(\frac{2\sqrt{2}}{3}\right)^{2 \times 2^n} \times \left(\frac{\sqrt{3}}{2}\right)^{4 \times 2^n - 4} \times \left(\frac{2\sqrt{6}}{5}\right)^{14 \times 2^n - 14}.$$

Proof: From equation (2) and using Table (1), we deduce

$$\begin{aligned}
 GII(NS_1[n]) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\
 &= \left(\frac{2\sqrt{1 \times 2}}{1+2}\right)^{2 \times 2^n} \times \left(\frac{2\sqrt{1 \times 3}}{1+3}\right)^{4 \times 2^n - 4} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{12 \times 2^n - 11} \times \left(\frac{2\sqrt{2 \times 3}}{2+3}\right)^{14 \times 2^n - 14} \\
 &= \left(\frac{2\sqrt{2}}{3}\right)^{2 \times 2^n} \times \left(\frac{\sqrt{3}}{2}\right)^{4 \times 2^n - 4} \times \left(\frac{2\sqrt{6}}{5}\right)^{14 \times 2^n - 14}.
 \end{aligned}$$

3. RESULTS FOR $NS_2[n]$ NANOSTAR DENDRIMERS

In this section, we focus on the second class of nanostar dendrimer, denoted by $NS_2[n]$, where n is the steps of growth in this type of dendrimer. The graph of $NS_2[n]$ nanostar dendrimer is shown in Figure 2.




Figure-2: The structure of $NS_2[n]$

Let G be the molecular graph of $NS_2[n]$ polypropylenimine octaamine dendrimer. By calculation, we obtain that G has $16 \times 2^n - 11$ edges. Also by calculation, we obtain that G has three types of edges based on the degree of end vertices of each edge as given Table 2.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(1, 2)	(2,2)	(2,3)
Number of edges	2×2^n	$8 \times 2^n - 5$	$6 \times 2^n - 6$

Table-2: Edge partition of $NS_2[n]$

In the following theorems, we derive the first multiplicative ABC index and multiplicative GA index of $NS_2[n]$.

Theorem 3: The first multiplicative atom bond connectivity index of polypropylenimine octaamine dendrimer $NS_2[n]$ is

$$ABC_1II(NS_2[n]) = \left(\sqrt{\frac{1}{2}}\right)^{16 \times 2^n - 11}.$$

Proof: Using equation (1) and Table 2, we derive

$$\begin{aligned}
 ABC_1II(NS_2[n]) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\
 &= \left(\sqrt{\frac{1+2-2}{1 \times 2}} \right)^{2 \times 2^n} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}} \right)^{8 \times 2^n - 5} \left(\sqrt{\frac{2+3-2}{2 \times 3}} \right)^{6 \times 2^n - 6} = \left(\sqrt{\frac{1}{2}} \right)^{16 \times 2^n - 11}.
 \end{aligned}$$

Theorem 4: The multiplicative geometric-arithmetic index of polypropylenine octaamine dendrimer $NS_2[n]$ is

$$GAI(NS_2[n]) = \left(\frac{2\sqrt{2}}{3} \right)^{2 \times 2^n} \times \left(\frac{2\sqrt{6}}{5} \right)^{6 \times 2^n - 6}.$$

Proof: Using equation (2) and Table 2, we have

$$\begin{aligned}
 GII(NS_2[n]) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\
 &= \left(\frac{2\sqrt{1 \times 2}}{1+2} \right)^{2 \times 2^n} \times \left(\frac{2\sqrt{2 \times 2}}{2+2} \right)^{8 \times 2^n - 5} \times \left(\frac{2\sqrt{2 \times 3}}{2+3} \right)^{6 \times 2^n - 6} \\
 &= \left(\frac{2\sqrt{2}}{3} \right)^{2 \times 2^n} \times \left(\frac{2\sqrt{6}}{5} \right)^{6 \times 2^n - 6}.
 \end{aligned}$$

4. RESULTS FOR $NS_3[n]$ POLYMER DENDRIMERS

In this section, we focus on the class of nanostar dendrimer, denoted by $NS_3[n]$, where n is the steps of growth in this type of dendrimer. The graph of $NS_3[n]$ polymer dendrimer is presented in Figure 3.




Figure-3: The structure of polymer dendrimer $NS_3[n]$

Let G be the molecular graph of $NS_3[n]$ polymer dendrimer. By calculation, we obtain that $|E(NS_3[n])| = 69 \times 2^n + 90$. Also by calculation, we obtain that G has four types of edges based on the degree of the end vertices of each edge as given in Table 3.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(1, 2)	(2, 2)	(2, 3)	(3, 3)
Number of edges	3×2^n	$27 \times 2^n - 24$	$33 \times 2^n + 114$	6×2^n

Table-3: Edge partition of $NS_3[n]$

We now compute the first multiplicative ABC index and multiplicative GA index of $NS_3[n]$.

Theorem 5: The first multiplicative atom bond connectivity index of polymer dendrimer $NS_3[n]$ is

$$ABCI(NS_3[n]) = \left(\frac{1}{\sqrt{2}} \right)^{63 \times 2^n + 90} \times \left(\frac{2}{3} \right)^{6 \times 2^n}.$$

Proof: From equation (1) and using Table 3, we obtain

$$\begin{aligned} ABC_1II(NS_3[n]) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\ &= \left(\sqrt{\frac{1+2-2}{1 \times 2}}\right)^{3 \times 2^n} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{27 \times 2^n - 24} \times \left(\sqrt{\frac{2+3-2}{2 \times 3}}\right)^{33 \times 2^n + 114} \times \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^{6 \times 2^n} \\ &= \left(\frac{1}{\sqrt{2}}\right)^{63 \times 2^n + 90} \times \left(\frac{2}{3}\right)^{6 \times 2^n}. \end{aligned}$$

Theorem 6: The multiplicative geometric-arithmetic index of polymer dendrimer $NS_3[n]$ is

$$GAII(NS_3[n]) = \left(\frac{2\sqrt{2}}{3}\right)^{3 \times 2^n} \times \left(\frac{2\sqrt{6}}{5}\right)^{33 \times 2^n + 114}.$$

Proof: From equation (2) and using Table 3, we have

$$\begin{aligned} GII(NS_3[n]) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\ &= \left(\frac{2\sqrt{1 \times 2}}{1+2}\right)^{3 \times 2^n} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{27 \times 2^n - 24} \times \left(\frac{2\sqrt{2 \times 3}}{2+3}\right)^{33 \times 2^n + 114} \times \left(\frac{2\sqrt{3 \times 3}}{3+3}\right)^{6 \times 2^n} \\ &= \left(\frac{2\sqrt{2}}{3}\right)^{3 \times 2^n} \times \left(\frac{2\sqrt{6}}{5}\right)^{33 \times 2^n + 114}. \end{aligned}$$

5. RESULTS FOR $NS_4[n]$ FULLERENE DENDRIMERS

In this section, we consider the class of fullerene dendrimer, denoted by $NS_4[n]$, where n is the steps of growth in this type of dendrimer. The molecular graph of $NS_4[n]$ fullerene dendrimer is presented in Figure 4.




Figure-4: The structure of fullerene dendrimer $NS_4[n]$

Let G be the graph of $NS_4[n]$ fullerene dendrimer. By calculation, we obtain that G has $20 \times 2^n + 89$ edges. Also by calculation, we obtain that G has six types of edges based on the degree of the end vertices of each edge as given in Table 4.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(1, 2)	(2, 2)	(2,3)	(3,3)	(3,4)	(4,4)
Number of edges	2×2^n	$2 \times 2^n + 2$	$16 \times 2^n - 8$	86	6	3

Table-4: Edge partition of $NS_4[n]$

We now compute the first multiplicative ABC index and multiplicative GA index of $NS_4[n]$.

Theorem 7: The first multiplicative atom bond connectivity index of fullerene dendrimer $NS_4[n]$ is

$$ABC_1H(NS_4[n]) = \left(\frac{1}{2}\right)^{10 \times 2^n - 3} \times \left(\frac{2}{3}\right)^{86} \times \left(\frac{5}{12}\right)^3 \times \left(\frac{\sqrt{6}}{4}\right)^3.$$

Proof: From equation (1) and using Table 4, we derive

$$\begin{aligned} ABC_1H(NS_4[n]) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\ &= \left(\sqrt{\frac{1+2-2}{1 \times 2}}\right)^{2 \times 2^n} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{2 \times 2^n + 2} \times \left(\sqrt{\frac{2+3-2}{2 \times 3}}\right)^{16 \times 2^n - 8} \times \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^{86} \times \left(\sqrt{\frac{3+4-2}{3 \times 4}}\right)^6 \times \left(\sqrt{\frac{4+4-2}{4 \times 4}}\right)^3 \\ &= \left(\frac{1}{2}\right)^{10 \times 2^n - 3} \times \left(\frac{2}{3}\right)^{6 \times 2^n} \times \left(\frac{5}{12}\right)^3 \times \left(\frac{\sqrt{6}}{4}\right)^3. \end{aligned}$$

Theorem 8: The multiplicative geometric-arithmetic index of fullerene dendrimer $NS_4[n]$ is

$$GAH(NS_4[n]) = \left(\frac{2\sqrt{2}}{3}\right)^{2 \times 2^n} \times \left(\frac{2\sqrt{6}}{5}\right)^{16 \times 2^n - 8} \times \left(\frac{4\sqrt{3}}{7}\right)^6.$$

Proof: From equation (2) and using Table 4, we derive

$$\begin{aligned} GAH(NS_4[n]) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\ &= \left(\frac{2\sqrt{1 \times 2}}{1+2}\right)^{2 \times 2^n} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{2 \times 2^n + 2} \times \left(\frac{2\sqrt{2 \times 3}}{2+3}\right)^{16 \times 2^n - 8} \times \left(\frac{2\sqrt{3 \times 3}}{3+3}\right)^{86} \times \left(\frac{2\sqrt{3 \times 4}}{3+4}\right)^6 \times \left(\frac{2\sqrt{4 \times 4}}{4+4}\right)^3 \\ &= \left(\frac{2\sqrt{2}}{3}\right)^{3 \times 2^n} \times \left(\frac{2\sqrt{6}}{5}\right)^{16 \times 2^n - 8} \times \left(\frac{4\sqrt{3}}{7}\right)^6. \end{aligned}$$

6. RESULTS FOR POLYMER DENDRIMERS $NS_5[n]$

In this section, we consider the class of polymer dendrimer $NS_5[n]$, where n is the steps of growth. The molecular graph of $NS_5[n]$ polymer dendrimer is shown in Figure 5.




Figure-5: The structure of polymer dendrimer $NS_5[n]$

Let G be the graph of $NS_5[n]$ polymer dendrimer. By calculation, we obtain that $NS_5[n]$ has $60 \times 2^n + 27$ edges. Also by calculation, we obtain that $E(NS_5[n])$ can be divided into four partitions as given in Table 5.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(1, 3)	(2, 2)	(2,3)	(3,3)
Number of edges	$6 \times 2^n + 3$	$6 \times 2^n + 6$	$48 \times 2^n - 6$	24

Table-5: Edge partition of $NS_5[n]$

We now compute the first multiplicative ABC index and multiplicative GA index of $NS_5[n]$.

Theorem 9: The first multiplicative atom bond connectivity index of polymer dendrimer $NS_5[n]$ is

$$ABC_1II(NS_5[n]) = \left(\sqrt{\frac{2}{3}}\right)^{6 \times 2^n + 51} \times \left(\frac{1}{2}\right)^{27 \times 2^n}.$$

Proof: From equation (1) and using Table 5, we deduce

$$\begin{aligned} ABC_1II(NS_5[n]) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\ &= \left(\sqrt{\frac{1+3-2}{1 \times 3}}\right)^{6 \times 2^n + 3} \times \left(\sqrt{\frac{2+2-2}{2 \times 2}}\right)^{6 \times 2^n + 6} \times \left(\sqrt{\frac{2+3-2}{2 \times 3}}\right)^{48 \times 2^n - 6} \times \left(\sqrt{\frac{3+3-2}{3 \times 3}}\right)^{24} \\ &= \left(\sqrt{\frac{2}{3}}\right)^{6 \times 2^n + 51} \times \left(\frac{1}{2}\right)^{27 \times 2^n}. \end{aligned}$$

Theorem 10: The multiplicative geometric-arithmetic index of polymer dendrimer $NS_5[n]$ is

$$GAII(NS_5[n]) = \left(\frac{\sqrt{3}}{2}\right)^{6 \times 2^n + 3} \times \left(\frac{2\sqrt{6}}{5}\right)^{48 \times 2^n - 6}.$$

Proof: Using equation (2) and using Table 5, we deduce

$$\begin{aligned} GAII(NS_5[n]) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\ &= \left(\frac{2\sqrt{1 \times 3}}{1+3}\right)^{6 \times 2^n + 3} \times \left(\frac{2\sqrt{2 \times 2}}{2+2}\right)^{6 \times 2^n + 6} \times \left(\frac{2\sqrt{2 \times 3}}{2+3}\right)^{48 \times 2^n - 6} \times \left(\frac{2\sqrt{3 \times 3}}{3+3}\right)^{24} \\ &= \left(\frac{\sqrt{3}}{2}\right)^{6 \times 2^n + 3} \times \left(\frac{2\sqrt{6}}{5}\right)^{48 \times 2^n - 6}. \end{aligned}$$

7. RESULTS FOR NANOSTAR DENDRIMERS D_n

We now focus on the class of nanostar dendrimer, denoted by D_n , where n is the steps of growth. The graph of D_n nanostar dendrimer is depicted in Figure 6.




Figure-6: The graph of nanostar dendrimer D_n

Let G be the graph of D_n nanostar dendrimer. The graph G has $33 \times 2^n - 45$ edges. By calculation, we obtain that the edge set $E(D_n)$ can be divided into three partitions as given Table 6.

$d_G(u), d_G(v) \setminus uv \in E(G)$	(2, 2)	(2,3)	(3,3)
Number of edges	$12 \times 2^n - 12$	$15 \times 2^n - 24$	$6 \times 2^n - 9$

Table-6: Edge partition of D_n

In the following theorems, we determine the first multiplicative ABC index and multiplicative GA index of D_n .

Theorem 11: The first multiplicative atom bond connectivity index of nanostar dendrimer D_n is

$$ABC_1II(D_n) = \left(\sqrt{\frac{1}{2}} \right)^{27 \times 2^n - 36} \times \left(\frac{2}{3} \right)^{6 \times 2^n - 9}.$$

Proof: Using equation (1) and using Table 6, we obtain

$$\begin{aligned} ABC_1II(D_n) &= \prod_{uv \in E(G)} \sqrt{\frac{d_G(u) + d_G(v) - 2}{d_G(u)d_G(v)}} \\ &= \left(\sqrt{\frac{2+2-2}{2 \times 2}} \right)^{12 \times 2^n - 12} \times \left(\sqrt{\frac{2+3-2}{2 \times 3}} \right)^{15 \times 2^n - 24} \left(\sqrt{\frac{3+3-2}{3 \times 3}} \right)^{6 \times 2^n - 9} \\ &= \left(\sqrt{\frac{1}{2}} \right)^{27 \times 2^n - 36} \times \left(\frac{2}{3} \right)^{6 \times 2^n - 9}. \end{aligned}$$

Theorem 12: The multiplicative geometric-arithmetic index of nanostar dendrimer D_n is

$$GAII(D_n) = \left(\frac{2\sqrt{6}}{5} \right)^{15 \times 2^n - 24}.$$

Proof: Using equation (2) and Table 6, we obtain

$$\begin{aligned} GAII(D_n) &= \prod_{uv \in E(G)} \frac{2\sqrt{d_G(u)d_G(v)}}{d_G(u) + d_G(v)} \\ &= \left(\frac{2\sqrt{2 \times 2}}{2+2} \right)^{12 \times 2^n - 12} \times \left(\frac{2\sqrt{2 \times 3}}{2+3} \right)^{15 \times 2^n - 24} \times \left(\frac{2\sqrt{3 \times 3}}{3+3} \right)^{6 \times 2^n - 9} \\ &= \left(\frac{2\sqrt{6}}{5} \right)^{15 \times 2^n - 24}. \end{aligned}$$

REFERENCES

1. S. Alikhani, R.Hasni and N.E. Arif, On the atom bond connectivity index of some families of dendrimers, *J. Comput. Theor. Nanosci.* 11(8) (2014) 1802-1805.
2. A.R.Ashrafi and P. Nikzad, Connectivity index of the family of dendrimer nanostars, *Digest Journal of Nanomaterials and Biostructures*, 4(2) (2009) 269-273.
3. K.C.Das, I. Gutman and B. Furtula, On atom bond connectivity index, *Filomat*. 26(4) (2012) 733-738.
4. N.De and S.K.A Nayeem, Computing the F -index of nanostar dendrimers, *Pacific Science Review A: Natural Science and Engineering* (2016) <http://dx.doi.org/10.1016/j.psra.2016.06.01>.
5. E. Estrada, L. Torres, L. Rodriguez and I. Gutman, An atom bond connectivity index: modeling the enthalpy of formation of alkanes, *Indian J. Chem.* 37A (1998) 849-855.
6. W.Gao and W.F. Wang, The fifth geometric-arithmetic index of bridge graph and carbon nanocanes, *J. Differ. Eq. Appl.* <https://doi.org/1080/10236198:2016:1197214>.
7. W.Gao, Y.J. Chen and W.F. Wang, The topological variable computation for a special type of cycloalkanes, *J. Differ. Eq. Appl.* <https://doi.org/1080/10236198:2016:1197214>.
8. W.Gao, Y. Wang and W. Wang, and Li Shi, The first multiplication atom bond connectivity index of molecular structures in drugs, *Saudi Pharmaceutical Journal*, (2017), <http://dx.doi.org/10.1016/j.jsps.2017.04.021>.

9. W.Gao, M.K.Jamil, W.Nazeer and M.Amin, Degree based multiplicative atom bond connectivity index of nanostructures, *IAENG, International Journal of Applied Mathematics*, 47:4, IJAM-47-4-04, (2017).
10. W.Gao, W.Wang, D.Dimitrov and Y. Wang, Nano properties analysis via fourth multiplicative ABC indicator calculating, *Arabian Journal of Chemistry* (2018) <https://doi.org/10.1016/j.arabjc.2017.12.024>.
11. N.M. Husin, R. Hasni and N.E. Arif, Atom bond connectivity and geometric-arithmetic indices of dendrimer nanostars, *Australian Journal of Basic and Applied Sciences*, 7(9) (2013) 10-14.
12. V.R. Kulli, On multiplicative connectivity indices of certain nanotubes, *Annals of Pure and Applied Mathematics*, 12(2) (2016), 169-176.
13. V.R. Kulli, Multiplicative connectivity indices of $TUC_4C_8[m,n]$ and $TUC_4[m,n]$ nanotubes, *Journal of Computer and Mathematical Sciences*, 7(11) (2016) 599-605.
14. V.R.Kulli, Two new multiplicative atom bond connectivity indices, *Annals of Pure and Applied Mathematics*, 13(1) (2017) 1-7.
15. V.R.Kulli, Multiplicative connectivity indices of nanostructures, *Journal of Ultra Scientist of Physical Sciences*, A 29(1) (2017) 1-10.
16. V.R.Kulli, Some new multiplicative geometric-arithmetic indices, *Journal of Ultra Scientist of Physical Sciences*, A, 29(2) (2017) 52-57.
17. V.R. Kulli, Atom bond connectivity reverse and product connectivity reverse indices of oxide and honeycomb networks, *International Journal of Fuzzy Mathematical Archive*, 15(1) (2018) 1-5.
18. V.R.Kulli, Some new fifth multiplicative Zagreb indices of PAMAM dendrimers, *Journal of Global Research in Mathematical Archives*, 5(2) (2018) 82-86.
19. V.R. Kulli, Multiplicative connectivity Revan indices of polycyclic aromatic hydrocarbons and benzenoid systems, *Annals of Pure and Applied Mathematics*, 16(2) (2018) 337-343.
20. V.R. Kulli, Multiplicative connectivity Banhatti indices of benzenoid systems and polycyclic aromatic hydrocarbons, *Journal of Computer and Mathematical Sciences*, 9(3) (2018) 212-220.
21. V.R. Kulli, Multiplicative connectivity reverse indices of two families of dendrimer nanostars, *International Journal of Current Research in Life Sciences*, 7(2) (2018) 1102-1108.
22. V.R. Kulli, Multiplicative connectivity Revan indices of polycyclic aromatic hydrocarbons and benzenoid systems, *International Journal of Mathematical Archive*, 9(3) (2018) 235-241.
23. V.R.Kulli and M.H. Akhbari, Multiplicative atom bond connectivity and multiplicative geometric-arithmetic indices of dendrimer nanostars, *Annals of Pure and Applied Mathematics*, 16(2) (2018) 429-436. DOI:<http://dx.doi.org/10.22457/apam.v16n2a20>.
24. J.B. Liu, A.Q. Baig, W.Khalid and M.R. Farahani, Multiplicative indices of carbon graphine t-levels, *Compt. Rend. Acad. Bulg. Sci.* 7(1) (2018) 10-21.
25. B.Zhao and H.L.Wu, Pharmacological Characteristics analysis of two molecular structures, *Applied Mathematics and Nonlinear Sciences*, 2(1) (2017) 93-110.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]