

## ANTI Q-FUZZY B – IDEALS IN B – ALGEBRA

Dr. A. PRASANNA<sup>1</sup>, \*M. PREMKUMAR<sup>2</sup> AND HAJEE. Dr. S. ISMAIL MOHIDEEN<sup>3</sup>

<sup>1</sup>Assistant Professor, PG and Research Department of Mathematics,  
Jamal Mohamed College (Autonomous), Tiruchirappalli-620020, Tamilnadu, India.

<sup>2</sup>Research Scholar and Assistant Professor,  
Department of Mathematics, Mahendra Engineering College (Autonomous),  
Tiruchengode, Namakkal-637 503, Tamilnadu, India.

<sup>3</sup>Principal, Head and Associate Professor,  
PG and Research Department of Mathematics,  
Jamal Mohamed College (Autonomous), Tiruchirappalli-620020, Tamilnadu, India.

(Received On: 13-04-18; Revised & Accepted On: 28-05-18)

### ABSTRACT

*In this paper, we introduce the notion of anti Q-fuzzy B-Ideals of B-algebras, lower level B – Ideal and prove some results on these. We show that a Q-fuzzy subset of a B-algebra is a Q-fuzzy B-ideal if and only if the complement of this Q-fuzzy subset is an anti Q-fuzzy B-ideal.*

**Keywords:** B-algebra, B-Ideal, Fuzzy B-Ideal, Anti Fuzzy B-Ideal, Q-Fuzzy B-Ideal, Anti Q-Fuzzy B-Ideal.

### 1. INTRODUCTION

After the introduction of fuzzy subsets by L.A. Zadeh<sup>[1]</sup>, several researchers explored on the generalization of the notion of fuzzy subset. K. Atanassov<sup>[2]</sup>, introduced the Intuitionistic fuzzy sets. F. Adam and N. Hassan<sup>[1]</sup>, introduced the Q-fuzzy soft set. Muthuraj, P.M. Sitharselvam, M.S. Muthuraman<sup>[6]</sup>, introduced the notion Anti Q-fuzzy group and its lower level subgroups. J.R.Cho and H.S.Kim<sup>[4]</sup> discussed relations between B-algebras and other topics, especially quasi-groups. Jiayinpeng<sup>[5]</sup>, introduced the Intuitionistic Fuzzy B-algebras. H.K.Park and H.S.Kin<sup>[8]</sup>, introduced the notion of Quadratic B-algebras. Sun ShinAhn and KeumseongBang<sup>[9]</sup> have discussed the fuzzy subalgebra in B-algebra. C.Yamini and S.Kailasavalli<sup>[10]</sup>, introduced the notion of Fuzzy B-ideals. R.Biswas<sup>[3]</sup>, introduced the concept of anti-fuzzy subgroups of groups. P.Muthuraj, M.Sridharan, M.S.Muthuraman and P.M.SitharSelvam<sup>[7]</sup>, introduce the notion of Anti Q-Fuzzy BG-Ideals in BG-Algebra. Modifying their idea, in this paper, we apply the idea to B-Algebra. We introduce the notion of Anti Q-Fuzzy B-ideals of B-Algebras, lower level cuts of a Q-Fuzzy set, and prove some results on these.

### 2. PRELIMINARIES

In this section we give some basic definitions and preliminaries of B-algebras and introduce Q-fuzzy B-ideal.

**Definition 2.1:** (Jung R. Cho and H.S.Kim [4]) A B-algebra is a non-empty set  $X$  with a constant  $0$  and a binary operation  $*$  satisfying the following axioms:

- (i)  $x * x = 0$
- (ii)  $x * 0 = x$
- (iii)  $(x * y) * z = x * (z * (0 * y))$ , for all  $x, y, z \in X$

For brevity we also call  $X$  a B-algebra. In  $X$  we can define a binary relation " $\leq$ " by  $x \leq y$  if and only if  $x * y = 0$ .

**Definition 2.2:** (Jung R. Cho and H.S.Kim [4]) A non-empty subset  $M$  of a B-algebra  $X$  is called a sub-algebra of  $X$  if  $x * y \in M$  for any  $x, y \in M$ .

**Corresponding Author:** \*M. Premkumar<sup>2</sup>

<sup>2</sup>Research Scholar and Assistant Professor, Department of Mathematics, Mahendra Engineering College (Autonomous), Tiruchengode, Namakkal-637 503, Tamilnadu, India.

**Definition 2.3:** (JiayinPeng [5]) Let  $\alpha$  be a fuzzy set in a B-algebra. Then  $\alpha$  is called a fuzzy subalgebra of X if  $\alpha(x * y) \geq \alpha(x) \wedge \alpha(y)$  for all  $x, y \in X$ .

**Definition 2.4:** (Jung R. Cho and H.S.Kim [4]) A non-empty subset  $N$  of a B-algebra  $X$  is called a B-ideal of  $X$  if it satisfies for  $x, y, z \in X$

- (i)  $0 \in N$
- (ii)  $(x * y) \in N$  and  $(z * x) \in N$  implies  $(y * z) \in N$

**Definition 2.5:** (L. A. Zadeh [1]) Let  $X$  be a non-empty set. A fuzzy subset  $\alpha$  of the set  $X$  is a mapping  $\alpha: X \rightarrow [0,1]$

**Definition 2.6:** (F. Adam and N. Hassan [1]) Let  $Q$  and  $G$  be any two sets. A mapping  $A: G \times Q \rightarrow [0,1]$  is called a Q-fuzzy set in  $G$ .

**Definition 2.7:** (R.Muthuraj *et.al* [7]) Let  $\alpha$  be a Q-fuzzy set in B-algebra. Then  $\alpha$  is called a Q-fuzzy sub-algebra of  $X$  if  $\alpha(x * y, q) \geq \min\{\alpha(x, q), \alpha(y, q)\}$ , for all  $x, y \in X$  &  $q \in Q$

**Definition 2.8:** (R.Muthuraj *et.al* [7]) Let  $\alpha$  be a Q-fuzzy set in set  $X$ . Then the complement  $\bar{\alpha}$  is the Q-fuzzy subset of  $X$  given by  $\bar{\alpha}(x, q) = 1 - \alpha(x, q)$  for all  $x \in X$  &  $q \in Q$ .

**Definition 2.9:** (R. Muthuraj *et.al* [6]) Let  $\alpha$  be a Q-fuzzy set in a set  $X$ . For  $S \in [0,1]$ , the set  $\alpha_s = \{x \in X \mid \alpha(x, q) \geq S \text{ for all } q \in Q\}$  is called a level subset of  $\alpha$ .

**Definition 2.10:** (R.Muthuraj *et.al* [7]) A Q-fuzzy set  $\alpha$  in  $X$  is called Q-fuzzy B-ideal of  $X$  if it satisfies the following axioms:

- (i)  $\alpha(0, q) \geq \alpha(x, q)$
- (ii)  $\alpha(y * z, q) \geq \min\{\alpha(x * y, q), \alpha(z * x, q)\}$ , for all  $x, y \in X$  and  $q \in Q$ .

### 3. ANTI Q-FUZZY B-IDEALS

**Definition 3.1:** A Q-fuzzy set  $\alpha$  of a B-algebra  $X$  is called and Anti Q-Fuzzy subalgebra of  $X$  if  $\alpha(x * y, q) \leq \max\{\alpha(x, q), \alpha(y, q)\}$  for all  $x, y \in X$  and  $q \in Q$ .

**Definition 3.2:** A Q-fuzzy set  $\alpha$  in  $X$  is called an anti Q-fuzzy B-ideal of  $X$  if it satisfies the following axioms:

- (i)  $\alpha(0, q) \leq \alpha(x, q)$
- (ii)  $\alpha(y * z, q) \leq \max\{\alpha(x * y, q), \alpha(z * x, q)\}$ , for all  $x, y \in X$  and  $q \in Q$

**Theorem 3.3:** Every anti Q-fuzzy B-ideal of a B-algebra  $X$  is order preserving.

**Proof:** Let  $\alpha$  be an anti Q-fuzzy B-ideal of a B-algebra  $X$ .

Let  $x, y \in X$  and  $q \in Q$  be such that  $y \leq x$  if and only if  $y * x = 0$

Now,

$$\begin{aligned} \alpha(y, q) &= \alpha(0 * y, q) \\ &\leq \max\{\alpha(x * 0, q), \alpha(y * x, q)\} \\ &\leq \max\{\alpha(x, q), \alpha(0, q)\} \\ &\leq \alpha(x, q) \\ \Rightarrow \alpha(y, q) &\leq \alpha(x, q) \end{aligned}$$

**Theorem 3.4:** A Q-fuzzy subset  $\alpha$  of a B-algebra  $X$  is a Q-fuzzy B-ideal of  $X$  if and only if its complement  $\bar{\alpha}$  is an anti Q-fuzzy B-ideal of  $X$ .

**Proof:** Let  $\alpha$  be a Q-fuzzy B-ideal of  $X$  and let  $x, y, z \in X$  and  $q \in Q$ .

To prove:

$\bar{\alpha}$  is an anti Q-fuzzy B-ideal of  $X$ .

- (i)  $\bar{\alpha}(0, q) = 1 - \alpha(0, q)$   
 $\leq 1 - \alpha(x, q)$   
 $= \bar{\alpha}(x, q)$
- (ii)  $\bar{\alpha}(y * z, q) = 1 - \alpha(y * z, q)$   
 $\leq 1 - \min\{\alpha(x * y, q), \alpha(z * x, q)\}$   
 $\leq 1 + \max\{-\alpha(x * y, q), -\alpha(z * x, q)\}$   
 $\leq \max\{1 - \alpha(x * y, q), 1 - \alpha(z * x, q)\}$   
 $= \max\{\bar{\alpha}(x * y, q), \bar{\alpha}(z * x, q)\}$   
 $\Rightarrow \bar{\alpha}(y * z, q) \leq \max\{\bar{\alpha}(x * y, q), \bar{\alpha}(z * x, q)\}$

Thus,  $\bar{\alpha}$  is an anti Q-fuzzy B-ideal of X.

The converse part can also be prepared similarly.

Hence, the proof.

**Definition 3.5:** Let  $\alpha$  be a Q-fuzzy subset of a B-algebra X. For  $S \in [0,1]$ , the set  $\alpha^S = \{x \in X / \mu(x, q) \leq S\}$  is called a lower level cut of  $\alpha$ .

Clearly,  $\alpha' = X$  and  $\alpha^S \cup \alpha^S = X$  for  $S \in [0,1]$ . If  $S_1 \leq S_2$  then  $\alpha^{S_1} \subseteq \alpha^{S_2}$ .

**Theorem 3.6:** Let  $\alpha$  be a Q-fuzzy subset of a B-algebra X. If  $\alpha$  is an anti Q-fuzzy B-ideal of X, then the lower level cut  $\alpha^S$  is a B-ideal of X for all  $S \in [0,1]$ ,  $S \geq \alpha(0, q)$ .

**Proof:** Let  $\alpha$  be an anti Q-fuzzy B-ideal of X. Then for all  $x, y \in X$  and  $q \in Q$ ,

- (i)  $\alpha(0, q) \leq \alpha(x, q)$
- (ii)  $\alpha(y * z, q) \leq \max\{\alpha(x * y, q), \alpha(z * x, q)\}$

To prove:

$\alpha^S$  is a B-ideal of X.

Let  $x, y, z \in \alpha^S$

- $\Rightarrow \alpha(x, q) \leq S$
- (i) Since  $\alpha(0, q) \leq \alpha(x, q)$   
 $\leq S$   
 $\Rightarrow \alpha(0, q) \leq S$   
 $\Rightarrow 0 \in \alpha^S$
- (ii)  $x * y \in \alpha^S$  &  $z * x \in \alpha^S$   
 $\Rightarrow \alpha(x * y, q) \leq S$  &  $\alpha(z * x, q) \leq S$   
 $\alpha(y * z, q) \leq \max\{\alpha(x * y, q), \alpha(z * x, q)\}$   
 $\leq \max\{S, S\}$   
 $= S$   
 $\Rightarrow \alpha(y * z, q) \leq S$   
 $\Rightarrow y * z \in \alpha^S$

Thus,  $\alpha^S$  is a B-ideal of X.

Hence, the proof

**Theorem 3.7:** Let  $\alpha$  be a Q-fuzzy subset of a B-algebra X. If for each  $S \in [0,1]$ ,  $S \geq \alpha(0, q)$  the lower level cut  $\alpha^S$  is a B-ideal of X, then S is an anti Q-fuzzy B-ideal of X.

**Proof:**

$\alpha^S$  is a B-ideal of X.

$0 \in \alpha^S$

$x * y \in \alpha^S$  &  $z * x \in \alpha^S \Rightarrow y * z \in \alpha^S$

To prove

$\alpha^S$  is an anti Q-fuzzy B-ideal of X.

For all  $x, y \in X$  and  $q \in Q$

- (i)  $x * y \in \alpha^S$  &  $z * x \in \alpha^S$   
 $\Rightarrow \alpha(x * y, q) \leq S$  &  $\alpha(z * x, q) \leq S$

Let  $\alpha(x * y, q) = S$  &  $\alpha(z * x, q) = S$

$y * z \in \alpha^S$   
 $\Rightarrow \alpha(y * z, q) \leq S$   
 $= \max\{S, S\}$   
 $\leq \max\{\alpha(x * y, q), \alpha(z * x, q)\}$   
 $\Rightarrow \alpha(y * z, q) \leq \max\{\alpha(x * y, q), \alpha(z * x, q)\}$

- (ii)  $0 \in \mu^t$

Since  $x * x = 0$

$\alpha(0, q) = \alpha(x * x, q)$   
 $\leq \max\{\alpha(x, q), \alpha(x, q)\}$   
 $= \alpha(x, q)$

$\Rightarrow \alpha(0, q) \leq \alpha(x, q)$

$\Rightarrow \alpha^S$  is an anti Q-fuzzy B-ideal of S.

Hence, the proof.

#### 4. CONCLUSION

This paper tried to define the Anti Q-Fuzzy B-Ideals on B-Algebra and proved some theorems on them. This Concept can further be generalized to normalization of Q-fuzzy B-Ideals in B-Algebra using n-fold translation and multiplication.

#### 5. REFERENCES

1. F. Adam and N. Hassan, Q-fuzzy soft set, Applied Mathematical Sciences, 8(2014).
2. K. Atanassov, Intuitionistic Fuzzy Sets, Fuzzy Sets Systems 20(1986).
3. R.Biswas, Fuzzy Subgroups and Anti Fuzzy Subgroups, Fuzzy Sets and Systems (1990).
4. Jung R.Cho and H.S. Kim, On B-algebras and quasigroups, Quasigroups and related systems 7(2001).
5. JiayinPeng, Intuitionistic Fuzzy B-algebras, Research Journal of Applied Sciences, Engineering and Technology 4(2012).
6. R. Muthuraj, P.M.SitharSelvam and M.S.Muthuraman, Anti Q-fuzzy group and its lower level subgroups, International Journal of Computer Applications 3(2010).
7. R.Muthuraj, M.Sridharan, M.S.Muthuraman and P.M.SitharSelvam, Anti Q-Fuzzy BG-Ideals in BG-Algebra, International Journal of Computer Applications 4 (2010).
8. H.K.Park and H.S.Kim, On quadratic B-algebras, Quasigroups and related systems 7(2001).
9. Sun ShinAhn and Keumseong Bang, On Fuzzy Subalgebras in B-algebras, Commun.KoreanMath.Soc. 18(2003).
10. C.Yamini and S.Kailasavalli, Fuzzy B-ideals on B-algebras, International Journal of Mathematical Archive 5(2014).
11. L. A. Zadeh, Fuzzy Sets, Information Control 8 (1965).

**Source of support: Nil, Conflict of interest: None Declared.**

**[Copy right © 2018. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**