

FUZZY HYPONORMAL OPERATOR IN FUZZY HILBERT SPACE

A. RADHARAMANI¹ AND A. BRINDHA*²

¹Department of Mathematics, Chikkanna Govt. Arts College, India.

²Department of Mathematics, Tiruppur Kumaran College for Women, India.

(Received On: 26-11-18; Revised & Accepted On: 05-01-19)

ABSTRACT

In this paper we introduce the definition of Fuzzy Hyponormal Operator acting on a Fuzzy Hilbert space (FH-space) and investigate certain properties of Fuzzy Hyponormal operator using spectrum and we have given some definitions related to Fuzzy Hyponormal operator in FH-space.

Key words: Adjoint Fuzzy operator, Fuzzy Hilbert space, Fuzzy Normal operator, Fuzzy Hyponormal operator, Spectrum of a fuzzy bounded linear operator, Self-Adjoint Fuzzy operator.

1. INTRODUCTION

The notion of Fuzzy Norm on a linear space is first introduced by Katsaras[3] in 1984. Then after any other mathematicians have studied fuzzy normed space from several points of view [5], [12], [15]. The definition of fuzzy inner product space (FIP-space) headmost started by R.Biswas[10] and after that according the chronological in [6], [7], [11], [4], [9]. Modulate the definition of fuzzy inner product space (FIP-space) has been inserted by M.Goudarzi and S.M.Vaezpour in [8], [14]. The definition of a fuzzy Hilbert space has been introduced by M.Goudarzi and S.M.Vaezpour [8] in 2009. The concept of adjoint fuzzy linear operators, self - adjoint fuzzy linear operators was introduced by Sudad.M.Rasheed [2] in 2007, whereas A.Radharamani *et al.* [1] defined concept of fuzzy normal operator and their properties.

In this paper we consider fuzzy normal operator in FH-space and introduce the definitions of Fuzzy Hyponormal operator, we establish some theorems from fuzzy hyponormal operator in FH-space.

The classification of this paper is as follows:

In section 2 provides some preliminary definitions, theorems and results, which are used in this paper. In section 3 we introduce the concept of Fuzzy hyponormal operator on FH-space and some properties of fuzzy hyponormal operator have been studied.

2. PRELIMINARIES

Definition 2.1: [8] A fuzzy inner product space (FIP-Space) is a triplet $(X, F, *)$, where X is a real vector space, $*$ is a continuous t -norm, F is a fuzzy set on $X^2 \times \mathbb{R}$ satisfying the following conditions for every $x, y, z \in X$ and $s, r, t \in \mathbb{R}$.

FI-1: $F(x, x, 0) = 0$ and $F(x, x, t) > 0$, for each $t > 0$.

FI-2: $F(x, x, t) \neq H(t)$ for some $t \in \mathbb{R}$ if and only if $x \neq 0$, where $H(t) = \begin{cases} 1, & \text{if } t > 0 \\ 0, & \text{if } t \leq 0 \end{cases}$

FI-3: $F(x, y, t) = F(y, x, t)$

FI-4: For any $\alpha \in \mathbb{R}$, $F(\alpha x, y, t) = \begin{cases} F(x, y, \frac{t}{\alpha}) & \alpha > 0 \\ H(t) & \alpha = 0 \\ 1 - F(x, y, \frac{t}{-\alpha}) & \alpha < 0 \end{cases}$

FI-5: $F(x, x, t) * F(y, y, s) \leq F(x+y, x+y, t+s)$

FI-6: $\sup_{s+r=t} [F(x, z, s) * F(y, z, r)] = F(x+y, z, t)$

FI-7: $F(x, y, \cdot): \mathbb{R} \rightarrow [0, 1]$ is continuous on $\mathbb{R} \setminus \{0\}$

FI-8: $\lim_{t \rightarrow \infty} F(x, y, t) = 1$

Corresponding Author: A. Brindha*²

²Department of Mathematics, Tiruppur Kumaran College for Women, India.

Definition 2.2: [14] Let $(E, F, *)$ be probabilistic inner product space.

1. A sequence $\{x_n\} \in E$ is called \mathcal{T} -converges to $x \in E$, if for any $\epsilon > 0$ and $\lambda > 0$, $\exists N \in \mathbb{Z}^+$, $N = N(\epsilon, \lambda)$ such that $F_{x_n-x, x_n-x}(\epsilon) > 1-\lambda$, whenever $n > N$.
2. A linear Functional $f(x)$ defined on E is called \mathcal{T}_F -continuous, if $x_n \xrightarrow{\mathcal{T}_F} x$ implies $f(x_n) \xrightarrow{\mathcal{T}_F} f(x)$ for any $\{x_n\}, x \in E$

Theorem 2.3: [8] Let $(X, F, *)$ be a FIP-space, where $*$ is strong t -norm, and for each $x, y \in X$, $\text{Sup}\{t \in \mathbb{R} : F(x, y, t) < 1\} < \infty$. Define $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{R}$ by $\langle x, y \rangle = \text{Sup}\{t \in \mathbb{R} : F(x, y, t) < 1\}$. Then $(X, \langle \cdot, \cdot \rangle)$ is an IP-space (inner product space), so that $(X, \|\cdot\|)$ is a N-space (normed space), where $\|\cdot\| = \langle x, x \rangle^{1/2}$, $\forall x \in X$.

Definition 2.4: [8] Let $(X, F, *)$ be a FIP-space with IP $\langle x, y \rangle = \text{Sup}\{t \in \mathbb{R} : F(x, y, t) < 1\}$, $\forall x, y \in X$. If X is complete in the $\|\cdot\|$, then X is called Fuzzy Hilbert – space (FH-space).

Theorem 2.5: [8] Let $(X, F, *)$ be a FH-space with IP $\langle x, y \rangle = \text{Sup}\{t \in \mathbb{R} : F(x, y, t) < 1\}$, $\forall x, y \in X$ for $x_n \in X$ and $x_n \xrightarrow{\|\cdot\|} x$, then $x_n \xrightarrow{\mathcal{T}_F} x$.

Theorem 2.6: (Riesz theorem) [8], [14]

Let $(X, F, *)$ be FH-space. For any \mathcal{T}_F -continuous functional, \exists unique $y \in X$ such that for all $x \in X$, we have $g(x) = \text{Sup}\{t \in \mathbb{R} : F(x, y, t) < 1\}$.

Theorem 2.7: [1] Let $(E, G, *)$ be a FIP space, where $*$ is strong t -norm, and $\text{Sup}\{x \in \mathbb{R} : G(u, v, x) < 1\} < \infty$ for all $u, v \in E$, then $\text{Sup}\{x \in \mathbb{R} : G(u+v, w, x) < 1\} = \text{Sup}\{x \in \mathbb{R} : G(u, w, x) < 1\} + \text{Sup}\{x \in \mathbb{R} : G(v, w, x) < 1\}$, $\forall u, v, w \in E$.

Remark 2.8: [1] Let $FB(E)$ the set of all fuzzy linear operators on E .

Theorem 2.9: [1] (Adjoint Fuzzy operator in FH-space)

Let $(E, G, *)$ be a FH-space, Let $S \in FB(E)$ be \mathcal{T}_F -continuous linear functional, then \exists unique $S^* \in FB(E)$ such that $\langle Su, v \rangle = \langle u, S^*v \rangle$ $\forall u, v \in E$.

Definition 2.10: [1] Let $(E, G, *)$ be a FH-space with IP: $\langle u, v \rangle = \text{Sup}\{x \in \mathbb{R} : G(u, v, x) < 1\}$, $\forall u, v \in E$ and let $S \in FB(E)$, then S is self –adjoint Fuzzy operator. If $S = S^*$ where S^* is adjoint Fuzzy operator of S .

Theorem 2.11: [1] Let $(E, G, *)$ be a FH-space with IP: $\langle u, v \rangle = \text{Sup}\{x \in \mathbb{R} : G(u, v, x) < 1\}$ and let $S \in FB(E)$, then S is self –adjoint Fuzzy operator.

Theorem 2.12: [1] Let $(E, G, *)$ be a FH-space with IP: $\langle u, v \rangle = \text{Sup}\{x \in \mathbb{R} : G(u, v, x) < 1\}$, $\forall u, v \in E$ and let S^* be the adjoint Fuzzy operator of $S \in FB(E)$, then:

- i. $(S^*)^* = S$
- ii. $(\alpha S)^* = \alpha S^*$
- iii. $(\alpha S + \beta T)^* = \alpha S^* + \beta T^*$ where α, β are scalars and $T \in FB(E)$.
- iv. $(ST)^* = T^* S^*$

Theorem 2.13: [1] Let $(E, G, *)$ be a FH-space with IP: $\langle u, v \rangle = \text{Sup}\{x \in \mathbb{R} : G(u, v, x) < 1\}$, $\forall u, v \in E$ and let $S \in FB(E)$ then, $\|Su\| = \|S^*u\|$ for all $u \in E$.

Remark 2.14: Let $(H, F, *)$ be a FH – space with IP: $\langle x, y \rangle = \text{Sup}\{u \in \mathbb{R} : F(x, y, u) < 1\}$ $\forall x, y \in H$ and If $U \in FB(H)$, it is easy to see that $U = 0 \Leftrightarrow \langle Ux, y \rangle = 0, \forall x, y \in H$.

Theorem 2.15: Let $U \in FB(H)$. Then $U = 0 \Leftrightarrow \langle Ux, x \rangle = 0, \forall x \in H$.

Proof: It is obvious that $U = 0 \Rightarrow \langle Ux, x \rangle = 0, \forall x \in H$.

Conversely, it suffices to show that $\langle Ux, y \rangle = 0$, for any $x, y \in H$. The proof of this depends on the following easily verified identity,

$$\langle U(\alpha x + \beta y), \alpha x + \beta y \rangle - |\alpha|^2 \langle Ux, x \rangle - |\beta|^2 \langle Uy, y \rangle = \alpha \bar{\beta} \langle Ux, y \rangle + \bar{\alpha} \beta \langle Uy, x \rangle \quad (1)$$

By hypothesis equation (1) holds and equals 0, for all α and β .

If we put $\alpha = 1$ and $\beta = 1$, then $\langle Ux, y \rangle + \langle Uy, x \rangle = 0$ (2)

And if we put $\alpha = i$ and $\beta = 1$, then $i \langle Ux, y \rangle - i \langle Uy, x \rangle = 0$ (3)

Dividing (3) by i and adding the result to (2), we get

$$\langle Ux, y \rangle + \langle Uy, x \rangle + \langle Ux, y \rangle - \langle Uy, x \rangle = 0 \Rightarrow 2\langle Ux, y \rangle = 0, \quad \text{so } \langle Ux, y \rangle = 0 \Rightarrow U = 0.$$

The proof is complete.

Definition 2.16: [1] Fuzzy Normal Operator

Let $(X, F, *)$ be a FH-space with IP: $\langle u, v \rangle = \sup\{x \in R: F(u, v, x) < 1\} \forall u, v \in X$ and let $S \in FB(X)$. Then S is a fuzzy normal operator if it commutes with its (fuzzy) adjoint i.e. $SS^* = S^*S$.

Definition 2.17: [18] Fuzzy Invariant

Let $(X, F, *)$ be a fuzzy normed linear space, let $T \in FB(X)$, A subspace M of a fuzzy normed linear space X is said to be fuzzy invariant under T , if $TM \subset M$.

Theorem 2.18: [18] Let T be a normal operator on a finite dimensional fuzzy Hilbert space H over R then,

- (i) $T - \lambda I$ is fuzzy normal.
- (ii) Every eigen vector of T is also eigen vector for T^*

Theorem 2.19: [18] Let $(H, F, *)$ be a fuzzy Hilbert space and $T \in FB(H)$ then $T=0$ iff $\sup\{x \in R: F(Tu, Tu, x) < 1\} = 0 \forall u \in H$.

Theorem 2.20: [18] A Closed subspace M of a fuzzy Hilbert space H reduces an operator T iff M is invariant under T and T^* .

Definition 2.21: [18] Let M be a closed subspace of a fuzzy Hilbert space H , and let $T \in FB(H)$, we say that T is reduced by M if both M and M^\perp are invariant under T if T is reduced by M , then we also say that M reduces T .

Theorem 2.22: [18] A closed subspace M of a fuzzy Hilbert space H reduces an operator T iff M is invariant under both T and T^* .

Definition 2.23: [18] Let M be a closed of a fuzzy Hilbert space X and $x \notin M$ is said that projection of $x \in X$ onto M if $z \in M$, $N(x - z, t) = \sup\left\{\frac{t}{t + \|x - y\|}; y \in M, t > 0\right\}$, we write $y = P_M(x)$.

Theorem 2.24: [18] Let M is subspace of a fuzzy Hilbert space X , then $X = M \oplus M^\perp$ i.e each $x \in X$ can be uniquely decomposed from $x = y + z$ with $y \in M, z \in M^\perp$.

Theorem 2.25: [18] Let M be a closed subspace of a fuzzy Hilbert space X over F , and let $T \in FB(H)$, then M is invariant under T iff M^\perp is invariant under T^* .

Definition 2.26: [18] Let $(X, N, *)$ be a fuzzy normed linear space over the field \mathbb{C} , where $x \neq \{0\}$ and $N: X \times (0, \infty) \rightarrow [0, 1]$ and $T: (X, N, *) \rightarrow (X, N, *)$ be a linear operator,

A regular value (eigen value) λ of T is complex number \exists (i). $\mathcal{R}_\lambda(T)$ exist (ii). $\mathcal{R}_\lambda(T)$ is fuzzy bounded linear operator on range of $T_\lambda = T - \lambda I$ (iii). $\mathcal{R}_\lambda(T)$ is defined on a set which is dense in X , where $\mathcal{R}_\lambda(T) = (T_\lambda)^{-1} = (T - \lambda I)^{-1}$ is called resolvent operator of T . it is denoted by $\rho(T)$ and the resolvent set $\rho(T)$ of T is the set of all regular value λ of T . Its complement $\sigma(T) = \mathbb{C} - \rho(T)$ in complex plane \mathbb{C} is called spectrum of T , a $\lambda \in \sigma(T)$ is called spectral value of T .

3. MAIN RESULTS

In this section we introduce the definition of fuzzy hyponormal operator in FH-space as well as some elementary properties of fuzzy hyponormal operator in FH-space are presented.

Definition 3.1: Let $(H, F, *)$ be a FH – space with IP: $\langle x, y \rangle = \sup\{x \in R: F(u, v, x) < 1\} \forall u, v \in H$ and let $T \in FB(H)$. Then T is a fuzzy hyponormal operator if $\|T^*u\| \leq \|Tu\| \forall u \in H$ or equivalently $T^*T - TT^* \geq 0$.

Note: The term operator will mean fuzzy bounded linear transformation.

Theorem 3.2: Let $(H, F, *)$ be a FH – space with IP: $\langle x, y \rangle = \sup\{x \in R: F(u, v, x) < 1\} \forall u, v \in H$ and let $T \in FB(H)$ be a fuzzy hyponormal operator on H . Then $\|(T - zI)u\| \geq \|(T^* - \bar{z}I)u\| \forall u \in H$, i.e. $T - zI$ is FHN.

Proof: Given T be a FHN – operator on H .

$$\begin{aligned}
 \text{Let } \|(T - \lambda I)u\|^2 &= \langle (T - \lambda I)u, (T - \lambda I)u \rangle \\
 &= \sup\{x \in R: F((T - \lambda I)u, (T - \lambda I)u, x) < 1\} \\
 &= \sup\{x \in R: F(u, (T - \lambda I)^*(T - \lambda I)u, x) < 1\} \\
 &\geq \sup\{x \in R: F(u, (T - \lambda I)^*(T - \lambda I)^*u, x) < 1\} \quad [\text{since, by definition of FHN}] \\
 &\geq \langle (T - \lambda I)^*u, (T - \lambda I)^*u \rangle \\
 \therefore \|(T - \lambda I)u\|^2 &\geq \|(T - \lambda I)^*u\|^2 \\
 \Rightarrow \|(T - \lambda I)u\| &\geq \|(T - \lambda I)^*u\| \\
 \text{i.e. } \|(T - \lambda I)u\| &\geq \|(T - \lambda I)^*u\| \\
 \text{Thus } \|(T - \lambda I)u\| &\geq \|(T^* - \bar{\lambda}I)u\|
 \end{aligned}$$

Theorem 3.3: Let T be a FHN – operator on the Hilbert space H. Then $Tu = \lambda u \Rightarrow T^*u = \bar{\lambda}u$.

Proof: Let u be an eigen vector of T corresponding to the eigen value λ .
 $\Rightarrow Tu = \lambda u$

$$\begin{aligned}
 \text{Now } \sup\{x \in R: F(Tu, Tu, x) < 1\} &= \sup\{x \in R: F(u, T^*Tu, x) < 1\} \\
 &= \sup\{x \in R: F(u, TT^*u, x) < 1\} \quad [\text{since, T is Fuzzy Normal operator}] \\
 &= \sup\{x \in R: F(T^*u, T^*u, x) < 1\}
 \end{aligned}$$

Since $T - \lambda I$ is fuzzy hyponormal, therefore $u \in H$

$$\sup\{x \in R: F((T - \lambda I)u, (T - \lambda I)u, x) < 1\} \geq \sup\{x \in R: F((T - \lambda I)^*u, (T - \lambda I)^*u, x) < 1\}$$

Since $Tu = \lambda u$

$$\Rightarrow Tu = \lambda u \Rightarrow Tu - \lambda u = 0 \Rightarrow (T - \lambda I)u = 0$$

$$\therefore T - \lambda I = 0.$$

$$\sup\{x \in R: F((T - \lambda I)u, (T - \lambda I)u, x) < 1\} = 0 \quad \forall u \in H \quad (3.1)$$

$$\begin{aligned}
 \text{From (2.19) } T = 0 \text{ iff } \sup\{x \in R: F(Tu, Tu, x) < 1\} &= 0 \\
 \Rightarrow \sup\{x \in R: F((T - \lambda I)^*u, (T - \lambda I)^*u, x) < 1\} &= 0 \quad \forall u \in H
 \end{aligned}$$

From (3.1) $(T - \lambda I)^* = 0$ then for each $u \in H$

$$\begin{aligned}
 (T - \lambda I)^*u &= 0 \\
 \Rightarrow (T^* - \bar{\lambda}I)u &= 0 \\
 \Rightarrow T^*u - \bar{\lambda}u &= 0 \\
 \Rightarrow T^*u &= \bar{\lambda}u
 \end{aligned}$$

\therefore u is an eigen vector of T^* corresponding to eigen value $\bar{\lambda}$.

Theorem 3.4: Let T be fuzzy hyponormal iff $\|T^*u\| \leq \|Tu\|$ for all $u \in H$.

Proof: Since T is a fuzzy hyponormal, $T^*T - TT^* \geq 0$

$$\begin{aligned}
 &\Rightarrow T^*T \geq TT^* \\
 \text{i.e. } TT^* &\leq T^*T
 \end{aligned}$$

$$\begin{aligned}
 \text{Let } \|T^*u\| &\leq \|Tu\| \\
 \Leftrightarrow \|T^*u\|^2 &\leq \|Tu\|^2 \\
 \Leftrightarrow \langle T^*u, T^*u \rangle &\leq \langle Tu, Tu \rangle \\
 \Leftrightarrow \sup\{x \in R: F(T^*u, T^*u, x) < 1\} &\leq \sup\{x \in R: F(Tu, Tu, x) < 1\} \\
 \Leftrightarrow \sup\{x \in R: F(TT^*u, u, x) < 1\} &\leq \sup\{x \in R: F(T^*Tu, u, x) < 1\} \\
 \Leftrightarrow \langle TT^*u, u \rangle &\leq \langle T^*Tu, u \rangle \\
 \Leftrightarrow \langle (TT^* - T^*T)u, u \rangle &\leq 0 \\
 \Leftrightarrow TT^* - T^*T &\leq 0 \\
 \Leftrightarrow TT^* &\leq T^*T \\
 \text{i.e. } T^*T - TT^* &\geq 0
 \end{aligned}$$

Theorem 3.5: Let T be FHN on H. Then $\|T\| = R_{sp}(T)$ (the spectral radius of T).

Proof: For $u \in H$, $\|u\| = 1$,

$$\begin{aligned}
 \|Tu\|^2 &= \langle Tu, Tu \rangle \\
 &= \sup\{x \in R: F(Tu, Tu, x) < 1\}, u \in H \\
 &= \sup\{x \in R: F(T^*Tu, u, x) < 1\}, u \in H \\
 &= \langle T^*Tu, u \rangle \\
 &\leq \|T^*Tu\| \\
 \therefore \|Tu\|^2 &\leq \|T^2u\|
 \end{aligned}$$

But then $\|T\|^2 \leq \|T^2\| \leq \|T\|^2$ which implies that $\|T\|^2 = \|T^2\|$.

$$\begin{aligned} \text{Now } \|T^n u\|^2 &= \langle T^n u, T^n u \rangle \\ &= \langle T^n u, T T^{n-1} u \rangle \\ &= \langle T^* T^n u, T^{n-1} u \rangle \\ &\leq \|T^* T^n u\| \|T^{n-1} u\| \\ &\leq \|T^* T T^{n-1} u\| \|T^{n-1} u\| \\ &\leq \|T^{n-1} u\| \|T^{n-1} u\| \end{aligned}$$

Then $\|T^n\|^2 \leq \|T^{n-1}\| \|T^{n-1}\|$

And combining this with the equality, a simple induction argument yields,
 $\|T^n\| = \|T\|^n$ for $n=1,2,3, \dots$

Since $R_{sp}(T) = \lim_{n \rightarrow \infty} \|T^n\|^{\frac{1}{n}}$
 $R_{sp}(T) = \lim_{n \rightarrow \infty} \|T\|$
 i.e. $R_{sp}(T) = \|T\|$ where $R_{sp}(T)$ means the spectral radius of T.

Theorem 3.6: Let $T \in FB(H)$ be a fuzzy hyponormal operator with $Tu_1 = \lambda_1 u_1$, $Tu_2 = \lambda_2 u_2$ and $\lambda_1 \neq \lambda_2$ then $\langle u_1, u_2 \rangle = 0$.

Proof: Since T be a fuzzy hyponormal operator with $Tu_1 = \lambda_1 u_1$, $Tu_2 = \lambda_2 u_2$ and $\lambda_1 \neq \lambda_2$ then by theorem (3.3)
 $T^* u_1 = \bar{\lambda}_1 u_1$ and $T^* u_2 = \bar{\lambda}_2 u_2$.

$$\begin{aligned} \text{Let } \lambda_1 \langle u_1, u_2 \rangle &= \langle \lambda_1 u_1, u_2 \rangle \\ &= \sup\{x \in R: F(\lambda_1 u_1, u_2, x) < 1\} \\ &= \sup\{x \in R: F(Tu_1, u_2, x) < 1\} \\ &= \sup\{x \in R: F(u_1, T^* u_2, x) < 1\} \\ &= \sup\{x \in R: F(u_1, \bar{\lambda}_2 u_2, x) < 1\} \\ &= \sup\{x \in R: F(\lambda_2 u_1, u_2, x) < 1\} \\ &= \langle \lambda_2 u_1, u_2 \rangle \\ &= \lambda_2 \langle u_1, u_2 \rangle \end{aligned}$$

Hence if $\lambda_1 \neq \lambda_2$ then $\langle u_1, u_2 \rangle = 0$. i.e. $u_1 \perp u_2$.

Theorem 3.7: Let $(H, F, *)$ be a FH-space with $IP: \langle u, v \rangle = \sup\{x \in R: F(u, v, x) < 1\} \forall u, v \in H$ and let $T \in FB(H)$ is a fuzzy hyponormal operator on H with
 $M \subset H$ invariant under T and let T_M be fuzzy hyponormal then M reduces T

Proof: Let $u \in M$ the eigen space of T, and corresponding eigen value be λ . so that $Tu = \lambda u$. Since T is fuzzy normal then by theorem 2.8 eigen values for T^* . that is $T^* u = \bar{\lambda} u$, $u \in H$.

Since M is a subspace, $\bar{\lambda} u \in M \Rightarrow T^* u \in M$
 $\Rightarrow M$ is invariant under T^* , but M is invariant under T.

By a known theorem 2.20, M reduces T.

Corollary 1: Let T be a fuzzy hyponormal on H and
 $M = \{x \in H; Tu = \lambda u\}$ then M reduces T and T_M is fuzzy normal.

Corollary 2; Let T be a fuzzy hyponormal on H and $M \subset H$ invariant under T then T_M is fuzzy hyponormal.

Theorem 7: Let T be a fuzzy hyponormal on H and λ_0 be an isolated point in the spectrum of T then $\lambda_0 \in \sigma_p(T)$ the point spectrum of T.

Proof: By theorem 1, we may assume that $\lambda_0 = 0$, choose $R > 0$ sufficiently small that 0 is the only point of $\sigma(T)$, contained in or on the circle $|\lambda| = R$

Define $E = \int_{|\lambda|=R} (T - \lambda I)^{-1} d\lambda$ then E is a non zero projection which commutes with T.

Thus EH is invariant under T and T_{EH} is fuzzy hyponormal, also $\sigma(T_{EH}) = \sigma(T) \cap \{|\lambda| = R\}$

Also $\sigma(T_{EH}) = \{0\}$

From the last corollary we may conclude that T_{EH} is the zero transformation.

It is clear that $EH = \{ u \in H; Tu = 0 \} \Rightarrow EH$ reduces T .

Theorem 8: Let $(H, F, *)$ be a FH-space with $IP: \langle u, v \rangle = \sup\{x \in R: F(u, v, x) < 1\} \forall u, v \in H$ and let $T \in FB(H)$ is a fuzzy hyponormal operator on H with a single limit point in its spectrum then T is fuzzy hyponormal.

Proof: We may assume that by theorem 1, that the limit point is zero.

By theorem 7, there exist an isolated point in the $\lambda_1 \in \sigma(T) \ni |z_1| = \|T\|$

Let $M_1 = \{ u \in H; Tu = \lambda_1 u \}$

M_1 is not empty by theorem 7, and since M_1 reduces T . we conclude from theorem 6, that $T_{M_1^\perp}$ does not have λ_1 in its spectrum. By corollary 2 that T_{M_1} is fuzzy normal.

We continue in this way, selecting points in $\sigma(T)$ ordered by absolute value, setting $M_i = \{ u \in H; Tu = \lambda_i u \}$ then $M_1 \oplus M_2 \oplus \dots \oplus M_n$ reduces T and $T_{M_1 \oplus M_2 \oplus \dots \oplus M_n}$ is fuzzy normal we observe that $T_{M_1 \oplus M_2 \oplus \dots \oplus M_n^\perp}$ is fuzzy hyponormal with its spectral radius equal to its fuzzy norm.

Thus, since 0 is the only point of $\sigma(T)$, the fuzzy hyponormal operator converge to T in the uniform operator topology. Hence T is fuzzy normal.

Cor: If T is an fuzzy hyponormal, completely continous operator then T is fuzzy normal

4. CONCLUSION

As the idea of fuzzy hyponormal operator in FH-space is relatively new and classic form of theorems play a role the prototype in our discussion of this paper. Some concepts and properties have been investigated about fuzzy hyponormal operator in fuzzy Hilbert space. The results of this paper will be helpful for researchers to develop fuzzy functional analysis.

ACKNOWLEDGEMENT

The authors are grateful to the referees for these valuable and constructive suggestions.

REFERENCES

1. A.Radharamani,A.Brindha,S.Maheswari, "Fuzzy Normal Operator in fuzzy Hilbert space and its properties", IOSR Journal of Engineering, Vol.8,no.7, pp.1-6, 2018.
2. Sudad.M.Rasheed, "Self- adjoint Fuzzy Operator in Fuzzy Hilbert Space and its Properties", Journal of Zankoy Sulaimani, Vol.19, no.1, pp. 233-238, 2017.
3. K.Katsaras, "Fuzzy topological vector space-II", Fuzzy Sets and Systems, Vol.12, pp.143-154, 1984.
4. B.Punnose and S.Kuriakose, "Fuzzy inner product space-A New Approach", Journal of Fuzzy math, Vol.14, no.2, pp.273-282, 2006.
5. C.Felbin, "Finite dimensional Fuzzy normedlinear space", Fuzzy Sets and Systems, Vol.48, pp.239-248, 1992.
6. J.K.Kohil and R.Kumar, "Linear mappings, Fuzzy linear spaces, Fuzzy inner product spaces and Fuzzy Co-inner product spaces", Bull Calcutta Math.Soc., Vol.87, pp.237-246, 1995.
7. J.K.Kohil and R.Kumar, "On Fuzzy Inner Product Spaces and Fuzzy Co- Inner Product Spaces", Fuzzy sets and system, Bull Calcutta Math.Soc., Vol.53, pp.227-232,1993.
8. M.Goudarzi and S.M.Vaezpour, "On the definition of Fuzzy Hilbert spaces and its Applications", J. NonlinearSci.Appl, Vol.2, no.1, pp.46-59, 2009.
9. P.Majumdar and S.K.Samanta, "On Fuzzy inner product spaces", J.Fuzzy Math., Vol.16, no.2, pp.377-392, 2008.
10. R.Biswas, "Fuzzy inner product spaces &Fuzzy norm functions", Information Sciences, Vol.53, pp.185-190, 1991.
11. R.Saadati and S.M.Vaezpoor, "Some results on fuzzy Banach spaces", J.appl.Math. and computing, Vol.17, no.1, pp.475-488,2005.
12. S.C.Cheng, J.N.Mordeson, "Fuzzy linear operators and Fuzzy normed linear spaces", Bull.Cal.Math.Soc, Vol.86, pp.429-436,1994.
13. T.Bag, S.K.Samanta, Operators Fuzzy Norm and some properties Fuzzy", Inf.Eng.Vol.7, pp.151-164, 2015.
14. Yongfusu., "Riesz Theorem in probabilistic inner product spaces", Inter.Math.Forum.,Vol.2,no.62,pp.3073-3078, 2007.

15. T.Bag, S.K.Samanta, "Finite Dimensional fuzzy normed linear spaces", J.Fuzzy Math, Vol.11, no.3,687-705, 2003.
16. G.F.Simmons, Introduction to Topology and Modern Analysis,(New Delhi:Tata McGraw-Hill,1963), pp.222,273-274.
17. Balmohan V Limaye, Function Analysis, (New Delhi: New Age International, 1996), pp.460-469.
18. Noori F.AI- Mayahi, Abbas M.Abbas, some properties of spectral theory in fuzzy Hilbert spaces, journal of AL-Q adisiyah for computer science and mathematics,8(2),2016,1-7

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2019. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]