

NON-EXISTENCE OF COSMIC STRINGS IN BIANCHI TYPE-V COSMOLOGY

DINKAR SINGH CHAUHAN*

**Department of Mathematics,
Mahadev Post-graduate College, Varanasi (U.P.) India.**

(Received On: 20-03-19; Revised & Accepted On: 03-05-19)

ABSTRACT

Bianchi type-V string cosmological model is investigated in the presence and absence of bulk viscous fluid in general relativity. It has been shown that cosmic strings do not exist in Bianchi type-V cosmology. The physical and geometrical aspects of the model are also discussed.

Keywords: Bianchi type-V model, Bulk viscous fluid, cosmic strings.

1. INTRODUCTION

The study of Bianchi type-V cosmological models plays an important role in the study of Universe and the study is more interesting as these models contain isotropic special cases and permit arbitrary small anisotropy levels at some point of time. The concept of string theory was developed to describe events of the early stage of the evolution of the Universe. The general relativistic treatment of strings was initially done by Stachel [28] and Letelier [15, 16]. Letelier [15] obtained the general solution of Einstein's field equations for a cloud of strings with spherical, plane and a particular case of cylindrical symmetry. Letelier [16] also obtained massive string cosmological models in Bianchi type-I and Kantowski-Sachs space-times. The gravitational effects of cosmic strings have been extensively discussed by Vilenkin [30] and Gott [10] in general relativity. The string cosmological models with magnetic field are investigated by Chakraborty [5], Tikeler & Patel [29]. Banerjee *et al.* [2] have investigated an axially symmetric Bianchi type-I string dust cosmological model with and without magnetic field. Krori *et al.* [12] and Wang [31] studied the exact solutions of string cosmology for Bianchi type-II, VI₀, VII and IX space-times.

Cosmic strings are important in the early stages of evolution of the universe before the particle creation. The present day observations do not rule out the possible existence of large scale networks of strings in the early universe. Reddy [21] have shown that, the cosmic strings which have received considerable attention in cosmology do not exist in the framework of Rosen's [23] bimetric theory of gravitation. Krori *et al.* [13] and Deo *et al.* [8] have shown that, the cosmic strings do not occur in Bianchi type-V cosmology.

The role of viscosity in cosmology has been investigated by Weinbergs [32], Nightingale [18] and Heller & Klimek [11]. Heller and Klimek [11] have investigated viscous fluid cosmological model without initial singularity. They have shown that the introduction of bulk viscosity effectively removes the initial singularity. Roy and Singh [22] have investigated LRS Bianchi type-V cosmological model with viscosity. Santos *et al.* [26] investigated isotropic homogeneous cosmological model with bulk viscosity assuming viscous coefficient as power function of mass density. Banerjee and Sanyal [3] have investigated Bianchi type-V cosmological models with viscosity and heat flow. Pradhan *et al.* [19] have found the integrability of cosmic string in Bianchi type-III space time in presence of bulk viscous fluid. Bianchi type-V cosmological models have been studied by several researchers such as Farnsworth [9], Collins [7], Maartens and Nel [17], Beesham [4], Camci *et al.* [6], Aydogdu and Salti [1], in different physical contexts. Following the work of Saha [24], Singh and Chaubey [27] have obtained the quadrature form of metric function for Bianchi type-V model with perfect fluid and viscous fluid.

In this paper, we have investigated non-existence of cosmic strings in Bianchi type-V cosmological model with and without bulk viscous fluid. This paper is organized as follows: The metric and field equations with bulk viscous fluid are presented in Section 2. Subsection 2.1 deals with solution of the field equations. Some geometrical and physical properties of the model is presented in Subsection 2.2. Field equations without bulk viscous fluid are presented in Section 3. Subsection 3.1 deals with solution of the field equations. Some geometrical and physical properties of the model is presented in Subsection 3.2 Finally the conclusions of the paper are presented in Section 4.

Corresponding Author: Dinkar Singh Chauhan*

2. STRING COSMOLOGICAL MODEL WITH BULK VISCOUS FLUID

We consider a spatially homogeneous and anisotropic Bianchi type-V metric in the form

$$ds^2 = -dt^2 + A^2 dx^2 + e^{2mx} [B^2 dy^2 + C^2 dz^2], \quad (1)$$

where the metric potentials A, B and C are function of cosmic time t alone and m is a constant.

The energy-momentum tensor for bulk viscous string dust is given by Letelier [16], Landau and Lifshitz [14] as

$$T_i^j = \rho v_i v^j - \lambda x_i x^j - \xi v_{;\ell}^\ell (g_i^j + v_i v^j), \quad (2)$$

where ρ is the proper energy density of the cloud of string with particle attached to them, λ is the string tension density and ξ is the coefficient of bulk viscosity. $v^i = (0, 0, 0, 1)$ is the four velocity of the particle and x^i is a unit space-like vector representing the direction of string. The vectors v^i and x^i satisfy the conditions

$$v_i v^i = -x_i x^i = -1, \quad v_i x^i = 0. \quad (3)$$

Choosing x^i parallel to $\frac{\partial}{\partial x}$, we have

$$x^i = \left(\frac{1}{A}, 0, 0, 0 \right). \quad (4)$$

If the particle density of the configuration is denoted by ρ_p , then

$$\rho = \rho_p + \lambda \quad (5)$$

The Einstein's field equations (in gravitational units $8\pi G = c=1$) are given by

$$R_i^j - \frac{1}{2} R g_i^j = -T_i^j. \quad (6)$$

The field equation (6) with Equation (2) subsequently lead to the following system of equations:

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} - \frac{m^2}{A^2} = \lambda + \xi v_{;\ell}^\ell, \quad (7)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{C}}{C} + \frac{\dot{A}\dot{C}}{AC} - \frac{m^2}{A^2} = \xi v_{;\ell}^\ell, \quad (8)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{m^2}{A^2} = \xi v_{;\ell}^\ell, \quad (9)$$

$$\frac{\dot{A}\dot{B}}{AB} + \frac{\dot{A}\dot{C}}{AC} + \frac{\dot{B}\dot{C}}{BC} - \frac{3m^2}{A^2} = \rho, \quad (10)$$

$$2\frac{\dot{A}}{A} - \frac{\dot{B}}{B} - \frac{\dot{C}}{C} = 0, \quad (11)$$

where

$$\theta = v_{;\ell}^\ell = \frac{\dot{A}}{A} + \frac{\dot{B}}{B} + \frac{\dot{C}}{C}. \quad (12)$$

2.1. SOLUTION OF THE FIELD EQUATIONS

Integrating (11) and engrossing the constant of integration in B or C, without loss of generality, we obtain

$$A^2 = B C \quad (13)$$

Subtracting (8) from (9) and taking integral of the resulting equation two times, we get

$$\frac{B}{C} = c_2 \exp \left[c_1 \int \frac{dt}{ABC} \right], \quad (14)$$

where c_1 and c_2 are constants of integration.

Equations (7) - (11) are five independent equations in six unknown parameters A, B, C, λ , ξ and ρ . For the complete determination of the system, we need one extra condition. Following Yadav [33], we assume the law of variation of scale factor as increasing function of time as given by

$$a = (t^k e^t)^{\frac{1}{n}}, \quad (15)$$

where k and n are positive constants. If we put $n = 2$, Eq. (15) reduces to $a(t) = \sqrt{t^k e^t}$ which is used by Saha *et al.* [25] and Pradhan & Amirhashchi [20] in studying two-fluid scenario for dark energy models in an FRW universe and accelerating dark energy models in Bianchi type-V space-time respectively.

Now the spatial volume V of the model is given by

$$V = a^3 = ABC \quad (16)$$

Equations (13), (15) and (16) lead to

$$A(t) = (t^k e^t)^{\frac{1}{n}} \quad (17)$$

Inserting Equations (17) and (13) into Equation (14), we get

$$B(t) = \sqrt{c_2} (t^k e^t)^{\frac{1}{n}} \exp \left[\frac{c_1}{2} \int \frac{dt}{(t^k e^t)^{\frac{3}{n}}} \right], \quad (18)$$

$$C(t) = \frac{1}{\sqrt{c_2}} (t^k e^t)^{\frac{1}{n}} \exp \left[\frac{-c_1}{2} \int \frac{dt}{(t^k e^t)^{\frac{3}{n}}} \right]. \quad (19)$$

Hence the metric (1) reduces to the form

$$ds^2 = -dt^2 + (t^k e^t)^{\frac{2}{n}} \left[dx^2 + e^{2mx} \left(c_2 \exp \left(c_1 \int \frac{dt}{(t^k e^t)^{\frac{3}{n}}} \right) dy^2 + \frac{1}{c_2} \exp \left(-c_1 \int \frac{dt}{(t^k e^t)^{\frac{3}{n}}} \right) dz^2 \right) \right] \quad (20)$$

2.2 SOME GEOMETRICAL AND PHYSICAL PROPERTIES OF THE MODEL

The proper energy density (ρ), string tension density (λ) and bulk viscosity coefficient (ξ) are respectively given by

$$\rho = \frac{3}{n^2} \left(\frac{k}{t} + 1 \right)^2 - \frac{c_1^2}{4 (t^k e^t)^{\frac{6}{n}}} - \frac{3m^2}{(t^k e^t)^{\frac{2}{n}}}, \quad (21)$$

$$\lambda = 0, \quad (22)$$

$$\xi = \frac{1}{n} \left(\frac{k}{t} + 1 \right) + \frac{n}{3 \left(\frac{k}{t} + 1 \right)} \left[\frac{c_1^2}{4 (t^k e^t)^{\frac{6}{n}}} - \frac{m^2}{(t^k e^t)^{\frac{2}{n}}} - \frac{2k}{nt^2} \right]. \quad (23)$$

The physical parameters such as spatial volume (V), Hubble parameter (H), expansion scalar (θ), shear scalar (σ) and anisotropy parameters (A_m) are given by

$$V = (t^k e^t)^{\frac{3}{n}}, \quad (24)$$

$$H = \frac{1}{n} \left(\frac{k}{t} + 1 \right), \quad (25)$$

$$\theta = \frac{3}{n} \left(\frac{k}{t} + 1 \right), \quad (26)$$

$$\sigma^2 = \frac{c_1^2}{4 (t^k e^t)^{\frac{6}{n}}}, \quad (27)$$

$$A_m = \frac{1}{6} \frac{c_1^2 n^2}{\left(\frac{k}{t} + 1 \right)^2 (t^k e^t)^{\frac{6}{n}}}. \quad (28)$$

From the above results, it can be seen that the spatial volume is zero at $t=0$. This shows that the universe starts evolving with zero volume at $t=0$ and expands with cosmic time t . We also observed that at initial epoch the values of proper energy density (ρ), coefficient of bulk viscosity (ξ), Hubble factor (H), expansion parameter (A_m) are very high and these values gradually decrease with the evolution of time; that is ρ , ξ , H , θ , σ , and A_m tend to zero as $t \rightarrow \infty$. Thus our

model starts with big-bang at $t = 0$ and the expansion in the model decreases as time increases. Since $\lim_{t \rightarrow \infty} \frac{\sigma}{\theta} = 0$.

Hence the model isotropizes for large values of t . The model (20) has real physical singularity at $t=0$. Since in the presence of bulk viscosity, the string tension density $\lambda=0$. Thus the string cosmological model with bulk viscous fluid for Bianchi type-V metric does not exist.

3. STRING COSMOLOGICAL MODEL WITHOUT VISCOUS FLUID

In this section we obtain a string cosmological model without bulk viscous fluid. The field equation (6) with Equation (2) (without bulk viscous fluid), lead to the following system of equations:

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} - \frac{m^2}{A^2} = \lambda, \quad (29)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{C}}{C} + \frac{\dot{A}\dot{C}}{AC} - \frac{m^2}{A^2} = 0, \quad (30)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{m^2}{A^2} = 0, \quad (31)$$

$$\frac{\dot{A}\dot{B}}{AB} + \frac{\dot{A}\dot{C}}{AC} + \frac{\dot{B}\dot{C}}{BC} - \frac{3m^2}{A^2} = \rho, \quad (32)$$

$$\frac{2\dot{A}}{A} - \frac{\dot{B}}{B} - \frac{\dot{C}}{C} = 0. \quad (33)$$

3.1. SOLUTION OF THE FIELD EQUATIONS

From Equations (30) and (31), we obtain

$$\left(\frac{\ddot{B}C - B\ddot{C}}{\dot{B}C - B\dot{C}} \right) + \frac{\dot{A}}{A} = 0, \quad (34)$$

Using Equation (33) in Equation (34), we get

$$\left(\frac{\ddot{B}C - B\ddot{C}}{\dot{B}C - B\dot{C}} \right) + \frac{1}{2} \left(\frac{B\dot{C} + \dot{B}C}{BC} \right) = 0. \quad (35)$$

Equation (35), after integration, reduces to

$$\left(\dot{B}C - B\dot{C} \right) = \frac{L}{\sqrt{BC}}, \quad (36)$$

where L is constant of integration.

Using Equation (33) in Equation (30), we get

$$\frac{2\ddot{B}}{B} + \frac{6\ddot{C}}{C} - \frac{\dot{B}^2}{B^2} + \frac{\dot{C}^2}{C^2} + \frac{4\dot{B}\dot{C}}{BC} - \frac{4m^2}{BC} = 0. \quad (37)$$

$$\text{Let } BC = \alpha \text{ and } \frac{B}{C} = \beta. \quad (38)$$

Using these in Equations (36) and (37), we obtain

$$\frac{\dot{\beta}}{\beta} = \frac{L}{\alpha^{3/2}} \quad (39)$$

$$\text{and } \frac{4\ddot{\alpha}}{\alpha} - \frac{2\ddot{\beta}}{\beta} - \frac{\dot{\alpha}^2}{\alpha^2} + \frac{3\dot{\beta}^2}{\beta^2} - \frac{3\dot{\alpha}\dot{\beta}}{\alpha\beta} - \frac{4m^2}{\alpha} = 0. \quad (40)$$

Using Equation (39) in Equation (40), we get

$$4\alpha\ddot{\alpha} + \frac{L^2}{\alpha} - \dot{\alpha}^2 - 4\alpha m^2 = 0,$$

which implies that

$$\ddot{\alpha} - \frac{\dot{\alpha}^2}{4\alpha} = m^2 - \frac{L^2}{4\alpha^2}. \quad (41)$$

Let $\dot{\alpha} = f(\alpha)$ which implies that $\ddot{\alpha} = ff'$, where $f' = \frac{df}{d\alpha}$. Hence Equation (41) takes the form

$$\frac{d}{d\alpha}(f^2) - \frac{1}{2\alpha}(f^2) = 2m^2 - \frac{L}{2\alpha^2}, \quad (42)$$

which after integration, reduces to

$$f^2 = 4m^2\alpha + \frac{L^2}{3\alpha} + M\sqrt{\alpha}, \quad (43)$$

where M is an integrating constant.

Equations (39) and (43) lead to

$$\log \beta = \int \frac{L\alpha^{-3/2}d\alpha}{\sqrt{4m^2\alpha + \frac{L^2}{3\alpha} + M\sqrt{\alpha}}}. \quad (44)$$

Thus the metric (1) reduces to

$$ds^2 = - \frac{dT^2}{\left[4m^2T + \frac{L^2}{3T} + M\sqrt{T}\right]} + TdX^2 + \beta Te^{2mX}dY^2 + \frac{T}{\beta}e^{2mX}dZ^2, \quad (45)$$

where $x=X, y=Y, z=Z, \alpha=T$ and β can be determined by Equation (44).

3.2. SOME GEOMETRICAL AND PHYSICAL PROPERTIES OF THE MODEL

The energy density (ρ), the string tension (λ), the scalar of expansion (θ), the magnitude of shear (σ^2), proper volume (V) and Hubble's parameter (H) for the model (45) are given by

$$\rho = \frac{3}{4} \frac{M}{T^{3/2}} \quad (46)$$

$$\lambda = 0 \quad (47)$$

$$V = T^{3/2} \quad (48)$$

$$\sigma^2 = \frac{L^2}{4T^3} \quad (49)$$

$$\theta = \frac{3}{2} \sqrt{\frac{4m^2}{T} + \frac{L^2}{3T^3} + \frac{M}{T^{3/2}}} \quad (50)$$

$$H = \frac{1}{2} \sqrt{\frac{4m^2}{T} + \frac{L^2}{3T^3} + \frac{M}{T^{3/2}}} \quad (51)$$

From Equations (48) and (50), we observe that the spatial volume is zero at $T=0$ and the expansion scalar is infinite, which shows that the universe starts evolving with zero volume with infinite rate of expansion which is big-bang scenario. The physical quantities such as proper energy density (ρ), shear scalar (σ) and Hubble factor (H) diverge

at $T=0$. As $T \rightarrow \infty$, volume becomes infinite where as ρ, θ, σ, H approach to zero. Since $\lim_{T \rightarrow \infty} \frac{\sigma}{\theta} = 0$. Hence the

model approaches isotropy for large values of T . Since in the absence of bulk viscosity the string tension density $\lambda=0$. Hence the string cosmological model in the absence of bulk viscosity for Bianchi type-V metric does not exist.

4. CONCLUSION

We have presented two categories of Bianchi type-V cosmological solutions of the field equations, one with bulk viscous fluid and other without bulk viscous fluid. In both the categories of the cosmological models, we have shown that cosmic strings do not exist in Bianchi type-V cosmology. We have also discussed some physical and geometrical properties of the cosmological models.

REFERENCES

1. Aydogdu, O. and Salti M., The momentum 4-vector in bulk viscous Bianchi type-V space time, Czech. J. Phys. 56 (2006), 789- 798.
2. Banerjee, A., Sanyal, A.K. and Chakraborty, S., String Cosmology in Bianchi I space-time, Pramana J. Phys. 34 (1990), 1-11.
3. Banerjee, A. and Sanyal, A.K., Irrotational Bianchi-V viscous fluid cosmology with heat flux, Gen. Rel. Grav. 20 (1998), 103 -113.
4. Beesham, A., Tilted Bianchi type-V cosmological model in the scale-covariant theory, Astrophys. Space Sci. 125 (1986), 99 -102.
5. Chakraborty, S., String cosmology in Bianchi VI0 space-time, Ind. J. Pure Appl. Phys. 29 (1991), 31-33.
6. Camci, U., Yavuz, I., Baysal, H., Taran, I. and Yilmaz, I., Generation of Bianchi type-V universe filled with a perfect fluid, Astrophys. Space Sci. 275 (2000), 391-400.
7. Collins, C.B., Tilting at cosmological singularities, Comm. Math. Phys. 39 (1974), 131-151.
8. Deo, S.D., Roughe, A.K., Non existence of Bianchi type-I, V and VI0 cosmic string in Bimetric relativity, Ultra Scientist J. Phys. Sci. 22 (2010), 535 - 538.
9. Fransworth, D.L., Some new general relativistic dust metrics possessing isometries, J. Math. Phys. 8 (1967), 2315 - 2317.
10. Gott, J.R., Gravitational Lensing effects of vacuum strings : Exact Solutions, Ap. Phys. J. 288 (1985), 422 - 427.
11. Heller, M. and Klimek, Z., Viscous universes without initial singularity, Astrophys. Space Sci. 33 (1975), L37 - L39.
12. Krori, K.D., Chaudhary, T., Mahanta, C.R. and Mazumder, A., Some exact solutions in string cosmology, Gen. Rel. Grav. 22 (1990), 123 - 130.
13. Krori, K.D., Chaundhury, T. and Mahanta, C.R., Strings in some Bianchi type cosmologies, Gen. Rel. Grav. 26 (1994), 265- 274.
14. Landau, L.D. and Lifhsitz, E.M., Fluid Mech. 6, 505 (1963).
15. Letelier, P.S., Clouds of strings in general relativity, Phys. Rev. D 20 (1979), 1294 -1302.
16. Letelier, P.S., String cosmologies, Phys. Rev. D 28 (1983), 2414 - 2419.
17. Maartens, R. and Nel, S.D., Decomposable differential operators in a cosmological context, Comm. Math. Phys. 59 (1978), 273 - 283.
18. Nightingale, J.D., Independent investigations concerning bulk viscosity in relativistic homogenous isotropic cosmologies, Astrophys. J. 185 (1973), 105 - 114.
19. Pradhan, A., Yadav, M.K. and Rai, A., Some Bianchi type-III string cosmological models with bulk viscosity, Int. J. Theo. Phys. 46 (2007), 2677 - 2687.
20. Pradhan, A. and Amirhashchi, H., Accelerating dark energy models in Bianchi type-V space-time, Mod. Phys. Lett. A 26 (2011), 2266 - 2275.
21. Reddy, D.R.K., Non-Existence of Cosmic strings in bimetric theory of gravitation, Astrophys. space Sci. 286 (2003), 397 - 400.
22. Roy, S.R. and Singh, J.P., LRS Bianchi type-V universes filled with matter and radiation, Astrophys. Space Sci. 96 (1983), 303 - 312.
23. Rosen's, N., A bimetric theory of gravitation, Gen. Rel. Grav. 4 (1973), 435 - 447.
24. Saha, B., Anisotropic cosmological models with a perfect fluid and a Λ - term, Astrophys. space Sci. 302 (2006), 83 - 91.
25. Saha, B., Amirhashchi, H. and Pradhan, A., Two-Fluid Scenario for dark energy models in an FRW universe-revisited, arXiv:1108.2133 [gr-qc] (2011).
26. Santos, N.O., Dias, R.S. and Banerjee, A., Isotropic homogenous universe with viscous fluid, J. Math. Phys. 26, 878 (1975).
27. Singh, T. and Chaubey, R., Bianchi type-V universe with a viscous fluid and Λ - term, Pramana J. Phys. 68 (2007), 721 - 734.
28. Stachel, J., Thickening the string. I. The string perfect dust, Phys. Rev. D 21(1980), 2171- 2181.
29. Tikekar, R. and Patel, L.K., Some exact solution of string cosmology in Bianchi III space-time, Gen- Rel. Grav. 24 (1992), 397 - 404.
30. Vilenkin, A., Cosmic strings, Phys. Rev. D 24 (1981), 858- 868.
31. Wang, X.X., Exact solutions for String cosmology, Chin. Phys. Lett. 20, 615 (2003).
32. Weinberg, S., Entropy generation and the survival of proto galaxies in an expanding universe, Astrophys. J. 168 (1971), 175 - 194.
33. Yadav, A.K., Bianchi-V string cosmological model and late time acceleration, arXiv: 1202.3644v3 [Physics. gen-ph] (2012).

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2019. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]