

ON SEMI GENERALIZED TOPOLOGICAL SPACES

ARUN KUMAR GALI*
 Department of Mathematics,
 Government First Grade College,
 Navanagar Bagalkot, Karnataka, INDIA.

(Received On: 01-01-20; Revised & Accepted On: 13-02-20)

ABSTRACT

The aim of this paper is to introduce and study two new classes of spaces, namely Semi generalised normal and Semi generalized regular spaces and obtained their properties by utilizing Semi generalised closed sets.

Keywords: Semi generalised closed set, Semi generalised open sets, Semi generalised regular space and Semi generalised normal space.

Mathematics subject classification (2010): 54A05.

1 INTRODUCTION

Maheshwari and Prasad[8] introduced the new class of spaces called s-normal spaces using semi-open sets. It was further studied by Noiri and Popa[10], Dorsett[6] and Arya [1]. Munshi[9], introduced g-regular and g-normal spaces using g-closed sets of Levine[7]. Later, Benchalli *et al.* [3] and Shik John[12] studied the concept of g*-pre-regular, g*-pre normal and w-normal, w-regular spaces in topological spaces. Recently, Benchalli *et al.* [2,11] introduced and studied the properties of regular weakly closed sets and regular weakly continuous functions.

2. PRELIMINARIES

Throughout this paper space (X, τ) and (Y, σ) (or simply X and Y) always denote topological space on which no separation axioms are assumed unless explicitly stated. For a subset A of a space X , $Cl(A)$, $Int(A)$, A^c , and $\alpha\text{-}Cl(A)$, denote the Closure of A , Interior of A and Complement of A and α -closure of A in X respectively.

Definition 2.1: A subset A of a topological space (X, τ) is called

- (i) Generalized closed set (briefly g-closed) [7] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in X .
- (ii) W-closed set [12] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open in X .

Definition 2.2: A topological space X is said to be a

1. α -regular [4], if for each α -closed set F of X and each point $x \notin F$, there exists disjoint α -open sets U and V such that $F \subseteq U$ and $x \in V$.
2. w-regular [12], if for each closed set F of X and each point $x \notin F$, there exists disjoint w-open sets U and V such that $F \subseteq U$ and $x \in V$.
3. g-regular [10], if for each g-closed set F of X and each point $x \notin F$, there exists disjoint open sets U and V such that $F \subseteq U$ and $x \in V$.
4. Semi generalized closed set [15] $Scl(A) \subseteq U$ When $A \subseteq U$ and U is Semi-open in X

Definition 2.3: A topological space X is said to be a

1. α -normal [4], if for any pair of disjoint α -closed sets A and B , there exists disjoint α -open sets U and V such that $A \subseteq U$ and $B \subseteq V$.
2. w-normal [12], if for any pair of disjoint w-closed sets A and B , there exists disjoint open sets U and V such that $A \subseteq U$ and $B \subseteq V$.
3. g-normal [10], if for any pair of disjoint g-closed sets A and B , there exists disjoint open sets U and V such that $A \subseteq U$ and $B \subseteq V$.

Corresponding Author: Arun Kumar Gali*

Department of Mathematics, Government First Grade College, Navanagar Bagalkot, Karnataka, INDIA.

Definition 2.4: [2] A topological space X is called $T_{\text{regular weakly}}$ -space if every Semi generalised closed set is closed set.

Definition 2.5: A map $f: (X, \tau) \rightarrow (Y, \tau)$ is said to be

- (i) Semi generalised continuous map [16] if $f^{-1}(V)$ is a Semi generalised closed set of (X, τ) for every closed set V of (Y, τ) .
- (ii) Semi generalised irresolute map [16] if $f^{-1}(V)$ is a Semi generalised closed set of (X, τ) for every Semi generalised closed set V of (Y, τ) .

3. SEMI GENERALIZED REGULAR SPACE

In this section, we introduce a new class of space called Semi generalised regular space using Semi generalised closed set and obtain some of their characterizations.

Definition 3.1. A topological space X is said to be Semi generalised regular space if for each Semi generalised closed set F and a point $x \notin F$, there exist disjoint open sets G and H such that $F \subseteq G$ and $x \in H$.

We have the following interrelationship between Semi generalised regularity and regularity.

Theorem 3.2: Every Semi generalised regular space is regular.

Proof: Let X be a Semi generalised regular space. Let F be any closed set in X and a point $x \notin F$. By [2], F is Semi generalised topological space-closed and $x \notin F$. Since X is a Semi generalised regular space, there exists a pair of disjoint open sets G and H such that $F \subseteq G$ and $x \in H$. Hence X is a regular space.

Remark 3.3: If X is a regular space and $T_{\text{Semi generalised topological space}}$, then X is Semi generalised regular space then we have the following characterization.

Theorem 3.4: The following statements are equivalent for a topological space X .

- (i) X is a Semi generalised regular space
- (ii) For each $x \in X$ and each Semi generalised topological space open neighbourhood U of x , there exists an open neighbourhood N of x such that $\text{cl}(N) \subseteq U$.

Proof: (i) implies (ii): Suppose X is a Semi generalised regular space. Let U be any Semi generalised neighbourhood of x . Then there exists Semi generalised open set G such that $x \in G \subseteq U$. Now $X - G$ is Semi generalised closed set and $x \notin X - G$. Since X is Semi generalised regular space, then there exist open sets M and N such that $X - G \subseteq M$, $x \in N$ and $M \cap N = \emptyset$ and so $N \subseteq X - M$. Now $\text{cl}(N) \subseteq \text{cl}(X - M) = X - M$ and $X - M \subseteq M$. This implies $X - M \subseteq U$. Therefore $\text{cl}(N) \subseteq U$.

(ii) implies (i): Let F be any Semi generalised topological space closed set in X and $x \in X - F$ and $X - F$ is a Semi generalised topological space open and so $X - F$ is a Semi generalised topological space neighbourhood of x . By hypothesis, there exists an open neighbourhood N of x such that $x \in N$ and $\text{cl}(N) \subseteq X - F$. This implies $F \subseteq X - \text{cl}(N)$ is an open set containing F and $N \cap (X - \text{cl}(N)) = \emptyset$. Hence X is Semi generalised regular space.

We have another characterization of Semi generalised regularity in the following.

Theorem 3.5: A topological space X is Semi generalised regular if and only if for each Semi generalised topological space closed set F of X and each $x \in X - F$ there exist open sets G and H of X such that $x \in G$, $F \subseteq H$ and $\text{cl}(G) \cap \text{cl}(H) = \emptyset$.

Proof: Suppose X is Semi generalised regular space. Let F be a Semi generalised topological space closed set in X with $x \notin F$. Then there exist open sets M and H of X such that $x \in M$, $F \subseteq H$ and $M \cap H = \emptyset$. This implies $M \cap \text{cl}(H) = \emptyset$. As X is Semi generalised regular, there exist open sets U and V such that $x \in U$, $\text{cl}(H) \subseteq V$ and $U \cap V = \emptyset$. so $\text{cl}(U) \cap V = \emptyset$.

Let $G = M \cap U$, then G and H are open sets of X such that $x \in G$, $F \subseteq H$ and $\text{cl}(G) \cap \text{cl}(H) = \emptyset$.

Conversely, if for each Semi generalised closed set F of X and each $x \in X - F$ there exist open sets G and H such that $x \in G$, $F \subseteq H$ and $\text{cl}(G) \cap \text{cl}(H) = \emptyset$. This implies $x \in G$, $F \subseteq H$ and $G \cap H = \emptyset$. Hence X is Semi generalised regular.

Now we prove that Semi generalised topological spaces- regularity is a hereditary property.

Theorem 3.6: Every subspace of a Semi generalised regular space is Semi generalised regular.

Proof: Let X be a Semi generalised regular space. Let Y be a subspace of X . Let $x \in Y$ and F be a Semi generalised closed set in Y such that $x \notin F$. Then there is a closed set and so Semi generalised closed set A of X with $F = Y \cap A$ and $x \notin A$. Therefore we have $x \in X$, A is Semi generalised closed in X such that $x \notin A$. Since X is Semi generalised regular, then there exist open sets G and H such that $x \in G$, $A \subseteq H$ and $G \cap H = \emptyset$. Note that $Y \cap G$ and $Y \cap H$ are open sets in Y . Also $x \in G$ and $x \in Y$, which implies $x \in Y \cap G$ and $A \subseteq H$ implies $Y \cap G \subseteq Y \cap H$, $F \subseteq Y \cap H$. Also $(Y \cap G) \cap (Y \cap H) = \emptyset$. Hence Y is Semi generalised regular space.

We have yet another characterization of Semi generalised topological spaces-regularity in the following.

Theorem 3.7: The following statements about a topological space X are equivalent:

- (i) X is Semi generalised regular
- (ii) For each $x \in X$ and each Semi generalised topological space open set U in X such that $x \in U$ there exists an open set V in X such that $x \in V \subseteq \text{cl}(V) \subseteq U$.
- (iii) For each point $x \in X$ and for each Semi generalised topological space closed set A with $x \notin A$, then there exists an open set V containing x such that $\text{cl}(V) \cap A = \emptyset$.

Proof:

- (i) implies(ii): Follows from Theorem 3.5.
- (ii) implies(iii): Suppose (ii) holds. Let $x \in X$ and A be an Semi generalised topological space closed set of X such that $x \notin A$. Then $X - A$ is a Semi generalised topological space open set with $x \in X - A$. By hypothesis, there exists an open set V such that $x \in V \subseteq \text{cl}(V) \subseteq X - A$. That is $x \in V$, $V \subseteq \text{cl}(V)$ and $\text{cl}(V) \subseteq X - A$. So $x \in V$ and $\text{cl}(V) \cap A = \emptyset$.
- (iii) implies(i): Let $x \in X$ and U be an Semi generalised topological space open set in X such that $x \in U$. Then $X - U$ is an Semi generalised topological space closed set and $x \notin X - U$. Then by hypothesis, there exists an open set V containing x such that $\text{cl}(V) \cap (X - U) = \emptyset$. Therefore $x \in V$, $\text{cl}(V) \subseteq U$ so $x \in V \subseteq \text{cl}(V) \subseteq U$.

The invariance of Semi generalised topological space regularity is given in the following.

Theorem 3.8: Let $f : X \rightarrow Y$ be a bijective, Semi generalised topological space irresolute and open map from a Semi generalised topological space regular space X into a topological space Y , then Y is Semi generalised topological spaces-regular.

Proof: Let $y \in Y$ and F be a Semi generalised topological space closed set in Y with $y \notin F$. Since F is Semi generalised topological space irresolute, $f^{-1}(F)$ is Semi generalised topological space closed set in X . Let $f(x) = y$ so that $x = f^{-1}(y)$ and $x \notin f^{-1}(F)$. Again X is Semi generalised-regular space, then there exist open sets U and V such that $x \in U$ and $f^{-1}(F) \subseteq V$, $U \cap V = \emptyset$. Since f is open and bijective, we have $y \in f(U)$, $F \subseteq f(V)$ and $f(U) \cap f(V) = f(U \cap V) = f(\emptyset) = \emptyset$. Hence Y is Semi generalised regular space.

Theorem 3.9: Let $f : X \rightarrow Y$ be a bijective, Semi generalised closed and open map from a topological space X into a Semi generalised regular space Y . If X is $T_{\text{Semi generalised topological spaces}}$, then X is Semi generalised regular.

Proof: Let $x \in X$ and F be an Semi generalised closed set in X with $x \notin F$. Since X is $T_{\text{Semi generalised topological spaces}}$, F is closed in X . Then $f(F)$ is Semi generalised closed set with $f(x) \notin f(F)$ in Y , since f is Semi generalised closed. As Y is Semi generalised regular, then there exist open sets U and V such that $x \in U$ and $f(x) \in U$ and $f(F) \subseteq V$. Therefore $x \in f^{-1}(U)$ and $F \subseteq f^{-1}(V)$. Hence X is Semi generalised regular space.

Theorem 3.10: If $f : X \rightarrow Y$ is w -irresolute, continuous injection and Y is Semi generalised topological spaces-regular space, then X is Semi generalised topological spaces-regular.

Proof: Let F be any closed set in X with $x \notin F$. Since f is w -irresolute, $f(F)$ is Semi generalised topological space closed set in Y and $f(x) \notin f(F)$. Since Y is Semi generalised regular, then there exists open sets U and V such that $f(x) \in U$ and $f(F) \subseteq V$. Thus $x \in f^{-1}(U)$, $F \subseteq f^{-1}(V)$ and $f^{-1}(U) \cap f^{-1}(V) = \emptyset$. Hence X is Semi generalised regular space.

4. SEMI GENERALISED NORMAL SPACES

In this section, we introduce the concept of Semi generalised normal spaces and study some of their characterizations.

Definition 4.1: A topological space X is said to be Semi generalised normal if for each pair of disjoint Semi generalised topological spaces closed sets A and B in X , then there exists a pair of disjoint open sets U and V in X such that $A \subseteq U$ and $B \subseteq V$.

We have the following interrelationship.

Theorem 4.2: Every Semi generalised normal space is normal.

Proof: Let X be a Semi generalised normal space. Let A and B be a pair of disjoint closed sets in X . From [2], A and B are Semi generalised topological spaces closed sets in X . Since X is Semi generalised normal, then there exists a pair of disjoint open sets G and H in X such that $A \subseteq G$ and $B \subseteq H$. Hence X is normal.

Remark 4.3: The converse need not be true in general as seen from the following example.

Example 4.4: Let $X = Y = \{a, b, c, d\}$, $\tau = \{X, \emptyset, \{a\}, \{c\}, \{a, c\}, \{b, c, d\}\}$. Then the space X is normal but not Semi generalised normal, since the pair of disjoint Semi generalised topological spaces closed sets namely, $A = \{a, d\}$ and $B = \{b, c\}$ for which there do not exist disjoint open sets G and H such that $A \subseteq G$ and $B \subseteq H$.

Remark 4.5: If X is normal and $T_{\text{Semi generalised topological spaces}}$, then X is Semi generalised -normal.

Hereditary property of Semi generalised normality is given in the following.

Theorem 4.6: A Semi generalised closed subspace of a Semi generalised normal space is Semi generalised normal. We have the following characterization.

Theorem 4.7: The following statements for a topological space X are equivalent:

- (i) X is Semi generalised topological spaces normal
- (ii) For each Semi generalised closed set A and each pre generalized pre regular weakly topological space open set U such that $A \subseteq U$, there exists an open set V such that $A \subseteq V \subseteq \text{cl}(V) \subseteq U$
- (iii) For any Semi generalised closed sets A, B , there exists an open set V such that $A \subseteq V$ and $\text{cl}(V) \cap B = \emptyset$.
- (iv) For each pair A, B of disjoint Semi generalised closed sets then there exist open sets U and V such that $A \subseteq U, B \subseteq V$ and $\text{cl}(U) \cap \text{cl}(V) = \emptyset$.

Proof:

- (i) implies (ii): Let A be a Semi generalised closed set and U be a Semi generalised open set such that $A \subseteq U$. Then A and $X - U$ are disjoint Semi generalised closed sets in X . Since X is Semi generalised normal, then there exists a pair of disjoint open sets V and W in X such that $A \subseteq V$ and $X - U \subseteq W$. Now $X - W \subseteq X - (X - U)$, so $X - W \subseteq U$ also $V \cap W = \emptyset$ implies $V \subseteq X - W$, so $\text{cl}(V) \subseteq \text{cl}(X - W)$ which implies $\text{cl}(V) \subseteq X - W$. Therefore $\text{cl}(V) \subseteq X - W \subseteq U$. So $\text{cl}(V) \subseteq U$. Hence $A \subseteq V \subseteq \text{cl}(V) \subseteq U$.
- (ii) implies (iii): Let A and B be a pair of disjoint Semi generalised closed sets in X . Now $A \cap B = \emptyset$, so $A \subseteq X - B$, where A is Semi generalised closed and $X - B$ is Semi generalised open. Then by (ii) there exists an open set V such that $A \subseteq V \subseteq \text{cl}(V) \subseteq X - B$. Now $\text{cl}(V) \subseteq X - B$ implies $\text{cl}(V) \cap B = \emptyset$. Thus $A \subseteq V$ and $\text{cl}(V) \cap B = \emptyset$.
- (iii) implies (iv): Let A and B be a pair of disjoint Semi generalised closed sets in X . Then from (iii) there exists an open set U such that $A \subseteq U$ and $\text{cl}(U) \cap B = \emptyset$. Since $\text{cl}(V)$ is closed, so Semi generalised closed set. Therefore $\text{cl}(V)$ and B are disjoint Semi generalised closed sets in X . By hypothesis, then there exists an open set V , such that $B \subseteq V$ and $\text{cl}(U) \cap \text{cl}(V) = \emptyset$.
- (iv) implies (i): Let A and B be a pair of disjoint Semi generalised closed sets in X . Then from (iv) then there exist an open sets U and V in X such that $A \subseteq U, B \subseteq V$ and $\text{cl}(U) \cap \text{cl}(V) = \emptyset$. So $A \subseteq U, B \subseteq V$ and $U \cap V = \emptyset$. Hence X is Semi generalised normal.

Theorem 4.8: Let X be a topological space. Then X is Semi generalised normal if and only if for any pair A, B of disjoint Semi generalised closed sets then there exist open sets U and V of X such that $A \subseteq U, B \subseteq V$ and $\text{cl}(U) \cap \text{cl}(V) = \emptyset$.

Theorem 4.9: Let X be a topological space. Then the following are equivalent:

- (i) X is normal
- (ii) For any disjoint closed sets A and B , then there exist disjoint Semi generalised topological spaces- open sets U and V such that $A \subseteq U, B \subseteq V$.
- (iii) For any closed set A and any open set V such that $A \subseteq V$, there exists a Semi generalised open set U of X such that $A \subseteq U \subseteq \alpha\text{cl}(U) \subseteq V$.

Proof:

- (i) implies (ii): Suppose X is normal. Since every open set is Semi generalised open [2], (ii) follows.
- (ii) implies (iii): Suppose (ii) holds. Let A be a closed set and V be an open set containing A . Then A and $X - V$ are disjoint closed sets. By (ii), then there exist disjoint Semi generalised open sets U and W such that $A \subseteq U$ and $X - V \subseteq W$, since $X - V$ is closed, so Semi generalised is closed. From [2], we have $X - V \subseteq \alpha\text{-int}(W)$ and $U \cap \alpha\text{-int}(W) = \emptyset$ and so we have $\alpha\text{-cl}(U) \cap \alpha\text{-int}(W) = \emptyset$. Hence $A \subseteq U \subseteq \alpha\text{-cl}(U) \subseteq X - \alpha\text{-int}(W) \subseteq V$. Thus $A \subseteq U \subseteq \alpha\text{-cl}(U) \subseteq V$.
- (iii) implies (i): Let A and B be a pair of disjoint closed sets of X . Then $A \subseteq X - B$ and $X - B$ is open. There exists a Semi generalised open set G of X such that $A \subseteq G \subseteq \alpha\text{-cl}(G) \subseteq X - B$. Since A is closed, it is w -closed, we have $A \subseteq \alpha\text{-int}(G)$. Take $U = \text{int}(\text{cl}(\text{int}(\alpha\text{-int}(G))))$ and $V = \text{int}(\text{cl}(\text{int}(X - \alpha\text{-cl}(G))))$. Then U and V are disjoint open sets of X such that $A \subseteq U$ and $B \subseteq V$. Hence X is normal.

We have the following characterization of Semi generalised topological spaces- normality and Semi generalised topological spaces- normality.

Theorem 4.10: Let X be a topological space. Then the following are equivalent:

- (i) X is α -normal.
- (ii) For any disjoint closed sets A and B , there exist disjoint Semi generalised topological space- open sets U and V such that $A \subseteq U, B \subseteq V$ and $U \cap V = \varnothing$.

Proof:

- (i) implies(ii): Suppose X is α - normal. Let A and B be a pair of disjoint closed sets of X . Since X is α -normal, there exist disjoint α – open sets U and V such that $A \subseteq U$ and $B \subseteq V$ and $U \cap V = \varnothing$.
- (ii) implies(i): Let A and B be a pair of disjoint closed sets of X . The by hypothesis there exist disjoint Semi generalised open sets U and V such that $A \subseteq U$ and $B \subseteq V$ and $U \cap V = \varnothing$. Since from [2], $A \subseteq \alpha\text{-int}U$ and $B \subseteq \alpha\text{-int}(V)$ and $\alpha\text{-int}U \cap \alpha\text{-int}V = \varnothing$. Hence X is α -normal.

Theorem 4.11: Let X be a α - normal, then the following hold good:

- (i) For each closed set A and every Semi generalised open set B such that $A \subseteq B$ there exists a α open set U such that $A \subseteq U \subseteq \alpha\text{-cl}(U) \subseteq B$.
- (ii) For every Semi generalised closed set A and every open set B containing A , there exist a α -open set U such that $A \subseteq U \subseteq \alpha\text{-cl}(U) \subseteq B$.

REFERENCES

1. S.P. Arya and T.M. Nour, Characterization of s - normal spaces, Indian. J.Pure and Appl. Math., 21(8),(1990), 717-719.
2. S.S.Benchalli and R.S Wali, on α - Closed sets in Topological Spaces, Bull. Malays. Math. sci. soc 30 (2007), 99-110.
3. S.S. Benchalli, T.D. Rayanagoudar and P.G. Patil, g^* - Pre Regular and g^* -Pre Normal Spaces, Int. Math. Forum 4/48(2010) 2399-2408.
4. S.S. Benchalli and P.G. Patil, Some New Continuous Maps in Topological Spaces, Journal of Advanced Studies in Topology 2/1-2 (2009) 53-63.
5. R. Devi, Studies on Generalizations of Closed Maps and Homeomorphisms in Topological Spaces, Ph.D. thesis, Bharatiyar University, Coimbatore (1994).
6. C. Dorsett, Semi normal Spaces, Kyungpook Math. J. 25 (1985) 173-180.
7. N. Levine, Generalized Closed sets in Topology, Rend. Circ. Math. Palermo 19/2(1970) 89-96.
8. S.N. Maheshwar and R. Prasad, On s -normal spaces, Bull. Math. Soc. Sci.Math. R.S. Roumanie 22 (1978) 27-28.
9. B.M. Munshi, Separation axioms, Acta Ciencia Indica 12 (1986) 140-146.
10. T. Noiri and V. Popa, On g -regular spaces and some functions, Mem. Fac. Sci. Kochi Univ. Math 20 (1999) 67-74. Journal of New Results in Science 5 (2014) 96-103 103
11. R.S.Wali, on some topics in general and fuzzy topological spaces Ph.d thesis Karnatak university Dharwad(2007).
12. M.Sheik. John, A Study on Generalizations of Closed Sets and Continuous Maps in Topological and Bitopological spaces, Ph.D Thesis, Bharathiar University, Coimbatore (2002).
13. R.S.Wali and Vivekananda Dembre; On pre generalised pre Closed Sets in Topological Spaces ;Journal of Computer and Mathematical Sciences, Vol.6(2),113-125, February 2015.
14. R.S.Wali and Vivekananda Dembre, Minimal weakly open sets and maximal weakly closed sets in topological spaces ; International Journal of Mathematical Archive; Vol-4(9)-Sept-2014.
15. P.Bhattacharya and B.K.Lahiri, Semi-generalized closed sets in topology, Indian J. Math., 29, 376-382 (1987).
16. P.Sundram, Ph.D Thesis Bharathiar University, Coimbatore-1991

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2020. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]