

THE CONCEPT OF $g^*\beta$ – CLOSED SETS IN TOPOLOGICAL SPACES

A. PUNITHA THARANI¹ AND SUJITHA.H^{*2}

¹Associate Professor, Department of Mathematics,
 St. Mary's College (Autonomous), Thoothukudi - 628001, TamilNadu, India.
 Affiliated to Manonmaniam Sundaranar University,
 Abishekapatti, Tirunelveli - 627012, TamilNadu, India.

²Research Scholar (Register number: 19122212092003), Department of Mathematics,
 St. Mary's College (Autonomous), Thoothukudi-628001, TamilNadu, India.
 Affiliated to Manonmaniam Sundaranar University,
 Abishekapatti, Tirunelveli - 627012, TamilNadu, India.

(Received On: 10-02-20; Revised & Accepted On: 01-03-20)

ABSTRACT

The aim of this paper is to introduce and study new class of sets called $g^*\beta$ – closed sets. This new class of sets lies between closed sets and βg -closed sets. Applying these sets, new spaces namely, ${}_{\beta}T^{**}_{1/2}$ -space, ${}_{\alpha\beta}T^*_c$ – space, ${}_{\beta}T^*_{1/2}$ – space, ${}_{\beta}T^*_{1/2}$ – space, ${}_{\beta}T^*_c$ – spaces are introduced.

Key words: $g^*\beta$ – irresolute map, ${}_{\beta}T^{**}_{1/2}$ -space, ${}_{\alpha\beta}T^*_c$ – space.

INTRODUCTION

The Kuratowski closure axioms or axioms for the closed sets are used to define every topological spaces. Hence we can understand how the important the concept of closed sets in the topological Spaces. In 1970, Levine [11] initiated the study of g – closed sets. Maki. *et.al* [14] defined αg – closed sets and $\alpha^{**}g$ – closed sets in 1994. S.P. Arya and T. Nour [3] defined gs – closed sets in 1990. Dontchev [9], Gnanambal [10] and Palaniappan and Rao [19] introduced gsp – closed sets, gpr – closed sets, and rg - closed sets respectively. M.K.R.S. Veerakumar [20] introduced g^* - closed sets in 1991. P.M. Helen [22] introduced g^{**} - closed sets. We introduce a new class of sets called $g^*\beta$ -closed sets which is properly placed in between the class of closed sets and the class of βg - closed sets. Levine [11] Devi. *et.al* [6,8] introduced $T_{1/2}$ - spaces, T_b - spaces and ${}_aT_b$ spaces respectively. The purpose of this paper is to introduce the concepts of, ${}_{\beta}T^{**}_{1/2}$ -space, ${}_{\alpha\beta}T^*_c$ – space, ${}_{\beta}T^*_{1/2}$ – space are introduced and investigated.

PRELIMINARIES

Throughout this paper (X, τ) , (Y, σ) and (Z, η) represent non-empty topological spaces of which no separation axioms are assumed unless otherwise mentioned. For a subset A of a space (X, τ) , $cl(A)$ and $int(A)$ denote the closure and the interior of A respectively. The class of all closed subsets of a space (X, τ) is denoted by $C(X, \tau)$. The smallest semi closed (resp. pre-closed and α -closed) set containing a subset of a space (X, τ) is called the semi-closure (resp. pre-closure and α -closure) of A is denoted by $scl(A)$ (resp. $pcl(A)$ and $\alpha cl(A)$).

Definition 2.1: A subset A of a topological space (X, τ) is called

- 1) a pre-open set [16] if $A \subseteq int(cl(A))$ and a preclosed set if $cl(int(A)) \subseteq A$.
- 2) a semi-open set [12] if $A \subseteq cl(int(A))$ and semi-closed set if $int(cl(A)) \subseteq A$.
- 3) a semi-preopen set [1] if $A \subseteq cl(int(cl(A)))$ and a semi preclosed set [1] if $int(cl(int(A))) \subseteq A$.
- 4) an α -open set [18] if $A \subseteq int(cl(int(A)))$ and an α -closed set [18] if $cl(int(cl(A))) \subseteq A$.
- 5) a regular-open set [16] if $int(cl(A))=A$ and regular-closed set [16] if $A=int(cl(A))$.

Corresponding Author: Sujitha.H^{*2}

²Research Scholar, Department of Mathematics,
 St. Mary's College (Autonomous), Thoothukudi - 628001, TamilNadu, India.

Definition 2.2: A subset A of a topological space (X, τ) is called

- 1) a generalised closed set (briefly g - closed) [11] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 2) g^* - closed if [20] $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is g^* -open in (X, τ) .
- 3) a generalised semi-closed set (briefly gs - closed) [3] is $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 4) an generalised semi pre-closed set (briefly gsp-closed) [9] if $spcl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 5) regular generalised closed set (briefly rg - closed) [19] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and regular open in (X, τ) .
- 6) α - generalised closed set (briefly αg - closed) [14] if $\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is g^* -open in (X, τ) .
- 7) g^{**} -closed [21] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 8) generalised pre regular-closed set (briefly gpr - closed)[10] if $pcl(A) \subseteq U$ whenever $A \subseteq U$ and U is regular open in (X, τ) .
- 9) weakly generalised closed set [18] (briefly wg - closed) if $cl(int(A)) \subseteq U$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 10) generalised pre-closed set (briefly gp - closed) [13] if $pcl(A)$ whenever $A \subseteq U$ and U is open in (X, τ) .
- 11) generalised α - closed (briefly $g\alpha$ - closed) [14] if $\alpha cl(A)$ whenever $A \subseteq U$ and U is α -open in (X, τ) .
- 12) Semi-generalised closed (briefly sg-closed) [5] if $scl(A)$ whenever $A \subseteq U$ and U is semiopen in (X, τ) .

Definition 2.3: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called

- 1) g- continuous [4] if $f^{-1}(V)$ is a g- closed set of (X, τ) for every closed set V of (Y, σ) .
- 2) αg - continuous [10] if $f^{-1}(V)$ is a αg - closed set of (X, τ) for every closed set V of (Y, σ) .
- 3) gs- continuous [7] if $f^{-1}(V)$ is a gs- closed set of (X, τ) for every closed set V of (Y, σ) .
- 4) gsp- continuous [9] if $f^{-1}(V)$ is a gsp- closed set of (X, τ) for every closed set V of (Y, σ) .
- 5) rg- continuous [19] if $f^{-1}(V)$ is a rg- closed set of (X, τ) for every closed set V of (Y, σ) .
- 6) gp- continuous [2] if $f^{-1}(V)$ is a gp- closed set of (X, τ) for every closed set V of (Y, σ) .
- 7) gpr- continuous [10] if $f^{-1}(V)$ is a gpr- closed set of (X, τ) for every closed set V of (Y, σ) .
- 8) g^* - continuous [20] if $f^{-1}(V)$ is a g^* - closed set of (X, τ) for every closed set V of (Y, σ) .
- 9) g^* -irresolute [20] if $f^{-1}(V)$ is a g^* - closed set of (X, τ) for every g^* -closed set of (Y, σ) .
- 10) wg- continuous [18] if $f^{-1}(V)$ is a wg- closed set of (X, τ) for every closed set V of (Y, σ) .
- 11) g^{**} - continuous [22] if $f^{-1}(V)$ is a g^{**} - closed set of (X, τ) for every closed set of (Y, σ) .
- 12) g^{**} -irresolute [22] if $f^{-1}(V)$ is a g^{**} - closed set of (X, τ) for every g^* -closed set V of (Y, σ) .

Definition 2.4: A topological space (X, τ) is said to be

- 1) a $T_{1/2}$ space [11] if every g- closed set in it is closed.
- 2) a T_b space [6] if every gs- closed set in it is closed.
- 3) a T_d space [6] if every gs- closed set in it is g- closed.
- 4) a $_aT_d$ space [4] if every αg - closed set in it is g- closed.
- 5) a $_aT_b$ space [8] if every αg - closed set in it is closed.
- 6) a $^*T_{1/2}$ space [20] if every g- closed set in it is g^* - closed set.
- 7) a $T^*_{1/2}$ space [20] if every g^* - closed set in it is closed.

3. BASIC PROPERTIES OF $g^*\beta$ -CLOSED SETS

We introduce the following definition

Definition 3.1: A subset A of (X, τ) is said to be $g^*\beta$ closed set if $\beta cl(A) \subseteq U$ whenever $A \subseteq U$ and U is g^* open in X. The family of all $g^*\beta$ - closed sets are denoted by $G^*\beta-C(X)$.

Proposition 3.2: Every closed set is $g^*\beta$ - closed.

Proof follows from the definition.

Proposition 3.3: Every β -closed set is $g^*\beta$ - closed set.

Proof follows from the definition.

Proposition 3.4: Every g^{**} -closed set is $g^*\beta$ - closed set.

Proof follows from the definition.

Proposition 3.5: Every g^* -closed set is $g^*\beta$ - closed set.

Proof follows from the definition.

Proposition 3.6: Every g-closed set is $g^*\beta$ - closed set.

Proof follows from the definition.

The converse of the above propositions need not be true in general.

Example 3.7: Let $X = \{a, b, c, d\}$. $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$.

Let $A = \{c\}$ is a $g^*\beta$ -closed set but not a closed set and g^{**} -closed set. So the class of a $g^*\beta$ -closed sets properly contains the class of closed sets and the class of g^{**} -closed sets. Also $\{c\}$ is not a g -closed set.

Example 3.8: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$. Let $A = \{a, b\}$ is a $g^*\beta$ -closed set but not a β -closed set and g^* - closed set of (X, τ) . So the class of $g^*\beta$ -closed sets properly contains the class of β -closed sets and the class of g^* -closed sets.

Proposition 3.9: Every $g^*\beta$ -closed set is (1) rg -closed (2) gp -closed (3) gpr - closed (4) gsp -closed (5) wg -closed. Proof follows from the definition.

The converse of the above propositions need not be true in general as seen in the following examples.

Example 3.10: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, c\}\}$. Let $A = \{a\}$ is gpr -closed set and a rg -closed set but not $g^*\beta$ -closed set.

Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, c\}\}$. Let $A = \{c\}$ is a gsp -closed, wg -closed and a gp -closed set but not a $g^*\beta$ -closed set of (X, τ) . Therefore the class of $g^*\beta$ -closed set is properly contained in the class of gpr -closed, rg -closed, gsp -closed, gp -closed and wg -closed.

Remark 3.11: $g^*\beta$ -closedness is independent of pre-closedness, semi pre-closedness, semiclosedness, $g\beta$ -closedness, β -closedness and sg -closedness. Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, c\}\}$. Let $A = \{a, b\}$ then A is $g^*\beta$ -closed set. A is neither β -closed nor semi-closed, infact it is not even a semipreclosed. Also it is not a sg -closed and $g\beta$ -closed.

Proposition 3.12: If A and B are $g^*\beta$ -closed sets, then $A \cup B$ is also a $g^*\beta$ -closed set. Proof follows from the fact that $\beta cl(A \cup B) = \beta cl(A) \cup \beta cl(B)$

Proposition 3.13: If A is both g^* -open and $g^*\beta$ -closed then A is β -closed. Proof follows from the definition of $g^*\beta$ -closed sets.

Proposition 3.14: A is $g^*\beta$ -closed set of (X, τ) if $\beta cl(A) \setminus A$ does not contain any non-empty g^* -closed set.

Proof: Let F be a g^* -closed set of (X, τ) such that $F \subseteq \beta cl(A) \setminus A$. Then $A \subseteq X \setminus F$. Since A is $g^*\beta$ -closed and $X \setminus F$ is g^* -open, $\beta cl(A) \subseteq X \setminus F$. This implies $F \subseteq X \setminus \beta cl(A)$. So $F \subseteq (X \setminus \beta cl(A)) \cap (\beta cl(A) \setminus A) \subseteq (X \setminus \beta cl(A)) \cap (\beta cl(A)) = \emptyset$. Hence $F = \emptyset$.

Proposition 3.15: If A is $g^*\beta$ -closed set of (X, τ) such that $A \subseteq B \subseteq \beta cl(A)$ then B is also a $g^*\beta$ -closed set of (X, τ) .

Proof: Let U be a g^* -open set of (X, τ) such that $B \subseteq U$. Then $A \subseteq U$. Since A is $g^*\beta$ -closed, then $\beta cl(A) \subseteq U$. Now $\beta cl(B) \subseteq \beta cl(\beta cl(A)) = \beta cl(A) \subseteq U$. Therefore B is also a $g^*\beta$ -closed set of (X, τ) .

Proposition 3.16: If A and B are two $g^*\beta$ -closed sets in a topological space X such that either $A \subset B$ or $B \subset A$ then both intersection and union of two $g^*\beta$ -closed set is $g^*\beta$ -closed set.

Proof: If A is contained in B or B is contained in A then $A \cup B = B$ or $A \cup B = A$ respectively. This shows that $A \cup B$ is $g^*\beta$ -closed as A and B are $g^*\beta$ -closed sets.

Similarly, $A \cap B$ is also a $g^*\beta$ -closed set.

Remark 3.17: Difference of two $g^*\beta$ -closed sets is not a $g^*\beta$ -closed set.

Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$. Here $A = \{a, b, d\}$ and $B = \{b, d\}$ are $g^*\beta$ -closed sets but $A - B = \{a\}$ is not.

Proposition 3.18: Let (X, τ) be a topological space then for each $x \in X$, the set $X \setminus \{x\}$ is $g^*\beta$ -closed or g^* -open.

Proof: If $X \setminus \{x\}$ is $g^*\beta$ -closed or g^* -open then we are done. Now suppose $X \setminus \{x\}$ is not g^* -open then X is the only g^* -open set containing $X \setminus \{x\}$ and also $\beta cl(X \setminus \{x\})$ is contained in X , as it is the biggest set containing all its subsets. Hence $X \setminus \{x\}$ is a $g^*\beta$ -closed in X .

4. $g^*\beta$ – CONTINUOUS AND $g^*\beta$ – IRRESOLUTE MAPS

We introduce the following definitions

Definition 4.1: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $g^*\beta$ – continuous if $f^{-1}(V)$ is $g^*\beta$ -closed set of (X, τ) for every closed set of (Y, σ) .

Theorem 4.2: Every continuous map is $g^*\beta$ – continuous.

Proof: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be continuous and let F be any closed set of Y , then $f^{-1}(F)$ is closed in X . Since every closed set is $g^*\beta$ -closed set, $f^{-1}(F)$ is $g^*\beta$ -closed set. Therefore f is $g^*\beta$ – continuous.

The following example supports that the converse of the above theorem need not be true in general.

Example 4.3: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$, $\sigma = \{\emptyset, Y, \{a, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ is defined as the identity map. The inverse image of all the closed sets of (Y, σ) is $g^*\beta$ -closed set in (X, τ) but not closed. Therefore f is $g^*\beta$ – continuous but not continuous.

Theorem 4.4: Every $g^*\beta$ – continuous map is (1) rg- continuous (2) gp- continuous (3) gpr- continuous (4) gsp- continuous (5) wg- continuous.

Proof: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be $g^*\beta$ – continuous map. Let V be a closed set of (Y, σ) . Since f is $g^*\beta$ – continuous by prop (3.9) $f^{-1}(V)$ is (1) rg-closed (2) gp-closed (3) gpr-closed (4) gsp-closed (5) wg-closed of (X, τ) . Therefore f is rg- continuous, gp- continuous, gpr- continuous, gsp- continuous, wg- continuous.

Example 4.5: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$, $\sigma = \{\emptyset, Y, \{b, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ is defined as the identity map. Then $f^{-1}(\{a\}) = \{a\}$ is not $g^*\beta$ -closed set in (X, τ) . But $\{a\}$ is rg-closed and gpr-closed. Therefore f is rg- continuous and gpr- continuous.

Example 4.6: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$, $\sigma = \{\emptyset, Y, \{b, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ is defined as the identity map. Then $f^{-1}(\{a\}) = \{a\}$ is not $g^*\beta$ -closed set in (X, τ) . But $\{a\}$ is gsp-closed. Therefore f is gsp- continuous.

Example 4.7: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, c\}\}$, $\sigma = \{\emptyset, Y, \{b, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ is defined as the identity map. Then $f^{-1}(\{a\}) = \{a\}$ is not $g^*\beta$ -closed set in (X, τ) . But $\{a\}$ is wg-closed and gp-closed. Therefore f is wg- continuous and gp- continuous. Thus the class of $g^*\beta$ -continuous maps is properly contained in the class of rg- continuous, gp- continuous, gsp- continuous and wg- continuous.

Theorem 4.8: Every g^* -continuous map is $g^*\beta$ -continuous.

Proof: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be g^* -continuous map. Let V be a closed set of (Y, σ) . Then $f^{-1}(V)$ is g^* -closed and hence by prop (3.5) it is $g^*\beta$ -closed set. Hence f is $g^*\beta$ -continuous map.

Example 4.9: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$, $\sigma = \{\emptyset, Y, \{b\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ is defined as the identity map. Here $A = \{a, c\}$ is closed in (Y, σ) . Then $f^{-1}(\{a, c\}) = \{a, c\}$ is $g^*\beta$ -closed set in (X, τ) but not g^* -closed in (X, τ) . Therefore f is $g^*\beta$ -continuous but not g^* -continuous.

Theorem 4.10: Every g-continuous map is $g^*\beta$ -continuous.

Proof: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be g-continuous map. Let V be a closed set of (Y, σ) . Then $f^{-1}(V)$ is g-closed and hence by prop (3.6) it is $g^*\beta$ -closed set. Hence f is $g^*\beta$ -continuous map.

Example 4.11: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$, $\sigma = \{\emptyset, Y, \{a, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ be the identity map. Here $A = \{b\}$ is closed in (Y, σ) . Then $f^{-1}(\{b\}) = \{b\}$ is $g^*\beta$ -closed set in (X, τ) but not g-closed in (X, τ) . Therefore f is $g^*\beta$ -continuous but not g-continuous.

Theorem 4.12: Every g^{**} -continuous map is $g^*\beta$ -continuous.

Proof: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be g^{**} -continuous map. Let V be a closed set of (Y, σ) . Then $f^{-1}(V)$ is g^{**} -closed and hence by prop (3.4) it is $g^*\beta$ -closed set. Hence f is $g^*\beta$ -continuous map.

The following example helps that the converse of the above theorem need not be true in general.

Example 4.13: $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$, $\sigma = \{\emptyset, Y, \{a, c\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ be the identity map. Here $A = \{b\}$ is closed in (Y, σ) . Then $f^{-1}(\{b\}) = \{b\}$ is $g^*\beta$ -closed set in (X, τ) but not g^{**} -closed in (X, τ) . Therefore f is $g^*\beta$ -continuous but not g^{**} -continuous.

Definition 4.14: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $g^*\beta$ -irresolute if $f^{-1}(V)$ is $g^*\beta$ -closed set of (X, τ) for every $g^*\beta$ -closed set of (Y, σ) .

Definition 4.15: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $g^*\beta$ -resolute if $f(U)$ is $g^*\beta$ -open in Y whenever U is $g^*\beta$ -open in X .

Definition 4.16: A function $f: (X, \tau) \rightarrow (Y, \sigma)$ is called $g^*\beta$ -homeomorphism if

- i. f is one and onto
- ii. f is $g^*\beta$ -irresolute and $g^*\beta$ -resolute.

Theorem 4.17: Every $g^*\beta$ -irresolute function is $g^*\beta$ -continuous.

Proof follows from the definition.

Theorem 4.18: Every g -irresolute function is $g^*\beta$ -continuous.

Proof follows from the definition.

Theorem 4.19: Every g^* -irresolute function is $g^*\beta$ -continuous.

Proof follows from the definition.

Theorem 4.20: Every g^{**} -irresolute function is $g^*\beta$ -continuous.

Proof follows from the definition.

Example 4.21: Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$, $\sigma = \{\emptyset, Y, \{a, b, d\}\}$, $f: (X, \tau) \rightarrow (Y, \sigma)$ be defined by $f(a) = b$, $f(b) = a$ and $f(c) = c$. $\{c\}$ is the only closed set of Y . $f^{-1}(\{c\}) = \{c\}$ is $g^*\beta$ -closed set in (X, τ) . Hence f is $g^*\beta$ -continuous map. But $f^{-1}(\{c\}) = \{c\}$ is not g -closed, g^* -closed and g^{**} -closed in X . Therefore f is not g -irresolute, g^* -irresolute, g^{**} -irresolute. Therefore f is $g^*\beta$ -continuous but not g -irresolute, g^* -irresolute and g^{**} -irresolute.

Example 4.22: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a, c\}\}$, $\sigma = \{\emptyset, Y, \{b\}\}$, let $f: (X, \tau) \rightarrow (Y, \sigma)$ be defined by $f(a) = b$, $f(b) = a$ and $f(c) = c$. $\{a, c\}$ is the only closed set of Y . $f^{-1}(\{a, c\}) = \{b, c\}$ is $g^*\beta$ -closed set in (X, τ) . Hence f is $g^*\beta$ -continuous map. But $\{c\}$ is $g^*\beta$ -closed set in Y , $f^{-1}(\{c\}) = \{c\}$ is not $g^*\beta$ -closed set in (X, τ) . Hence $g^*\beta$ -continuous but not $g^*\beta$ -irresolute.

Theorem 4.23: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ and $g: (Y, \sigma) \rightarrow (Z, \rho)$ be any two functions. Then

- i. $g \circ f$ is $g^*\beta$ -continuous if g is continuous and f is $g^*\beta$ -continuous.
- ii. $g \circ f$ is $g^*\beta$ -irresolute if both f and g are $g^*\beta$ -irresolute.
- iii. $g \circ f$ is $g^*\beta$ -continuous if g is $g^*\beta$ -continuous and f is $g^*\beta$ -irresolute.

5. APPLICATIONS OF $g^*\beta$ -CLOSED SETS

As applications of $g^*\beta$ -closed sets, new spaces namely new spaces namely ${}_{\beta}T_{1/2}^{**}$ -space, ${}_{\alpha\beta}T_c^*$ -space, ${}_{\beta}T_{1/2}^*$ -space, are introduced.

Definition 5.1: A space (X, τ) is called ${}_{\beta}T_{1/2}^{**}$ -space if every $g^*\beta$ -closed set is closed.

Theorem 5.2: Every ${}_{\beta}T_{1/2}^{**}$ -space is a $T_{1/2}$ -space.

Proof follows from the definition.

Theorem 5.3: Every ${}_{\beta}T_{1/2}^{**}$ -space is a $T_{1/2}^*$ -space.

Proof follows from the definition.

The converse need not be true in general as seen in the following example.

Example 5.4: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$. $G^*\beta Cl(x, \tau) = \{\emptyset, X, \{b, c\}\} = C(X, \tau)$. Therefore (X, τ) is a $T_{1/2}^*$ -space but not a ${}_{\beta}T_{1/2}^{**}$ -space. Since $\{a, c\}$ is $g^*\beta$ -closed set but not closed in (X, τ) .

Theorem 5.5: Every T_b -space is a $\beta T_{1/2}^{**}$ -space.

Proof follows from the definition.

The converse need not be true in general as seen in the following example.

Example 5.6: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$. (X, τ) is a $\beta T_{1/2}^{**}$ -space but not a T_b -space. Since $A = \{b\}$ is $g^*\beta$ -closed set but not closed in (X, τ) .

Remark 5.7: T_d -ness is independent of $\beta T_{1/2}^{**}$ -ness as it can be seen from the following examples.

Example 5.8: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$. (X, τ) is a $\beta T_{1/2}^{**}$ -space but not a T_d -space. Since $A = \{a\}$ is $g^*\beta$ -closed set but not g -closed in (X, τ) .

Example 5.9: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b, c\}\}$. (X, τ) is a T_d -space but not a $\beta T_{1/2}^{**}$ -space. Since $A = \{c\}$ is $g^*\beta$ -closed but not closed.

Theorem 5.10: The following conditions are equivalent in topological space (X, τ) .

- i. (X, τ) is a $\beta T_{1/2}^{**}$ -space.
- ii. Every singleton set of X is either g^* -closed or open.

Proof:

(i) \Rightarrow (ii): Let (X, τ) be a $\beta T_{1/2}^{**}$ -space. Let $x \in X$ and suppose $\{x\}$ is not g^* -closed. Then $X \setminus \{x\}$ is not g^* -open. This implies that X is the only g^* -open set containing $X \setminus \{x\}$. Therefore $X \setminus \{x\}$ is closed since (X, τ) is a $\beta T_{1/2}^{**}$ -space. Therefore $\{x\}$ is open in (X, τ) .

(ii) \Rightarrow (i): Let A be a $g^*\beta$ -closed of (X, τ) . $A \subseteq \beta cl(A) \subseteq cl(A)$ and let $x \in \beta cl(A)$ this implies $x \in cl(A)$. By (ii) $\{x\}$ is g^* -closed or open.

Case-(i): Let $\{x\}$ be g^* -closed. If x does not belong to A then $\beta cl(A) \setminus A$ contains a nonempty g^* -closed set $\{x\}$. But it is not possible by proposition (3.14). Therefore $x \in A$.

Case-(ii): Let $\{x\}$ be open. Now $x \in cl(A)$, then $\{x\} \cap A = \emptyset$. Therefore $x \notin A$ and so $cl(A) \subseteq A$ and hence $A = cl(A)$ or A is closed. Therefore (X, τ) is a $\beta T_{1/2}^{**}$ -space.

We introduce the following definition.

Definition 5.11: A space (X, τ) is called $_{\alpha\beta}T_c^*$ -space if every βg -closed set is $g^*\beta$ -closed.

Theorem 5.12: Every βT_b -space is a $_{\alpha\beta}T_c^*$ -space but not conversely.

Example 5.13: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$ is $_{\alpha\beta}T_c^*$ -space but not a βT_b -space but not a βT_b -space since $\{a, c\}$ is βg -closed but not closed.

Definition 5.14: A subset A of (X, τ) is called $g^*\beta$ -open set if its complement is $g^*\beta$ -closed set of (X, τ) .

Theorem 5.15: If (X, τ) is a $_{\alpha\beta}T_c^*$ -space for each $x \in X$, $\{x\}$ is either βg -closed or $g^*\beta$ -open.

Proof: Let $x \in X$ suppose that $\{x\}$ is not βg -closed of (X, τ) . Then $\{x\}$ is not closed set since every closed set is a βg -closed set. Therefore, $X \setminus \{x\}$ is not open. Therefore $X \setminus \{x\}$ is a βg -closed set since X is the only open set which contains $X \setminus \{x\}$. Since (X, τ) is a $_{\alpha\beta}T_c^*$ -space, $X \setminus \{x\}$ is $g^*\beta$ -closed

or $\{x\}$ is $g^*\beta$ -open.

We introduce the following definition.

Definition 5.16: A space (X, τ) is called $^{**}\beta T_{1/2}$ -space, if every $g^*\beta$ -closed set is g^* -closed.

Theorem 5.17: Every $\beta T_{1/2}^{**}$ -space is a $^{**}\beta T_{1/2}$ -space.

Proof: Let (X, τ) be a $\beta T_{1/2}^{**}$ -space. Let A be a $g^*\beta$ -closed set of (X, τ) . Since (X, τ) is a $\beta T_{1/2}^{**}$ -space, A is closed. Since every closed set is g^* -closed, (X, τ) is a $^{**}\beta T_{1/2}$ -space.

Theorem 5.18: Every T_b -space is a ${}^{\beta}T_{1/2}^{**}$ -space.

Proof: Let (X, τ) be a T_b -space. Then by theorem 5.5, it is ${}^{\beta}T_{1/2}^{**}$ -space. Therefore by theorem 5.19, it is ${}^{\beta}T_{1/2}^{**}$ -space. The converse need not be true in general as seen in the following example.

Example 5.19: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$. (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space but not a T_b -space. Since $A = \{a\}$ is g^* -closed set but not closed in (X, τ) .

Theorem 5.20: Every ${}^{\beta}T_{1/2}^{**}$ -space is a ${}^*T_{1/2}$ -space.

Proof: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let A be a g -closed set of (X, τ) . Then by prop (3.6) A is $g^*\beta$ -closed. Since (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space, A is g^* -closed. Therefore it is a ${}^*T_{1/2}$ -space.

The converse of the above theorem need not be true as seen in the following example.

Example 5.21: Let $X = Y = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}, \{a, c\}\}$. (X, τ) is a ${}^*T_{1/2}$ -space but not a ${}^{\beta}T_{1/2}^{**}$ -space. Since $A = \{c\}$ is $g^*\beta$ -closed but not g^* -closed.

Theorem 5.22: Every ${}^{\beta}T_{1/2}^{**}$ -space is a ${}^{\beta}T_{1/2}^{**}$ -space.

Proof: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let A be a g^{**} -closed set of (X, τ) . Then by prop (3.4) A is $g^*\beta$ -closed. Since (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space, A is g^* -closed. Therefore it is a ${}^{\beta}T_{1/2}^{**}$ -space.

The converse of the above theorem need not be true as seen in the following example.

Example 5.23: Let $X = \{a, b, c, d\}$, $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$. Here (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space but not a ${}^{\beta}T_{1/2}^{**}$ -space. Since $A = \{c\}$ is $g^*\beta$ -closed but not g^* -closed.

Theorem 5.24: If (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space for each $x \in X$, $\{x\}$ is either closed or g^* -open.

Proof: Suppose (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let $x \in X$ and let $\{x\}$ not be closed set. Then $X \setminus \{x\}$ is not open set. Therefore $X \setminus \{x\}$ is a ${}^{\beta}T_{1/2}^{**}$ -space, $X \setminus \{x\}$ is g^* -closed set. Therefore $\{x\}$ is g^* -open.

Definition 5.25: A space (X, τ) is called ${}^{\beta}T_{1/2}^{**}$ -space if every $g^*\beta$ -closed set is g -closed.

Theorem 5.26: Every ${}^{\beta}T_{1/2}^{**}$ -space is a ${}^{\beta}T_{1/2}^{**}$ -space.

Proof: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let A be a $g^*\beta$ -closed set of (X, τ) . Then A is closed. Since (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space. But every closed set is a g -closed set, therefore A is g -closed. Therefore (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space. The converse of the above theorem need not be true as seen in the following example.

Example 5.27: Let $X = \{a, b, c\}$, $\tau = \{\emptyset, X, \{a\}\}$. (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space but not a ${}^{\beta}T_{1/2}^{**}$ -space. Since $A = \{a, b\}$ is $g^*\beta$ -closed set but not closed in (X, τ) .

Theorem 5.28: The space (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space iff it is a ${}^{\beta}T_{1/2}^{**}$ -space and a $T_{1/2}$ -space.

Proof: Necessity: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let A be a g -closed set of (X, τ) . Then by prop (3.6) A is $g^*\beta$ -closed. Also since (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space, A is closed set. Therefore (X, τ) is a $T_{1/2}$ -space. By theorem (5.26) (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space.

Sufficiency: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space and a $T_{1/2}$ -space. Let A be a $g^*\beta$ -closed set. Then A is g -closed. Since (X, τ) is a $T_{1/2}$ -space, A is a closed set. Therefore (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space.

Theorem 5.29: Every ${}^{\beta}T_{1/2}^{**}$ -space is a ${}^{\beta}T_{1/2}^{**}$ -space.

Proof: Let (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space. Let A be a $g^*\beta$ -closed set. Then A is g^* -closed since (X, τ) is a ${}^{\beta}T_{1/2}^{**}$ -space. But every g^* -closed is g -closed and hence A is a g -closed set. Therefore (X, τ) be a ${}^{\beta}T_{1/2}^{**}$ -space.

Example 5.30: Let $X = \{a, b, c\}$, $\tau = \{\varphi, X, \{a\}\}$. (X, τ) is a ${}^*T_{1/2}^*$ -space but not a ${}^{**}T_{1/2}$ -space. Since $A = \{b\}$ is $g^*\beta$ -closed but not g^* -closed.

We introduce the following definition

Definition 5.31: A space (X, τ) is called ${}^*T_c^*$ -Space if every g s-closed of (X, τ) is a $g^*\beta$ -closed.

Theorem 5.32: A T_c -space is a ${}^*T_c^*$ -Space.

Proof: Let (X, τ) be a T_c -space. Let A be a g s-closed set of (X, τ) . Then A is g^* -closed.

Since (X, τ) be a T_c -space, by proposition (3.5), then A is $g^*\beta$ -closed set. Therefore (X, τ) is a ${}^*T_c^*$ -Space.

Example 5.33: Let $X = \{a, b, c\}$, $\tau = \{\varphi, X, \{c\}\}$. (X, τ) is a ${}^*T_c^*$ -space but not a T_c -space. Since $A = \{b\}$ is g s-closed but not g^* -closed.

Theorem 5.34: A T_b -space is a ${}^*T_c^*$ -Space.

Proof: Let (X, τ) be a T_b -space. Let A be a g s-closed set of (X, τ) . Then A is closed. Since (X, τ) be a T_b -space, by proposition (3.2), A is $g^*\beta$ -closed set. Therefore (X, τ) is a ${}^*T_c^*$ -Space.

Example 5.35: Let $X = \{a, b, c\}$, $\tau = \{\varphi, X, \{c\}\}$. (X, τ) is a ${}^*T_c^*$ -space but not a T_b -space. Since $A = \{b\}$ is g s-closed but not a closed set.

Theorem 5.36: If (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^*$ -Space then it is a *T_d -Space.

Proof: Let (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^*$ -Space. Let A be a βg -closed set of (X, τ) . Then A is also g s-closed. Since (X, τ) is a ${}^*T_c^*$ -Space, A is a $g^*\beta$ -closed set. Also since (X, τ) is a ${}^*T_{1/2}^*$ -Space, A is g -closed set. Therefore (X, τ) is a *T_d -Space.

The following example helps that the converse of the above theorem need not be true in general.

Example 5.37: Let $X = \{a, b, c\}$, $\tau = \{\varphi, X, \{a\}, \{b\}, \{a, b\}\}$. (X, τ) is a *T_d -space but not a ${}^*T_c^*$ -space. Since $A = \{b\}$ is g s-closed but not a $g^*\beta$ -closed set.

Theorem 5.38: If (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^{**}$ -Space then it is a T_b -Space.

Proof: Let (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^{**}$ -Space. Let A be a βg -closed set of (X, τ) . Then A is also g s-closed. Since (X, τ) is a ${}^*T_c^*$ -Space, A is a $g^*\beta$ -closed set. But every $g^*\beta$ -closed set is closed. Also since (X, τ) is a ${}^*T_{1/2}^{**}$ -Space, A is closed set. Therefore (X, τ) is a T_b -Space.

Example 5.39: Let $X = \{a, b, c\}$, $\tau = \{\varphi, X, \{a\}, \{b\}, \{a, b\}\}$. (X, τ) is a T_b -space but not a ${}^*T_c^*$ -space. Since $A = \{b\}$ is g s-closed but not a $g^*\beta$ -closed set.

Theorem 5.40: If (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^*$ -Space then it is a T_d -Space.

Proof: Let (X, τ) is a ${}^*T_c^*$ -Space and ${}^*T_{1/2}^*$ -Space. Let A be a g s-closed set of (X, τ) .

Since (X, τ) is a ${}^*T_c^*$ -Space, A is a $g^*\beta$ -closed set. Also since (X, τ) is a ${}^*T_{1/2}^*$ -Space, A is g -closed set. Therefore (X, τ) is a T_d -Space.

Theorem 5.41: If (X, τ) is a ${}^*T_c^*$ -Space, then for each $x \in X$, $\{x\}$ is either semi-closed or $g^*\beta$ -open.

Proof: Suppose (X, τ) be a ${}^*T_c^*$ -Space. Let $x \in X$ and let $\{x\}$ not be semiclosed. Then $X \setminus \{x\}$ is g s-closed. Also $X \setminus \{x\}$ is g s-closed. Since (X, τ) is a $g^*\beta$ -closed, $\{x\}$ is $g^*\beta$ -open.

Theorem 5.42: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $g^*\beta$ -continuous map. If (X, τ) is a ${}^*T_{1/2}^{**}$ -space, then f is continuous.

Theorem 5.43: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $g^*\beta$ -continuous map. If (X, τ) is a ${}^{**}T_{1/2}$ -space, then f is g^* -continuous.

Theorem 5.44: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $g^*\beta$ - continuous map. If (X, τ) is a ${}^*T_{1/2}$ - space, then f is g -continuous.

Theorem 5.45: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a gs - continuous map. If (X, τ) is a βT_c^* - space, then f is $g^*\beta$ -continuous.

Theorem 5.46: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a g^* - irresolute map and a β - closed map. Then $f(A)$ is a $g^*\beta$ -closed set of (Y, σ) for every $g^*\beta$ - closed set A of (X, τ) .

Proof: Let A be a $g^*\beta$ - closed set of (X, τ) . Let U be a g^* - open set of (Y, σ) such that $f(A) \subseteq U$. Since f is g^* - irresolute, $f^{-1}(U)$ is g^* - open in (X, τ) . Now $f^{-1}(U)$ is g^* - open and A is $g^*\beta$ - closed set of (X, τ) , then $\beta cl(A) \subseteq f^{-1}(U)$. Then $f(\beta cl(A)) = \beta cl[f(\beta cl(A))]$. Therefore $\beta cl[f(A)] \subseteq \beta cl[f(\beta cl(A))] = f(\beta cl(A)) \subseteq U$. Therefore $f(A)$ is a $g^*\beta$ -closed set of (Y, σ) for every $g^*\beta$ - closed set A of (X, τ) .

Theorem 5.47: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $g^*\beta$ - irresolute and closed. If (X, τ) is a $\beta T_{1/2}^{**}$ -space, then (Y, σ) is also a $\beta T_{1/2}^{**}$ -space.

Theorem 5.48: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be a $g^*\beta$ - closed map if $f(A)$ is $g^*\beta$ - closed set of (Y, σ) for every $g^*\beta$ - closed set of (X, τ) .

Theorem 5.49: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be onto $g^*\beta$ - irresolute and pre- g^* -closed. If (X, τ) is a ${}^{**}T_{1/2}$ - space, then (Y, σ) is also a ${}^{**}T_{1/2}$ - space.

Proof follows from the definition of $g^*\beta$ - irresolute and pre- g^* -closed.

Theorem 5.50: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be onto gs - irresolute and $g^*\beta$ -closed map. If (X, τ) is a βT_c^* - space, then (Y, σ) is also a βT_c^* - space.

Proof follows from the definition of gs - irresolute and $g^*\beta$ closed map.

Theorem 5.51: Let $f: (X, \tau) \rightarrow (Y, \sigma)$ be onto $g^*\beta$ - irresolute and g -closed. If (X, τ) is a ${}^*T_{1/2}$ - space, then (Y, σ) is also a ${}^*T_{1/2}$ - space.

Proof follows from the definition of $g^*\beta$ - irresolute and g closed map.

REFERENCES

1. Andrijevic. D, Semi-preopen sets, Mat. Vesnik, 38(1) (1986), 24-32.
2. Arockiarani. I, Balachandran. K and Dontchev. J, Some characterization of gp -irresolute and gp -continuous topological spaces, Mem.Fac.Sci.kchi.Univ.Ser.A.Math., 20(1999), 93-104.
3. Arya. S. P and T. Nour, Characterizations of s -normal spaces, Indian J.Pure.Appl.Math., 21(8)(1990),717-719.
4. Balachandran. K, Sundram. P and Maki. H, On generalized continuous maos in topological spaces, Mem.Fac.KochiUniv.Ser.A.Math., 12(1991), 5-13.
5. Bhattacharya. P and Lahiri. B. K., semi-generalised closed sets in topology, Indian J.Math., 29(3) (1987), 375-382.
6. Devi. R, Maki. H and Balachandran. K, semi-generalized closed maps and generalized closed maps, Mem.Fac.Sci.KochiUniv.Ser.A.Math., 14(1993), 41-54.
7. Devi. R, Maki. H and Balachandran. K, semi-generalized homeomorphism and generalized semi homeomorphism topological spaces, Indian J.Pure.Appl.Math., 26(3) 1995, 271-284.
8. Devi. R, Maki. H and Balachandran. K, Generalised α -closed maps and α -generalized closed maps, Indian J.Pure.Appl.Math., 29(1) (1998), 37-49.
9. Dontchev. J, On generalizing semipre open set, Mem.Fac.Sci.KochiUniv.Ser.A.Math., 16(1995), 35-48.
10. Gnanambal. Y, On Generalized Preregular closed sets in topological spaces, Indian J.Pure.Appl.Math., 28(3) (1997), 351-360.
11. Levine. N, Generalized closed sets in topology, Rend.Circ.Math.Palermo, 19(2) (1970), 89-96.
12. Levine. N, Semi-open sets and semi-continuity in topological spaces, Amer.Math.Monthly, 70(1963), 36-41.
13. Maki. H, Umehara. J and Noiri. T, Every topological spaces in pre- $T_{1/2}$, Mem.Fac.Sci.Kochi Univ.Ser.A.Math., 17(1996), 33-42.
14. Maki. H, Devi. R and Balachandran. K, Associated topologies of Generalized α -closed set and α -generalized closed sets Mem.Fac.Sci.Koc
15. Maki. H, Devi. R and Balachandran. K, Generalized α -closed sets in topology, Bull.FukuokaUniv.Ed.Part III, 42(1993), 13-21.
16. Mashhour.A. S., M.E.Abd El-Monsef and S.N.El-Deeb, On Pre-continuous and weak pre continuous mappings, Proc.Math. And Phys.soc.Egypt, 53(1982), 47-53.
17. Nagaveni. N, studies On Generalizations of Homeomorphisms in Topological Spaces, Ph.D, thesis, Bharathiar University, Coimbatore, 1999.

18. Njastad. O, On Some classes of nearly open sets, Pacific J.Math., 15(1965), 961-970.
19. Palaniappan. N and Rao. K. C., Regular generalized closed sets, Kyungpook Math.J., 33(2) (1993), 211-219.
20. Pauline Mary Helen. M, g^{**} -closed sets in Topological spaces, International Journal of Mathematical Archive -3(5), 2012, 2005-2019.
21. Punitha Tharani, Priscilla Pacifica, pg^{**} -Closed sets in Topological Spaces, International Journal of Mathematical Archieve-6 (7), 2015, 128- 137.
22. Veerakumar. M. K. R. S, Between Closed sets and g -closed sets, Mem. Fac. Sci. Koch. Univ. Ser. A, Math., 17 (19916). 33-42.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2020. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]