

CO-FUZZY IDEALS OF FINITE Γ -NEAR RING

DR. D. BHARATHI¹, P. VENKATRAO*² AND K. BALAKOTESWARA RAO³

^{1,2}Department of Mathematics, S. V. University, Tirupati – (A.P.), India.

³Department of Mathematics, P. R. R & V S Govt. College, Vidavaluru, India.

(Received On: 03-03-21; Revised & Accepted On: 12-04-21)

ABSTRACT

In this paper we introduce and study the concept of ideals of finite Γ -near rings, co-fuzzy ideals of finite Γ -near rings and related theorems. We prove intersection of two co-fuzzy ideals is always a co-fuzzy ideal but union of two co-fuzzy ideals need not be a co-fuzzy ideal of finite Γ -near ring and it is explained by a suitable example. We also introduce homomorphic image and homomorphic pre image of co-fuzzy ideals of finite Γ -near rings and related theorems.

Key Words : finite Γ -near ring, fuzzy set, level subset, union of fuzzy sets, intersection of fuzzy sets, co-fuzzy ideals and homomorphism of co-fuzzy ideals.

1. INTRODUCTION

The fuzzy set theory was developed by Zadeh.L.A. [12] in 1965. The fuzzification of algebraic structure was introduced by Rosenfield.A[7] and he introduced the notation of fuzzy of subgroups in 1971. Swamy K.L.N and Swamy U.M [8] studied fuzzy prime ideals [4,6] Later Satyanarayana. Bh [9] defined Γ -near rings and also he studied ideal theory in Γ -near rings. The notation of fuzzy ideals and its properties were applied to various areas like semi groups [11, 10, 5] and semi rings [1,2] Jun.Y.B [3] considered the fuzzification of ideals of Γ -near rings and investigated the related properties.

In this paper, we have defined co-fuzzy ideal concept of finite Γ –near ring with less than or equal and maximum conditions and also investigated several properties with the new definitions. Throughout this chapter \mathcal{N} stands for zero symmetric finite Γ – near ring.

2. PRELIMINARIES

In this section we recall some of the fundamental definitions, which are necessary for this paper.

Definition 2.1: A triplet $(\mathcal{N}, +, \cdot)$ is said to be a near ring if

1. $(\mathcal{N}, +)$ is a group
2. (\mathcal{N}, \cdot) is a semi-group
3. $(a + b) \cdot c = a \cdot c + b \cdot c$ (Right distribution law) $\forall a, b, c \in \mathcal{N}$.

Definition 2.2: A near ring $(\mathcal{N}, +, \cdot)$ is said to be finite near ring if \mathcal{N} has finite number of elements.

Definition 2.3: Let \mathcal{N} be a non-empty finite set. If

1. $(\mathcal{N}, +)$ is a group. (Not necessarily abelian group)
2. Γ is the set of binary operations on \mathcal{N} such that $(\mathcal{N}, +, \alpha)$ is a near ring. Where $\alpha \in \Gamma$.
3. $a\alpha(b\beta c) = (a\alpha b)\beta c, \forall a, b, c \in \mathcal{N}$ and $\alpha, \beta \in \Gamma$. Then $(\mathcal{N}, +, \Gamma)$ is called finite Γ -near ring.

Corresponding Author: P. Venkatrao*²,

²Department of Mathematics, S. V. University, Tirupati - (A.P.), India.

Definition 2.4: If X be a non empty set and $f : X \rightarrow [0, 1]$ is a mapping then the pair (X, f) is called fuzzy set and f is called fuzzy sub set of X .

Definition 2.5: Let f be a fuzzy sub set of the set X . Then the set $\{x \in X / f(x) < s\}$ Where $s \in [0, 1]$ is called level sub set of f . It is denoted by f_s . This f_s is also s -cut of f .

$$\therefore f_s = \{x \in X / f(x) < s\}$$

Definition 2.6: Let f and g be two fuzzy sub sets of the set X . If $f(x) \leq g(x)$, $\forall x \in X$ then f is said to be contained in g . It is denoted by $f \subseteq g$.

Definition 2.7: Let f and g be two fuzzy sub sets of the set X . Then their intersection and union are denoted by $f \cap g$ and $f \cup g$ respectively and defined as follows,

$$(f \cap g)(x) = \text{Min} \{f(x), g(x)\}, \quad \forall x \in X.$$

and

$$(f \cup g)(x) = \text{Max} \{f(x), g(x)\}, \quad \forall x \in X.$$

Definition 2.8: Let M_1 and M_2 be two non empty sets and $\mu: M_1 \rightarrow M_2$ is a mapping. If f is a fuzzy sub set of M_1 then g be a fuzzy sub set of M_2 defined by

$$g(y) = \begin{cases} \text{Inf}_{z \in \mu^{-1}(y)} f(z) & \text{if } \mu^{-1}(y) \neq \phi \\ 0 & \text{if } \mu^{-1}(y) = \phi \end{cases}$$

Where $\mu^{-1}(y) = \{x \in M_1 / \mu(x) = y\}$

If g is a fuzzy sub set of M_2 then f be a fuzzy sub set of M_1 defined by

$$f(x) = g(\mu(x)), \quad \forall x \in M_1.$$

Definition 2.9: For any sub set A of the set X , the co-fuzzy characteristic set δ_A is defined as follows,

$$\delta_A(x) = \begin{cases} 0 & \text{if } x \in A \\ 1 & \text{if } x \notin A \end{cases}$$

3. CO - FUZZY IDEALS OF FINITE Γ -NEAR RINGS

In this section we define ideals and co-fuzzy ideals of a finite Γ -near rings. We prove the intersection of two co-fuzzy ideals of a finite Γ -near ring is always a co-fuzzy ideal. But, the union of two co-fuzzy ideals of a finite Γ -near ring need not be a co-fuzzy ideal and it is explained by a suitable example. We also prove some of related theorems on ideals and co-fuzzy ideals of a finite Γ -near rings.

Definition 3.1: A sub set S of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be a left ideal of \mathcal{N} if

1. $x + y \in S, \quad \forall x, y \in S$
2. $y + x - y \in S, \quad \forall x \in S \text{ and } y \in \mathcal{N}$
3. $a\alpha(x + b) - aab \in S, \quad \forall x \in S, a, b \in \mathcal{N} \text{ and } \alpha \in \Gamma.$

Definition 3.2: A sub set S of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be a right ideal of \mathcal{N} if

1. $x + y \in S, \quad \forall x, y \in S$
2. $y + x - y \in S, \quad \forall x \in S \text{ and } y \in \mathcal{N}$
3. $x\alpha a \in S, \quad \forall x \in S, a \in \mathcal{N} \text{ and } \alpha \in \Gamma.$

Definition 3.3: A sub set S of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be an ideal of \mathcal{N} if

1. $x + y \in S, \quad \forall x, y \in S$
2. $y + x - y \in S, \quad \forall x \in S \text{ and } y \in \mathcal{N}$
3. $a\alpha(x + b) - aab \in S, \quad \forall x \in S, a, b \in \mathcal{N} \text{ and } \alpha \in \Gamma.$
4. $x\alpha a \in S, \quad \forall x \in S, a \in \mathcal{N} \text{ and } \alpha \in \Gamma.$

Definition 3.4: A fuzzy sub set f of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be a co-fuzzy left ideal of \mathcal{N} if

1. $f(x + y) \leq \text{Max} \{f(x), f(y)\},$
2. $f(y + x - y) \leq f(x),$
3. $f(a\alpha(x + b) - aab) \leq f(x), \quad \forall x, y, a, b \in \mathcal{N} \text{ and } \alpha \in \Gamma.$

Definition 3.5: A fuzzy sub set f of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be a co-fuzzy right ideal of \mathcal{N} if

1. $f(x + y) \leq \text{Max} \{f(x), f(y)\}$,
2. $f(y + x - y) \leq f(x)$,
3. $f(x\alpha a) \leq f(x)$, $\forall x, y, a \in \mathcal{N}$ and $\alpha \in \Gamma$.

Definition 3.6: A fuzzy sub set f of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be a co-fuzzy ideal of \mathcal{N} if

1. $f(x + y) \leq \text{Max} \{f(x), f(y)\}$,
2. $f(y + x - y) \leq f(x)$,
3. $f(a\alpha(x + b) - a\alpha b) \leq f(x)$,
4. $f(x\alpha a) \leq f(x)$, $\forall x, y, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$.

Definition 3.7: A L -fuzzy sub set f_L of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be L -fuzzy ideal of \mathcal{N} if

1. $f_L(x + y) \leq \text{Max} \{f_L(x), f_L(y)\}$
2. $f_L(y + x - y) \leq f_L(x)$
3. $f_L(a\alpha(x + b) - a\alpha b) \leq f_L(x)$
4. $f_L(x\alpha a) \leq f_L(x)$ $\forall x, y, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$.

Here L is a complete lattice satisfying infinite distribute laws.

Definition 3.8: A finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is said to be zero symmetric if $x\alpha 0 = 0\alpha x = 0$ $\forall x \in \mathcal{N}$

Note 3.9:

1. $f(x + y) \leq \text{Max} \{f(x), f(y)\}$
2. $f(x - y) \leq \text{Max} \{f(x), f(-y)\}$
3. $f(0) \leq \text{Max} \{f(x), f(-x)\}$
4. $f(0) = f(x\alpha 0) \leq f(x)$, $\forall x, y \in \mathcal{N}$

Example 3.10: Let $S = \{1, 2, 3, 4\}$ and $\mathcal{N} = P(S)$ is the power set of S .

Let $\Gamma = \{\{1\}, \{2, 3\}\}$

Define $f: \mathcal{N} \rightarrow [0, 1]$ such that $f(A) = \begin{cases} \frac{1}{2} & \text{if } A \neq \phi \\ \frac{1}{3} & \text{if } A = \phi \end{cases}$

Here f is both co-fuzzy left and co-fuzzy right ideal of finite Γ -near ring $(\mathcal{N}, \Delta, \Gamma)$.

Hence f is co-fuzzy ideal of finite Γ -near ring $(\mathcal{N}, \Delta, \Gamma)$.

Example 3.11: Let \mathcal{N} be the set of all 2×2 matrices defined over Z_5 . Where $Z_5 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}\}$ is the set of all residue classes modulo 5 and $\Gamma = \{\bar{1}, \bar{2}, \bar{3}\}$. Then $(\mathcal{N}, +, \Gamma)$ is finite Γ -near ring.

Define $f: \mathcal{N} \rightarrow [0, 1]$ such that $f(A) = \begin{cases} 0.3 & \text{if } A = \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} \\ 0.5 & \text{if otherwise} \end{cases}$

Then f is a co-fuzzy right ideal of \mathcal{N} . But not co-fuzzy left ideal of \mathcal{N} .

Define $f: \mathcal{N} \rightarrow [0, 1]$ such that $f(A) = \begin{cases} 0.2 & \text{if } A = \begin{bmatrix} 0 & a \\ 0 & b \end{bmatrix} \\ 0.6 & \text{otherwise} \end{cases}$

Then f is a co-fuzzy left ideal of \mathcal{N} . But not co-fuzzy right ideal of \mathcal{N} .

Now, we prove some important theorems on co-fuzzy ideals.

Theorem 3.12: A fuzzy sub set f of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ is a co-fuzzy ideal of \mathcal{N} if and only if for each $\delta \in \text{Im}(f)$, the level set f_δ of f is an ideal of \mathcal{N} .

Proof: Let f be a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

Now, we have to prove that the level set f_δ of f is an ideal of $\mathcal{N} \forall \delta \in \text{Im}(f)$

By the definition of the level set $f_\delta = \{x \in \mathcal{N}, f(x) < \delta\}$

1. Let $x, y \in f_\delta$
 Then $f(x) < \delta$ and $f(y) < \delta$
 Now $f(x + y) \leq \text{Max}\{f(x), f(y)\} < \delta$
 $\therefore f(x + y) < \delta$
 $\Rightarrow x + y \in f_\delta$
2. Let $x \in f_\delta$ and $y \in \mathcal{N}$
 Then $f(x) < \delta$
 Now $f(y + x - y) \leq f(x) < \delta$
 $\therefore f(y + x - y) < \delta$
 $\Rightarrow y + x - y \in f_\delta$
3. Let $x \in f_\delta$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$
 Then $f(x) < \delta$
 Now $f(a\alpha(x + b) - aab) \leq f(x) < \delta$
 $\therefore f(a\alpha(x + b) - aab) < \delta$
 $\Rightarrow a\alpha(x + b) - aab \in f_\delta$
 Then the level set f_δ of f is a left ideal of $\mathcal{N}, \forall \delta \in \text{Im}(f)$
4. Let $x \in f_\delta$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$
 Then $f(x) < \delta$
 We have $f(x\alpha a) \leq f(x) < \delta$
 $\therefore f(x\alpha a) < \delta$
 $\Rightarrow x\alpha a \in f_\delta$

Then the level set f_δ of f is a right ideal of \mathcal{N} .

Hence the level set f_δ of f is a ideal of $\mathcal{N}, \forall \delta \in \text{Im}(f)$

Conversely assume that f_δ is an ideal of \mathcal{N}

Now we prove that f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

1. Let $x, y \in \mathcal{N}$
 If possible, let there exists $x_0, y_0 \in \mathcal{N}$ such that
 $f(x_0 + y_0) > \text{Max}\{f(x_0), f(y_0)\}$
 Let $\delta = \frac{1}{2}\{f(x_0 + y_0) + \text{Max}\{f(x_0), f(y_0)\}\}$
 Then $\delta < \frac{1}{2}\{f(x_0 + y_0) + f(x_0 + y_0)\}$
 $\Rightarrow \delta < f(x_0 + y_0)$
 $\Rightarrow x_0 + y_0 \notin f_\delta$
 And $\delta > \frac{1}{2}\{2 \text{Max}\{f(x_0), f(y_0)\}\}$
 $\Rightarrow \delta > \text{Max}\{f(x_0), f(y_0)\}$
 $\Rightarrow \delta > f(x_0)$ and $\delta > f(y_0)$
 $\Rightarrow x_0, y_0 \in f_\delta$
 $\therefore x_0 + y_0 \notin f_\delta$ and $x_0, y_0 \in f_\delta$
 This is a contradiction to f_δ is an ideal of \mathcal{N}
 Then our assumption is wrong.
 Hence $f(x + y) \leq \text{Max}\{f(x), f(y)\}, \forall x, y \in \mathcal{N}$

2. Let $x, y \in \mathcal{N}$

If possible, let there exists $x_0, y_0 \in \mathcal{N}$ such that

$$f(y_0 + x_0 - y_0) > f(x_0)$$

$$\text{Let } \delta = \frac{1}{2}\{f(y_0 + x_0 - y_0) + f(x_0)\}$$

$$\text{Then } \delta < \frac{1}{2}\{2f(y_0 + x_0 - y_0)\} \text{ and } \delta > \frac{1}{2}\{2f(x_0)\}$$

$$\Rightarrow \delta < f(y_0 + x_0 - y_0) \text{ and } \delta > f(x_0)$$

$$\Rightarrow y_0 + x_0 - y_0 \notin f_\delta \text{ and } x_0 \in f_\delta$$

This is a contradiction to f_δ is an ideal of \mathcal{N}

Then our assumption is wrong and hence $f(y + x - y) \leq f(x)$,

$$\forall x, y \in \mathcal{N}$$

3. Let $x, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

If possible, let there exists $x_0, a_0, b_0 \in \mathcal{N}$ and $\alpha \in \Gamma$ such that

$$f(a_0\alpha(x_0 + b_0) - a_0\alpha b_0) > f(x_0)$$

$$\text{Let } \delta = \frac{1}{2}\{f(a_0\alpha(x_0 + b_0) - a_0\alpha b_0) + f(x_0)\}$$

$$\text{Then } \delta < \frac{1}{2}\{2f(a_0\alpha(x_0 + b_0) - a_0\alpha b_0)\} \text{ and } \delta > \frac{1}{2}\{2f(x_0)\}$$

$$\Rightarrow \delta < f(a_0\alpha(x_0 + b_0) - a_0\alpha b_0) \text{ and } \delta > f(x_0)$$

$$\Rightarrow a_0\alpha(x_0 + b_0) - a_0\alpha b_0 \notin f_\delta \text{ and } x_0 \in f_\delta$$

This is a contradiction to f_δ is an ideal of \mathcal{N}

Then our assumption is wrong and hence

$$f(a\alpha(x + b) - a\alpha b) \leq f(x), \forall x, a, b \in \mathcal{N} \text{ and } \alpha \in \Gamma.$$

Then f is a co-fuzzy left ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

4. Let $x, a \in \mathcal{N}$ and $\alpha \in \Gamma$

If possible, let there exists $x_0, a_0 \in \mathcal{N}$ and $\alpha \in \Gamma$ such that

$$f(x_0\alpha a_0) > f(x_0)$$

$$\text{Let } \delta = \frac{1}{2}\{f(x_0\alpha a_0) + f(x_0)\}$$

$$\text{Then } \delta < f(x_0\alpha a_0) \text{ and } \delta > f(x_0)$$

$$\Rightarrow x_0\alpha a_0 \notin f_\delta \text{ and } x_0 \in f_\delta$$

This is a contradiction to f_δ is an ideal of \mathcal{N}

Then our assumption is wrong and hence $f(x\alpha a) \leq f(x), \forall x, a \in \mathcal{N} \text{ and } \alpha \in \Gamma$

Then f is a co-fuzzy right ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Hence f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

Theorem 3.13: Let $S(\neq \emptyset)$ be a non empty sub set of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$. Then the fuzzy set

$f_S: \mathcal{N} \rightarrow [0,1]$ defined by $f_S(x) = \begin{cases} 0 & \text{if } x \in S \\ 1 & \text{if } x \notin S \end{cases}$ is a co-fuzzy ideal of \mathcal{N} if and only if S is an ideal of \mathcal{N} .

Proof: Let $S(\neq \emptyset)$ be a non empty sub set of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Define $f_S: \mathcal{N} \rightarrow [0,1]$ such that $f_S(x) = \begin{cases} 0 & \text{if } x \in S \\ 1 & \text{if } x \notin S \end{cases}$

Let f_S be a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Now we prove that S is an ideal of \mathcal{N}

1. Let $x, y \in S$

$$\text{Then } f_S(x) = 0 \text{ and } f_S(y) = 0$$

$$\text{Now } f_S(x + y) \leq \text{Max}\{f_S(x), f_S(y)\} = 0$$

$$\therefore f_S(x + y) = 0$$

$$\Rightarrow x + y \in S$$

2. $x \in S$ and $y \in \mathcal{N}$
Then $f_S(x) = 0$
Now $f_S(y + x - y) \leq f_S(x) = 0$
 $\therefore f_S(y + x - y) = 0$
 $\Rightarrow y + x - y \in S$
3. Let $x \in S$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$
Then $f_S(x) = 0$
Now $f_S(a\alpha(x + b) - a\alpha b) \leq f_S(x) = 0$
 $\therefore f_S(a\alpha(x + b) - a\alpha b) = 0$
 $\Rightarrow a\alpha(x + b) - a\alpha b \in S$
Then S is a left ideal of \mathcal{N}
4. Let $x \in S$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$
Then $f_S(x) = 0$
We have $f_S(x\alpha a) \leq f_S(x) = 0$
 $\therefore f_S(x\alpha a) = 0$
 $\Rightarrow x\alpha a \in S$
Then S is a right ideal of \mathcal{N}
Hence S is an ideal of \mathcal{N}

Conversely assume that S is an ideal of \mathcal{N}

We prove that f_S is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

1. Let $x, y \in \mathcal{N}$

(i) Let $x, y \in S$

Then $f_S(x) = 0$ and $f_S(y) = 0$ and $x + y \in S$

$$\Rightarrow f_S(x + y) = 0 \leq \max\{f_S(x), f_S(y)\}$$

$$\therefore f_S(x + y) \leq \max\{f_S(x), f_S(y)\}$$

(ii) Let $x, y \notin S$ and $x + y \in S$

Then $f_S(x) = f_S(y) = 1$ and $f_S(x + y) = 0$

$$\text{Now } f_S(x + y) = 0 \leq \max\{f_S(x), f_S(y)\}$$

$$\therefore f_S(x + y) \leq \max\{f_S(x), f_S(y)\}$$

(iii) Let $x, y \notin S$ and $x + y \notin S$

Then $f_S(x) = f_S(y) = f_S(x + y) = 1$

$$\text{Now } f_S(x + y) = 1 \leq \max\{f_S(x), f_S(y)\}$$

$$\therefore f_S(x + y) \leq \max\{f_S(x), f_S(y)\}$$

2. Let $x, y \in \mathcal{N}$

(i) Let $x \in S$ and $y \in \mathcal{N}$

Then $f_S(x) = 0$ and $y + x - y \in S$

$$\Rightarrow f_S(y + x - y) = 0 \leq f_S(x)$$

$$\therefore f_S(y + x - y) \leq f_S(x)$$

(ii) Let $x \notin S$ and $y \in \mathcal{N}$

Then $f_S(x) = 1$

Now $f_S(y + x - y) \leq 1 \leq f_S(x)$

$$\therefore f_S(y + x - y) \leq f_S(x)$$

3. Let $x, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

(i) Let $x \in S$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f_S(x) = 0$ and $a\alpha(x + b) - a\alpha b \in S$

$$\Rightarrow f_S(a\alpha(x + b) - a\alpha b) = 0$$

Now $f_S(a\alpha(x + b) - a\alpha b) = 0 \leq f_S(x)$

$$\therefore f_S(a\alpha(x + b) - a\alpha b) \leq f_S(x)$$

(ii) Let $x \notin S$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f_S(x) = 1$

Now $f_S(a\alpha(x+b) - a\alpha b) \leq 1 = f_S(x)$

$\therefore f_S(a\alpha(x+b) - a\alpha b) \leq f_S(x)$

Thus f_S is a co-fuzzy left ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

4. Let $x, a \in \mathcal{N}$ and $\alpha \in \Gamma$

(i) Let $x \in S$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f_S(x) = 0$ and $x\alpha a \in S$

$\Rightarrow f_S(x\alpha a) = 0$

Now $f_S(x\alpha a) = 0 \leq f_S(x)$

$\therefore f_S(x\alpha a) \leq f_S(x)$

(ii) Let $x \notin S$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f_S(x) = 1$

Now $f_S(x\alpha a) \leq 1 \leq f_S(x)$

$\therefore f_S(x\alpha a) \leq f_S(x)$

Thus f_S is a co-fuzzy right ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Hence f_S is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Theorem 3.14: If f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ then the set $S = \{x \in \mathcal{N} / f(x) = f(0)\}$ is an ideal of \mathcal{N}

Proof: Let f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Given that $S = \{x \in \mathcal{N} / f(x) = f(0)\}$

We know that $f(0) \leq f(x)$, $\forall x \in \mathcal{N}$

Now we prove that S is an ideal of \mathcal{N}

1. Let $x, y \in S$

Then $f(x) = f(0)$ and $f(y) = f(0)$

Since f is a fuzzy ideal, $f(x+y) \leq \text{Max}\{f(x), f(y)\} = f(0)$

$f(x+y) \leq f(0)$

$\Rightarrow f(x+y) = f(0)$

$\Rightarrow x+y \in S$

2. Let $x \in S$ and $y \in \mathcal{N}$

Then $f(x) = f(0)$

Since f is a fuzzy ideal, $f(y+x-y) \leq f(x) = f(0)$

$\therefore f(y+x-y) \leq f(0)$

$\Rightarrow f(y+x-y) = f(0)$

$\Rightarrow y+x-y \in S$

3. Let $x \in S$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f(x) = f(0)$

Since f is a fuzzy left ideal, $f(a\alpha(x+b) - a\alpha b) \leq f(x) = f(0)$

$\therefore f(a\alpha(x+b) - a\alpha b) \leq f(0)$

$\Rightarrow f(a\alpha(x+b) - a\alpha b) = f(0)$

$\Rightarrow a\alpha(x+b) - a\alpha b \in S$

Thus S is a left ideal of \mathcal{N} .

4. Let $x \in S$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f(x) = f(0)$

Since f is a fuzzy right ideal, $f(x\alpha a) \leq f(x) = f(0)$

$\therefore f(x\alpha a) \leq f(0)$

$\Rightarrow f(x\alpha a) = f(0)$

$\Rightarrow x\alpha a \in S$

Thus S is a right ideal of \mathcal{N} .

Hence S is an ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

Theorem 3.15: Let $S (\neq \emptyset)$ is an ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$. Then for all $t \in (0, 1]$ there exists a co-fuzzy ideal f of \mathcal{N} Such that $\mathcal{N}_f = S$, Where $\mathcal{N}_f = \{x \in \mathcal{N} / f(x) = f(0)\}$.

Proof: Given that $S (\neq \phi)$ is an ideal of \mathcal{N} . Define $f: \mathcal{N} \rightarrow [0,1]$ such that $f(x) = \begin{cases} 0 & \text{if } x \in S \\ t & \text{if } x \notin S \end{cases}$

We prove that f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

1. Let $x, y \in \mathcal{N}$

(i) Let $x, y \in S$

Then $f(x) = f(y) = 0$

Since S is an ideal of \mathcal{N} , $x + y \in S$

$\Rightarrow f(x + y) = 0$

$\Rightarrow f(x + y) = 0 \leq \text{Max}\{f(x), f(y)\}$

$\therefore f(x + y) \leq \text{Max}\{f(x), f(y)\}$

(ii) Let $x, y \notin S$ and $x + y \in S$

Then $f(x) = f(y) = t$ and $f(x + y) = 0$

Now $f(x + y) = 0 \leq \text{Max}\{f(x), f(y)\}$

$\therefore f(x + y) \leq \text{Max}\{f(x), f(y)\}$

(iii) Let $x, y \notin S$ and $x + y \notin S$

Then $f(x) = f(y) = t$ and $f(x + y) = t$

Now $f(x + y) = t \leq \text{Max}\{f(x), f(y)\}$

$\therefore f(x + y) \leq \text{Max}\{f(x), f(y)\}$

2. Let $x, y \in \mathcal{N}$

(i) Let $x \in S$ and $y \in \mathcal{N}$

Then $f(x) = 0$

Since S is an ideal of \mathcal{N} , $y + x - y \in S$

$\Rightarrow f(y + x - y) = 0 \leq f(x)$

$\Rightarrow f(y + x - y) \leq f(x)$

(ii) Let $x \notin S$ and $y \in \mathcal{N}$

Then $f(x) = t$ and $y + x - y \notin S$

$\Rightarrow f(y + x - y) = t \leq f(x)$

$\Rightarrow f(y + x - y) \leq f(x)$

3. Let $x, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

(i) Let $x \in S$, $a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f(x) = 0$ and $a\alpha(x + b) - a\alpha b \in S$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) = 0 \leq f(x)$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) \leq f(x)$

(ii) Let $x \notin S$ and $a\alpha(x + b) - a\alpha b \in S$

Then $f(x) = t$ and $f(a\alpha(x + b) - a\alpha b) = 0$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) = 0 \leq t = f(x)$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) \leq f(x)$

(iii) Let $x \notin S$ and $a\alpha(x + b) - a\alpha b \notin S$

Then $f(x) = t$ and $f(a\alpha(x + b) - a\alpha b) = t$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) = t = f(x)$

$\Rightarrow f(a\alpha(x + b) - a\alpha b) \leq f(x)$

Thus f is a co-fuzzy left ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

4. Let $x, a \in \mathcal{N}$ and $\alpha \in \Gamma$

(i) Let $x \in S$, $a \in \mathcal{N}$ and $\alpha \in \Gamma$

Then $f(x) = 0$ and $x\alpha a \in S$

$\Rightarrow f(x\alpha a) = 0$

$\Rightarrow f(x\alpha a) = 0 \leq f(x)$

$\Rightarrow f(x\alpha a) \leq f(x)$

(ii) Let $x \notin S$ and $x\alpha a \in S$

Then $f(x) = t$ and $f(x\alpha a) = 0$

$\Rightarrow f(x\alpha a) = 0 \leq t = f(x)$

$\Rightarrow f(x\alpha a) \leq f(x)$

(iii) Let $x \notin S$ and $x\alpha a \notin S$

Then $f(x) = t$ and $f(x\alpha a) = t$

$\Rightarrow f(x\alpha a) = t = f(x)$

$\Rightarrow f(x\alpha a) \leq f(x)$

Thus f is a co-fuzzy right ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

Hence f is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$.

$$\begin{aligned}\text{Finally, } \mathcal{N}_f &= \{x \in \mathcal{N} / f(x) = f(0)\} \\ &= \{x \in \mathcal{N} / f(x) = 0\} \\ &= S \\ \therefore \mathcal{N}_f &= S\end{aligned}$$

Theorem 3.16: If f and g are two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$, then their intersection $(f \cap g)$ is also co-fuzzy ideal of \mathcal{N} .

Proof: Given that f and g are two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

Now we prove that $(f \cap g)$ is co-fuzzy ideal of \mathcal{N} .

Let $x, y, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

1. $(f \cap g)(x + y) = \text{Min} \{f(x + y), g(x + y)\}$
 $\leq \text{Min} \{\text{Max} \{f(x), f(y)\}, \text{Max} \{g(x), g(y)\}\}$
 $\leq \text{Max} \{\text{Min} \{f(x), g(x)\}, \text{Min} \{f(y), g(y)\}\}$
 $= \text{Max} \{(f \cap g)(x), (f \cap g)(y)\}$
 $\therefore (f \cap g)(x + y) \leq \text{Max} \{(f \cap g)(x), (f \cap g)(y)\}$
2. $(f \cap g)(y + x - y) = \text{Min} \{f(y + x - y), g(y + x - y)\}$
 $\leq \text{Min} \{f(x), g(x)\}$
 $= (f \cap g)(x)$
 $\therefore (f \cap g)(y + x - y) \leq (f \cap g)(x)$
3. $(f \cap g)(a\alpha(x + b) - a\alpha b) = \text{Min} \{f(a\alpha(x + b) - a\alpha b), g(a\alpha(x + b) - a\alpha b)\}$
 $\leq \text{Min} \{f(x), g(x)\}$
 $= (f \cap g)(x)$
 $\therefore (f \cap g)(a\alpha(x + b) - a\alpha b) \leq (f \cap g)(x)$
 Thus $(f \cap g)$ is a co-fuzzy left ideal of \mathcal{N}
4. $(f \cap g)(x\alpha a) = \text{Min} \{f(x\alpha a), g(x\alpha a)\}$
 $\leq \text{Min} \{f(x), g(x)\}$
 $= (f \cap g)(x)$
 $\therefore (f \cap g)(x\alpha a) \leq (f \cap g)(x)$
 Thus $(f \cap g)$ is a co-fuzzy right ideal of \mathcal{N} .
 Hence $(f \cap g)$ is a co-fuzzy ideal of \mathcal{N} .

Remark 3.17: The union of two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$

need not be a co-fuzzy ideal of \mathcal{N} . Now, we prove a sufficient condition for the union of two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ to be a co-fuzzy ideal of \mathcal{N} .

Theorem 3.18: If f and g are two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$, then their union $(f \cup g)$ is fuzzy ideal of \mathcal{N} if $f \subseteq g$ or $g \subseteq f$.

Proof: Given that f and g are two co-fuzzy ideals of a finite Γ -near ring $(\mathcal{N}, +, \Gamma)$ such that $f \subseteq g$ or $g \subseteq f$,

Without loss of generality, We assume that $f \subseteq g$

Now we prove that $(f \cup g)$ is co-fuzzy ideal of \mathcal{N} .

Let $x, y, a, b \in \mathcal{N}$ and $\alpha \in \Gamma$

1. $(f \cup g)(x + y) = \text{Max} \{f(x + y), g(x + y)\}$
 $\leq \text{Max} \{\text{Max} \{f(x), f(y)\}, \text{Max} \{g(x), g(y)\}\}$
 $\leq \text{Max} \{\text{Max} \{f(x), g(x)\}, \text{Max} \{f(y), g(y)\}\}$
 $= \text{Max} \{(f \cup g)(x), (f \cup g)(y)\}$
 $\therefore (f \cup g)(x + y) \leq \text{Max} \{(f \cup g)(x), (f \cup g)(y)\}$
2. $(f \cup g)(y + x - y) = \text{Max} \{f(y + x - y), g(y + x - y)\}$
 $\leq \text{Max} \{f(x), g(x)\}$
 $= (f \cup g)(x)$
 $\therefore (f \cup g)(y + x - y) \leq (f \cup g)(x)$
3. $(f \cup g)(a\alpha(x + b) - a\alpha b) = \text{Max} \{f(a\alpha(x + b) - a\alpha b), g(a\alpha(x + b) - a\alpha b)\}$
 $\leq \text{Max} \{f(x), g(x)\}$
 $= (f \cup g)(x)$
 $\therefore (f \cup g)(a\alpha(x + b) - a\alpha b) \leq (f \cup g)(x)$
 Thus $(f \cup g)$ is a co-fuzzy left ideal of \mathcal{N}

4. $(f \cup g)(x\alpha a) = \text{Max} \{f(x\alpha a), g(x\alpha a)\}$
 $\leq \text{Max} \{f(x), g(x)\}$
 $= (f \cup g)(x)$
 $\therefore (f \cup g)(x\alpha a) \leq (f \cup g)(x)$
 Thus $(f \cup g)$ is a co-fuzzy right ideal of \mathcal{N}
 Hence $(f \cup g)$ is a co-fuzzy ideal of \mathcal{N}

Remark 3.19: The converse of the above theorem need not be true. i.e, even if $(f \cup g)$ is a co-fuzzy ideal of a finite Γ -near ring, either one may not contained in other.

Example: Let $S = \{1,2,3,4\}$ and $\mathcal{N} = P(S)$, Where $P(S)$ is the power set of S .

Then $(\mathcal{N}, \Delta, \cap)$ is a finite near ring.

Let us take $\Gamma = \{\{1\}, \{1,2,3\}\}$, then $(\mathcal{N}, \Delta, \Gamma)$ is a finite Γ -near ring.

Define $f: \mathcal{N} \rightarrow [0,1]$ such that $f(A) = \begin{cases} 0.6 & \text{if } A \neq \phi \\ 0.3 & \text{if } A = \phi \end{cases}$

and define $g: \mathcal{N} \rightarrow [0,1]$ such that $g(A) = \begin{cases} 0.7 & \text{if } A \neq \phi \\ 0.2 & \text{if } A = \phi \end{cases}$

Then $(f \cup g): \mathcal{N} \rightarrow [0,1]$ such that $(f \cup g)(A) = \begin{cases} 0.7 & \text{if } A \neq \phi \\ 0.3 & \text{if } A = \phi \end{cases}$ is a co-fuzzy ideal of a finite Γ -near ring $(\mathcal{N}, \Delta, \Gamma)$. But $f \not\subseteq g$ and $g \not\subseteq f$.

4. CO-FUZZY HOMOMORPHISM OF FINITE Γ -NEAR RINGS

In this section, we define co-fuzzy homomorphism between two finite Γ -near rings and we prove that the homomorphic image of co-fuzzy ideal is a co-fuzzy ideal and inverse image of a co-fuzzy ideal is also a co-fuzzy ideal.

Definition 4.1: Let $(\mathcal{N}_1, +, \Gamma)$ and $(\mathcal{N}_2, +, \Gamma)$ be two finite Γ -near rings. Then the function $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is said to be a homomorphism from \mathcal{N}_1 to \mathcal{N}_2 if

1. $f(a + b) = f(a) + f(b)$
2. $f(a\alpha b) = f(a)\alpha f(b), \forall a, b \in \mathcal{N}_1$ and $\alpha \in \Gamma$

Definition 4.2: Let $(\mathcal{N}_1, +, \Gamma_1)$ be a finite Γ_1 -near ring and $(\mathcal{N}_2, +, \Gamma_2)$ be a finite Γ_2 -near ring, and $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ and $g: \Gamma_1 \rightarrow \Gamma_2$ be two functions. Then the pair (f, g) is said to be a homomorphism from \mathcal{N}_1 to \mathcal{N}_2 if

1. $f(a + b) = f(a) + f(b)$
2. $f(a\alpha b) = f(a)g(\alpha)f(b), \forall a, b \in \mathcal{N}_1$ and $\alpha \in \Gamma_1$

Definition 4.3: Let $(\mathcal{N}_1, +, \Gamma)$ and $(\mathcal{N}_2, +, \Gamma)$ be two finite Γ -near rings and $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is a homomorphism from \mathcal{N}_1 to \mathcal{N}_2 . If μ is a fuzzy sub set of \mathcal{N}_1 then its image $f(\mu)$ is a fuzzy sub set of \mathcal{N}_2 is defined by

$$(f(u))(y) = \begin{cases} \inf_{z \in f^{-1}(y)} \mu(z) & \text{if } f^{-1}(y) \neq \phi \\ 0 & \text{if } f^{-1}(y) = \phi \end{cases} \quad \forall y \in \mathcal{N}_2$$

Definition 4.4: Let $(\mathcal{N}_1, +, \Gamma)$ and $(\mathcal{N}_2, +, \Gamma)$ be two finite Γ -near rings. The function $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is a homomorphism from \mathcal{N}_1 to \mathcal{N}_2 . If σ is a co-fuzzy sub set of \mathcal{N}_2 then its inverse image $f^{-1}(\sigma)$ is a fuzzy sub set of \mathcal{N}_1 defined by $(f^{-1}(\sigma))(x) = \sigma(f(x)), \forall x \in \mathcal{N}_1$.

Theorem 4.5: Let $(\mathcal{N}_1, +, \Gamma)$ and $(\mathcal{N}_2, +, \Gamma)$ be two finite Γ -near rings. The function $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is a homomorphism from \mathcal{N}_1 to \mathcal{N}_2 . If σ is a co-fuzzy ideal of \mathcal{N}_2 then its inverse image $f^{-1}(\sigma)$ is a co-fuzzy ideal of \mathcal{N}_1 .

Proof: Let $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is a homomorphism from the finite Γ -near rings $(\mathcal{N}_1, +, \Gamma)$ to $(\mathcal{N}_2, +, \Gamma)$ and $\sigma: \mathcal{N}_2 \rightarrow [0,1]$ is a co-fuzzy sub set of \mathcal{N}_2 .

Let σ be a co-fuzzy ideal of \mathcal{N}_2

Now we prove that $f^{-1}(\sigma)$ is a co-fuzzy ideal of \mathcal{N}_1

Let $x, y, a, b \in \mathcal{N}_1$ and $\alpha \in \Gamma$

1. $(f^{-1}(\sigma))(x + y) = \sigma(f(x + y))$
 $= \sigma(f(x) + f(y))$
 $\leq \text{Max} \{ \sigma(f(x)), \sigma(f(y)) \}$
 $= \text{Max} \{ (f^{-1}(\sigma))(x), (f^{-1}(\sigma))(y) \}$
 $\therefore (f^{-1}(\sigma))(x + y) \leq \text{Max} \{ (f^{-1}(\sigma))(x), (f^{-1}(\sigma))(y) \}$
2. $(f^{-1}(\sigma))(y + x - y) = \sigma(f(y + x - y))$
 $= \sigma(f(y) + f(x) - f(y))$
 $\leq \sigma(f(x))$
 $= (f^{-1}(\sigma))(x)$
 $\therefore (f^{-1}(\sigma))(y + x - y) \leq (f^{-1}(\sigma))(x)$
3. $(f^{-1}(\sigma))(a\alpha(x + b) - a\alpha b) = \sigma(f(a\alpha(x + b) - a\alpha b))$
 $= \sigma(f(a)\alpha(f(x) + f(b)) - f(a)\alpha f(b))$
 $\leq \sigma(f(x))$
 $= (f^{-1}(\sigma))(x)$
 $\therefore (f^{-1}(\sigma))(a\alpha(x + b) - a\alpha b) \leq (f^{-1}(\sigma))(x)$

Thus $(f^{-1}(\sigma))$ is a co-fuzzy left ideal of \mathcal{N}_1

4. $(f^{-1}(\sigma))(x\alpha a) = \sigma(f(x\alpha a))$
 $= \sigma(f(x)\alpha f(a))$
 $\leq \sigma(f(x))$
 $= (f^{-1}(\sigma))(x)$

$\therefore (f^{-1}(\sigma))(x\alpha a) \leq (f^{-1}(\sigma))(x)$

Thus $(f^{-1}(\sigma))$ is a co-fuzzy right ideal of \mathcal{N}_1

Hence $(f^{-1}(\sigma))$ is a co-fuzzy ideal of \mathcal{N}_1 .

Theorem 4.6: Let $(\mathcal{N}_1, +, \Gamma)$ and $(\mathcal{N}_2, +, \Gamma)$ be two finite Γ -near rings. The function $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is a onto homomorphism from \mathcal{N}_1 to \mathcal{N}_2 . If μ is a co-fuzzy ideal of \mathcal{N}_1 then its image $f(\mu)$ is a co-fuzzy ideal of \mathcal{N}_2 .

Proof: Let $f: \mathcal{N}_1 \rightarrow \mathcal{N}_2$ is an onto homomorphism from the finite Γ -near rings $(\mathcal{N}_1, +, \Gamma)$ to $(\mathcal{N}_2, +, \Gamma)$ and $\mu: \mathcal{N}_1 \rightarrow [0,1]$ is a fuzzy sub set of \mathcal{N}_1

Let μ be a co-fuzzy ideal of \mathcal{N}_1

Now we prove that $f(\mu)$ is a co-fuzzy ideal of \mathcal{N}_2

We have

$$(f(\mu))(y) = \begin{cases} \inf_{z \in f^{-1}(y)} \mu(z) & \text{if } f^{-1}(y) \neq \emptyset \\ 0 & \text{if } f^{-1}(y) = \emptyset \end{cases} \quad \forall y \in \mathcal{N}_2$$

Let $x, y, a, b \in \mathcal{N}_2$ and $\alpha \in \Gamma$

Since f is onto from \mathcal{N}_1 to \mathcal{N}_2 then there exists $x_0, y_0, a_0, b_0 \in \mathcal{N}_1$ such that

$f(x_0) = x, f(y_0) = y, f(a_0) = a$ and $f(b_0) = b$

$$1. (f(\mu))(x+y) = \inf_{z \in f^{-1}(x+y)} \mu(z) \quad (1)$$

$$\begin{aligned} \text{Let } \mu(x_0) &= \inf_{z \in f^{-1}(x)} \mu(z) \text{ and } \mu(y_0) = \inf_{z \in f^{-1}(y)} \mu(z) \\ \Rightarrow f(x_0) &= x \text{ and } f(y_0) = y \\ \text{Now } f(x_0 + y_0) &= f(x_0) + f(y_0) = x + y \\ \Rightarrow x_0 + y_0 &\in f^{-1}(x+y) \end{aligned}$$

$$\begin{aligned} \text{From equation (1), } (f(\mu))(x+y) &= \inf_{z \in f^{-1}(x+y)} \mu(z) \\ &\leq \mu(x_0 + y_0) \\ &\leq \max\{\mu(x_0), \mu(y_0)\} \\ &= \max\{\inf_{z \in f^{-1}(x)} \mu(z), \inf_{z \in f^{-1}(y)} \mu(z)\} \\ &= \max\{(f(\mu))(x), (f(\mu))(y)\} \\ \therefore (f(\mu))(x+y) &\leq \max\{(f(\mu))(x), (f(\mu))(y)\} \end{aligned}$$

$$2. (f(\mu))(y+x-y) = \inf_{z \in f^{-1}(y+x-y)} \mu(z) \quad (2)$$

$$\begin{aligned} \text{Now } f(y_0 + x_0 - y_0) &= f(y_0) + f(x_0) - f(y_0) = y + x - y \\ \therefore y_0 + x_0 - y_0 &\in f^{-1}(y+x-y) \end{aligned}$$

$$\begin{aligned} \text{From equation (2), } (f(\mu))(y+x-y) &= \inf_{z \in f^{-1}(y+x-y)} \mu(z) \\ &\leq \mu(y_0 + x_0 - y_0) \\ &\leq \mu(x_0) \\ &= \inf_{z \in f^{-1}(x)} \mu(z) \\ &= (f(\mu))(x) \\ \therefore (f(\mu))(y+x-y) &\leq (f(\mu))(x) \end{aligned}$$

$$3. (f(\mu))(a\alpha(x+b) - a\alpha b) = \inf_{z \in f^{-1}(a\alpha(x+b) - a\alpha b)} \mu(z) \quad (3)$$

$$\begin{aligned} \text{Now } f(a_0\alpha(x_0 + b_0) - a_0\alpha b_0) &= f(a_0)\alpha(f(x_0) + f(b_0)) - f(a_0)\alpha f(b_0) \\ &= a\alpha(x+b) - a\alpha b \\ \therefore a_0\alpha(x_0 + b_0) - a_0\alpha b_0 &\in f^{-1}(a\alpha(x+b) - a\alpha b) \end{aligned}$$

$$\begin{aligned} \text{From equation (3), } (f(\mu))(a\alpha(x+b) - a\alpha b) &= \inf_{z \in f^{-1}(a\alpha(x+b) - a\alpha b)} \mu(z) \\ &\leq \mu(a_0\alpha(x_0 + b_0) - a_0\alpha b_0) \\ &\leq \mu(x_0) \\ &= \inf_{z \in f^{-1}(x)} \mu(z) \\ &= (f(\mu))(x) \\ \therefore (f(\mu))(a\alpha(x+b) - a\alpha b) &\leq (f(\mu))(x) \end{aligned}$$

Thus $(f(\mu)): \mathcal{N}_2 \rightarrow [0,1]$ is a co-fuzzy left ideal of \mathcal{N}_2

$$4. (f(\mu))(x\alpha a) = \inf_{z \in f^{-1}(x\alpha a)} \mu(z) \quad (4)$$

$$\begin{aligned} \text{Now } f(x_0\alpha a_0) &= f(x_0)\alpha f(a_0) = x\alpha a \\ \therefore x_0\alpha a_0 &\in f^{-1}(x\alpha a) \end{aligned}$$

$$\begin{aligned} \text{From equation (4), } (f(\mu))(x\alpha a) &= \inf_{z \in f^{-1}(x\alpha a)} \mu(z) \\ &\leq \mu(x_0\alpha a_0) \\ &\leq \mu(x_0) \\ &= \inf_{z \in f^{-1}(x)} \mu(z) \\ &= (f(\mu))(x) \\ \therefore (f(\mu))(x\alpha a) &\leq (f(\mu))(x) \end{aligned}$$

Then $(f(\mu)): \mathcal{N}_2 \rightarrow [0,1]$ is a co-fuzzy right ideal of \mathcal{N}_2

Hence $(f(\mu)): \mathcal{N}_2 \rightarrow [0,1]$ is a co-fuzzy ideal of \mathcal{N}_2 .

5. CONCLUSION

In this article, we inspected the idea of co-fuzzy ideals of a finite Γ -near ring. We proved some necessary and sufficient conditions for a fuzzy subset of finite Γ -near ring to be co-fuzzy ideal of the ring. The intersection and union of co-fuzzy ideals and homomorphism theorems have been proved. This concept may be extended to Bipolar co-fuzzy ideals in finite Γ -near rings.

6. REFERENCES

1. Dutta, T.K., Sardar. S.K. and Sarbani Goswami,(2011) Fuzzy Ideals Extension in Γ – semirings, arXiv : 1104.4791v1 – [math. GM].
2. Jun. Y.B.Neggars.J. and Kim.H.S.,(1996) Normal L – Fuzzy ideals in semirings. Fuzzy Sets and Systems. Vol.82, 383 – 386.
3. Jun.Y.B.. On fuzzy prime ideals of G-rings. Soochow J. Math, 21(1). (1995). 41-48.
4. Jun.Y.B., Lee. C.Y., Fuzzy prime ideals in G-rings. Pusan Kyongnam Math J., 9(1), (1993), 105-111.
5. Magil.K.D.Jr. and S.subbiah(1975) Maximal ideals of semigroups of endomorphisms, Bull. Austral,Math. Soc. Vol.12, 211-225.
6. Malik. D.S and Mordeson N.John. (1990) Fuzzy prime ideals of a ring. Fuzzy Sets and Systems. Vol.37, 93-98.
7. Rosen field.A., Fuzzy groups, J. Math, Anal.Appl.35.512–517(1971).
8. Swamy, K.L.N and Swamy, U.M..(1988) Fuzzy Prime Ideals of Rings, Journal of Mathematical Analysis and Applications. Vol.134, No.1. 94–103.
9. Satyanarayana Bh., A note on G–near–rings, Indian J.Math., 41(3).(1999). 427-133.
10. Steinfeld, O., (1978) Quasi-ideals in rings and semigroups. Akademiai kiado, Budapest.
11. Yaqoob.N., Bipolar – Valued Fuzzy Ideals in LA – semigroups. Journal of Advanced Studies in Topology. 3(2012). 60–71.
12. Zadeh L.A., (1965) Fuzzy sets. Information and Control Vol.8. 338–353.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2021. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]