SOME FIXED POINT THEOREMS FOR δ -COMPATIBLE MAPPINGS IN METRIC SPACES

¹S. De Sarkar* and ²Kalishankar Tiwary

¹Department of Mathematics, University of North Bengal, Rajarammohanpur, Siliguri, (Darjeeling), West Bengal E-mail: suparnadesarkar@yahoo.co.in

²Department of Mathematics, City College South (Evening), 23/49 Gariahat Road, Kolkata 700 029, West Bengal E-mail: tiwarykalishankar@yahoo.co.in

(Received on: 14-11-11; Accepted on: 30-11-11)

ABSTRACT

S ome common fixed point theorems for point-valued and set-valued compatible mappings in metric spaces are proved.

Key words: Fixed points, δ -compatible mappings.

AMS Subject Classification [2010]: 47H10.

1. INTRODUCTION:

In the last two decades research workers have generalized/extended Banach contraction theorem in various ways. A large number of research papers are available which have generalized the theorem of Banach involving more than one mapping. In these papers they have used the notion of commutativity, weak commutativity and compatibility. For this one can see the papers [3, 4, 5, 6, 8, 9]. It is shown by examples in [3] and [10] that commuting mappings are weakly commuting and weakly commuting mappings are compatible. The converse of these results is not true. Some authors have also made attempts to prove some fixed point theorems for set-valued mappings. Jungck and Rhoades have generalized some results of [1], [2] via compatibility for set-valued mappings.

In this paper we have proved some common fixed point theorems for point valued and set-valued compatible mappings which coincide with the results already proved for fixed point theorems for point-valued mappings.

2. KNOWN DEFINITIONS AND RESULTS:

Let (X,d) be a metric space and let $f,g:(X,d) \rightarrow (X,d)$. The mappings f and g are said to be weakly commutative [1] if and only if $d(fgx, gfx) \leq d(fx, gx)$ for $x \in X$. The mappings f and g are said to be compatible [cf. 6] if and only if whenever $\{x_n\}$ is a sequence in X such that fx_n , $gxn \rightarrow t$ then $d(fgx_n, gfxn) \rightarrow 0$.

Let B(X) denote the family of bounded subsets of X and let $\delta:B(X)\times B(X)\to [0,\infty)$ by $\delta(A,B)=sup\{d(a,b):a\in A,b\in B\}$. Fisher $\{[1], [2]\}$ made several discussions on δ and obtained some of its properties. It is to be note that for any three $A,B,C\in B(X), \delta(A,B)\leq \delta(A,C)+\delta(C,B)$ and $\delta(A,B)=0$ iff $A=B=\{a\}$. If $x\in X$, we write $\delta(x,A)$ for $\delta(\{x\},A)$.

Let $\{A_n\}$ be a sequence in B(X), we say that $\{A_n\} \rightarrow A \subset X$, and we write $A_n \rightarrow A$ if and only if

(i) $a \in A$ implies that $a = lima_n$ for some sequence $\{a_n\}$ in X where $a_n \in A_n$ for n = 1, 2, 3, ... and

(ii) for any $\epsilon > 0$ there exists $m \epsilon N$, N is a set of natural numbers, such that $A_n \epsilon A_{\epsilon} = \{x \epsilon A : d(x, a) < \epsilon \text{ for some } a \epsilon A \text{ for } n > m\}$.

Corresponding author: ¹S. De Sarkar, *E-mail: suparnadesarkar@yahoo.co.in

¹S. De Sarkar* and ²Kalishankar Tiwary/ SOME FIXED POINT THEOREMS FOR δ-COMPATIBLE MAPPINGS IN METRIC SPACES/ IJMA- 2(12), Dec.-2011, Page: 2813-2817

The following lemmas are collected from Fisher $\{[1], [2]\}$ which we need to prove our theorem.

Lemma: 1 Suppose $\{A_n\}$ and $\{B_n\}$ are sequences in B(X) and (X, d) is a complete metric space. If $A_n \rightarrow A \in B(X)$ and $B_n \rightarrow B \in B(X)$ then $\delta(A_n, B_n) \rightarrow \delta(A, B)$.

Lemma: 2 If $\{A_n\}$ is a sequence of non-empty bounded sets in the complete metric space (X, d) and if $\lim_{n\to\infty} \delta(A_n, \{y\}) = 0$ for some $y \in X$ then $A_n \to \{y\}$.

Let $F: X \rightarrow B(X)$. *F* is said to be continuous at $x \in X$ if $x_n \rightarrow x$ implies $Fx_n \rightarrow Fx$. Jungck and Rhoades [6] formulated the notion of compatibility for point valued and set-valued mappings in the following way:

Let *I*: $X \to X$ and *F*: $X \to B(X)$. *F* and *I* are δ -compatible if and only if $IF(x) \in B(X)$ for $x \in X$ and δ (*IFx*, *FIxn*) $\to 0$ whenever $\{x_n\}$ is a sequence in *X* such that $Ix_n \to t$ and $Fx_n \to \{t\}$ for some $t \in X$.

Further Jungck and Rhoades [6] proved the following proposition.

Proposition: 1 Let (X,d) be a complete metric space. Suppose I:X \rightarrow X,F:X \rightarrow B(X) and I and F are δ -compatible. (i) Suppose the sequence {*Fx_n*} converges to {*z*} and {*Ix_n*} converges to *z*. If *I* is continuous, then *FIx_n\rightarrow*{*Iz*}. (ii) If {*Iu*}=*Fu* for some *u* \in X, then *FIu*=*IFu*.

By $\bigcup FX$ we mean $\bigcup F(X) = \{y \in X : y \in Fx \text{ for some } x \in X\}.$

3. THEOREMS AND COROLLARIES:

Theorem: 1 Let F,G be two set-valued mappings from a complete metric space X into B(X). Let I, J be mappings from X into itself satisfying the inequality

$$\delta(Fx,Gy) \leq \max\{\delta(Ix,Jy), \delta(Ix,Fx), \delta(Jy,Gy)\} - w[\max\{\delta(Ix,Jy), \delta(Ix,Fx), \delta(Jy,Gy)\}]$$

for all $x, y \in X$. Suppose the mappings F and I are δ -compatible and G and J are δ -compatible and $UGX \subset I(X)$ and $UFX \subset J(X)$. If one of the mappings I and J is continuous then F, G, I and J have a unique common fixed point z. Moreover Moreover $Fz=Gz=\{z\}$ and $w: [0, \infty) \rightarrow [0, \infty)$ such that w is continuous and w(r) < r for r > 0.

Proof: Let $x_0 \in X$ be arbitrary. Since J contains the range of F and I contains the range of G, we construct a sequence $\{x_n\}$ inductively in X such that

 $Jx_{2n+1} \in Y_{2n} = Fx_{2n}, Ix_{2n+2} \in Y_{2n+1} \in Gx_{2n+1}, n=0,1,2,...$

Let

$$\delta_n = \delta(Fx_n, Gx_{n+1})$$
 if $n = 2m, m = 0, 1, 2, ...$

$$= \delta(Gx_n, Fx_{n+1})$$
 if $n=2m+1$.

Then

$$\delta_{2m} = \delta(Fx_{2m}, Gx_{2m+1}) \le \max\{d(Ix_{2m}, Jx_{2m+1}), \delta(Ix_{2m}, Fx_{2m}), \delta(Jx_{2m+1}, Gx_{2m+1})\} - w[\max\{d(Ix_{2m}, Jx_{2m+1}), \delta(Ix_{2m}, Fx_{2m}), \delta(Jx_{2m+1}, Gx_{2m+1})]$$

i.e.
$$\delta_{2m} \le max \{ \delta (Gx_{2m-1}, Fx^{2m}), \delta (Gx_{2m-1}, Fx^{2m}), \delta (Fx_{2m}, Gx^{2m+1}) \} \\ -w[max \{ \delta (Gx_{2m-1}, Fx_{2m}), \delta (Gx_{2m-1}, Fx_{2m}), \delta (Fx_{2m}, Gx_{2m+1}) \}$$

$$= max \{\delta_{2m-1}, \delta_{2m+1}, \delta_{2m}\} - w[max\{\delta_{2m-1}, \delta_{2m+1}, \delta_{2m}\}].$$

If $\delta_{2m} > \delta_{2m-1}$, then $\delta_{2m} \le \delta_{2m} - w \delta_{2m}$, a contradiction. So, $\delta_{2m} \le \delta_{2m-1}$. Then $\delta_{2m} \le \delta_{2m-1} - w (\delta_{2m-1})$. So, we have

$$\delta_n \le \delta_{n-l} - w(\delta_{n-1}) \tag{1}$$

i.e.
$$w(\delta_{n-1}) \leq \delta_{n-1} - \delta_n$$
.

So,
$$\sum_{i=0}^{n} w(\delta_i) \le \delta_0 - \delta_1 \le \delta_0.$$

1 S. De Sarkar * and 2 Kalishankar Tiwary/ SOME FIXED POINT THEOREMS FOR δ -COMPATIBLE MAPPINGS IN METRIC SPACES/ IJMA- 2(12), Dec.-2011, Page: 2813-2817

So, the series of non-negative terms $\sum_{i=0}^{n} w(\delta_i)$ is convergent. Hence, $\lim_{n\to\infty} w(\delta_n) = 0$. From (1) for $n \ge l$,

we get $\{\delta_n\}$ is a decreasing sequence of non-negative terms which implies that $\lim_{n\to\infty} w\delta_n = a$ for some $a \in \mathbb{R}^+$. Since w is continuous $w(a) = \lim_{n \to \infty} w(\delta_n) = 0 = 0$. This implies that a = 0. Hence,

$$\lim_{n \to \infty} \delta_n = 0. \tag{2}$$

We next show that the sequence $\{Fx_0=Y_0, Gx_1=Y_1, Fx_2=Y_2, Gx_3=Y_3, ...\}$ is a Cauchy sequence. In order to show that $\{Y_n\}$ is a Cauchy sequence, it is sufficient to show that $\{Y_{2n}\}$ is a Cauchy sequence. For this, there exists a positive number ε such that for each even integer 2k there are even positive integer 2m(k) and 2n(k) such that 2m(k) > 2n(k) > 2k and $\delta(Y_{2m(k)}, Y_{2n(k)}) < \varepsilon$.

For each even integer 2k, let 2m(k) be the least even integer exceeding 2n(k) satisfying the above inequality so that

$$\delta(Y_{2m(k)-2}, Y_{2n(k)}) \leq \epsilon \text{ and } \delta(Y_{2m(k)}, Y_{2n(k)}) > \epsilon$$

Then for each even integer 2k,

$$\begin{split} \delta(Y_{2m(k)}, Y_{2n(k)}) &\leq \delta(Y_{2n(k)}, Y_{2m(k)-2}) + \delta(Y_{2m(k)-2}, Y_{2m(k)-1}) + \delta(Y_{2m(k)-1}, Y_{2m(k)}) \\ &= \delta(Y_{2m(k)-2}, Y_{2n(k)}) + \delta_{2m(k)-2} + \delta_{2m(k)-1}. \end{split}$$

From (2), (3) and the above inequality

$$\lim_{n \to \infty} \delta(Y_{2m(k)}, Y_{2(k)}) = \epsilon \tag{4}$$

On using triangle inequality we get,

$$\begin{split} &|\delta(Y_{2m(k)},Y_{2n(k)+1}) - \delta(Y_{2m(k)},Y_{2n(k)})| \le \delta(Y_{2n(k)},Y_{2n(k)+1}) = \delta_{2n(k)}, \\ &|\delta(Y_{2m(k)+1},Y_{2n(k)+1}) - \delta(Y_{2m(k)},Y_{2n(k)+1})| \le \delta(Y_{2m(k)},Y_{2n(k)+1}) = \delta_{2m(k)}, \\ &|\delta(Y_{2m(k)+1},Y_{2n(k)+2}) - \delta(Y_{2m(k)+1},Y_{2n(k)+1})| \le \delta(Y_{2n(k)+1},Y_{2n(k)+2}) = \delta_{2n(k)+1}. \end{split}$$

From (2), (4) and the above inequality passing on limit as $k \rightarrow \infty$ we get,

* *

1.0 / 7.7

$$\epsilon = \lim_{k \to \infty} \delta(Y_{2m(k), Y_{2n(k)+1}}) = \lim_{k \to \infty} \delta(Y_{2m(k)+1, Y_{2n(k)+1}}) = \lim_{k \to \infty} \delta(Y_{2m(k)+1, Y_{2n(k)+2}}).$$
(A)

Now.

 $\delta(Y_{2m(k)+1}, Y_{2n(k)+2}) = \delta(Gx_{2m(k)+1}, Fx_{2n(k)+2})$

 $\leq max\{ \delta(Ix_{2n(k)+2}, Jx_{2m(k)+1}), \delta(Ix_{2n(k)+2}, Fx_{2n(k)+2}), \delta(Jx_{2m(k)+1}, Gx_{2m(k)+1})\}$ $-w[max\{\delta(Ix_{2n(k)+2},Jx_{2m(k)+1}),\delta(Ix_{2n(k)+2},Fx_{2n(k)+2}),\delta(Jx_{2m(k)+1},Gx_{2m(k)+1})\}]$

 $\leq max\{ \delta(Gx_{2n(k)+1}, Fx_{2m(k)}), \delta(Gx_{2n(k)+1}, Fx_{2n(k)+2}), \delta(Fx_{2m(k)}, Gx_{2n(k)+1}) \}$ - $W[max\{\delta(Gx_{2n(k)+1}, Fx_{2m(k)}), \delta(Gx_{2n(k)+1}, Fx_{2n(k)}), \delta(Fx_{2m(k)}, Gx_{2n(k)+1})\}]$

 $= max\{ \delta(Y_{2m(k)}, Y_{2n(k)+1}), \delta(Y_{2n(k)+1}, Y_{2n(k)+2}), \delta(Y_{2m(k)}, Y_{2m(k)+1}) \}$ $-w[max\{\delta(Y_{2m(k)},Y_{2n(k)+1}),\delta(Y_{2n(k)+1},Y_{2n(k)+2}),\delta(Y_{2m(k)+1},Y_{2m(k)+1})\}].$

Using (A) and passing on limit as $k \rightarrow \infty$, we have

 $\varepsilon \leq \varepsilon - w(\varepsilon)$ i.e. $w(\varepsilon) \leq 0$ a contradiction and so $\{Y_n\}$ is a Cauchy sequence.

If y_n is an arbitrary point in Y_n for n=1,2,..., it follows that $d(y_m,y_n) \leq \delta(Y_m,Y_n) < \varepsilon$ for m,n>N. It follows that the sequence $\{Y_n\}$ is a Cauchy sequence in the complete metric space X, the point y_n being independent of the particular choice of Yn and $\{y_n\}$ is Cauchy and because X is a complete metric space let y_n converge to $y \in X$. In particular the sequence $\{Ix_{2n}\}$ and $\{Jx_{2n+1}\}$ will converge to y and further by Fisher [2] the sequence of sets $\{Fx_{2n}\}$ and $\{Gx_{2n+1}\}$ will converge to the set $\{y\}$.

(3)

¹S. De Sarkar* and ²Kalishankar Tiwary/ SOME FIXED POINT THEOREMS FOR δ-COMPATIBLE MAPPINGS IN METRIC SPACES/ IJMA- 2(12), Dec.-2011, Page: 2813-2817

We now suppose that the mapping *I* is continuous. Then it follows that the sequence $I^2x_{2n} \rightarrow Iy$. Since *F* and *I* are δ -compatible, by proposition 1 [7], $FIx_n \rightarrow \{Iy\}$.

Now,

$$\delta(FIx_{2n}, Gx_{2n+1}) \le max\{ \delta(I^{2}x_{2n}, Jx_{2n+1}), \delta(I^{2}x_{2n}, FIx_{2n}), \delta(Jx_{2n+1}, Gx_{2n+1})\} \\ -w[max\{ \delta(I^{2}x_{2n}, Jx_{2n+1}), \delta(I^{2}x_{2n}, FIx_{2n}), \delta(Jx_{2n+1}, Gx_{2n+1})\}].$$

Letting *n* tend to infinity we have

$$d(Iy,y) \le max\{d(Iy,y), d(Iy,Iy), d(y,y)\} - w[max\{dI(Iy,y), d(Iy,Iy)\}] = d(Iy,y) - w(d(Iy,y)), d(Iy,Iy)\} = d(Iy,y) - w(d(Iy,y)), d(Iy,Iy) = d(Iy,y) - w(Iy,Iy), d(Iy,Iy), d(Iy,Iy) = d(Iy,Y) - w(Iy,Iy), d(Iy,Iy) = d(Iy,Y) - w(Iy,Iy), d(Iy,Iy), d(Iy,Iy) = d(Iy,Y) - w(Iy,Iy), d(Iy,Iy) = d(Iy,Y) - w(Iy,Iy), d(Iy,Iy) = d(Iy,Y) - w(Iy,Iy), d(Iy,Iy), d(Iy,Iy),$$

and it follows that *Iy*=*y*. Further,

$$\delta(fy,Gx_{2n+1}) \leq \max\{d(Iy,Jx_{2n+1}),\delta(Iy,Fy),\delta(Jx_{2n+1},Gx_{2n+1})\} - \max\{d(Iy,Jx_{2n+1}),\delta(Iy,Fy),\delta(Jx_{2n+1},Gx_{2n+1})\}\}.$$

Letting n tend to infinity we get

 $\delta(Fy,y) \le \max\{d(y,y), \delta(y,Fy), \delta(y,y)\} \cdot w[\max\{d(y,y), \delta(y,Fy), \delta(y,y)\}].$

i.e. $\delta(Fy,y) \leq \delta(Fy,y) - w(\delta(Fy,y)),$

and so $Fy=\{y\}$. This means that y is in the range of F and since the range of J contains the range of F, there must exist a point y' in X such that Jy=y and so GJy=Gy.

Now,

$$\begin{split} \delta(y,Gy') &= \delta(Fy,Gy') \leq max\{d(Iy,Jy'), \delta(Iy,Fy), \delta(Jy',Gy')\} - w[max\{d(Iy,Jy'), \delta(Iy,Fy), \delta(Jy',Gy')\}] \\ &= max\{d(y,y), d(y,y), (y,Gy')\} - w[max\{d(y,y), d(y,y), \delta(y,Gy')\}] \\ &= \delta(y,Gy') - w(\delta(y,Gy')), \end{split}$$

and this gives $\delta(y, Gy')=0$, i.e. $Gy'=\{y\}$.

Next,

$$\begin{split} \delta(Jy,Gy) &\leq \delta(Jy,GJy') \\ &\leq \delta(Jy,JGy') + \delta(JGy',GJy') \\ &= \delta(Jy,JGy'), \text{ since J and G are compatible} \\ &= d(Jy,Jy) = 0. \end{split}$$

So $Gy=\{Jy\}$. Next,

$$\begin{split} \delta(y,Gy) &= \delta(Fy,Gy) \\ &\leq max\{d(Iy,Jy),\delta(Iy,Fy),\delta(Jy,Gy)\} - w[max\{d(Iy,Jy),\delta(Iy,Fy),\delta(Jy,Gy)\}] \\ &= max\{\delta(y,Gy),d(y,y),d(Jy,Jy)\} - w[max\{d(Iy,Jy),\delta(Iy,Fy),\delta(Jy,Gy)\}] \\ &= \delta(y,Gy) - w\delta(y,Gy). \end{split}$$

So, $Gy=\{y\}$. This gives that $\{Jy\}=Gy=\{y\}$.

We have therefore proved that y is a common fixed point of F,G,I and J and that $Fy=Gy=\{y\}$. This result of course holds if J is continuous instead of I.

Now suppose that F, G, I and J have a second fixed point z. Then

 $\delta(Fy, Gz) \leq \max\{d(Iy, Jz), \delta(Iy, Fy), \delta(Jz, Gz)\} \cdot w[\max\{d(Iy, Jz), \delta(Iy, Fy), \delta(Jz, Gz)\}]$

i.e. $d(y, z) = max\{d(y,z), d(y,y), d(z,z)\} - w[max\{d(y,z), d(y,y), d(z,z)\}]$

i.e. $d(y, z) \le d(y, z) - w(d(y, z)),$

and so y=z. This means that y is the unique common fixed of F,G,I and J.

This completes the proof of the theorem.

The following Corollary follows immediately. © 2011, IJMA. All Rights Reserved

¹S. De Sarkar* and ²Kalishankar Tiwary/ SOME FIXED POINT THEOREMS FOR δ-COMPATIBLE MAPPINGS IN METRIC SPACES/ IJMA- 2(12), Dec.-2011, Page: 2813-2817

Corollary: 1 Let F and G be mappings of a complete metric space (X, d) into B(X) satisfying the inequality

 $\delta(Fx, Gy) \leq \max\{d(x, y), \delta(x, Fx), \delta(y, Fy)\} - w[\max\{d(x, y), \delta(x, Fx), \delta(y, Fy)\}]$

for all x, $y \in X$. Then F and G have unique common fixed point z. Further $Fz=Gz=\{z\}$ and z is the unique common fixed point of F and G.

Theorem: 2 Let f, g, I and J be mappings of a complete metric space (X, d) into itself satisfying the inequality

 $d(fx, gy) \leq \max\{d(Ix, Jy), d(Ix, fx), d(Jy, gy)\} - w[\max\{d(Ix, Jy), d(Ix, fx), d(Jy, gy)\}]$

for all x, $y \in X$. If the mappings f and I are compatible and the mappings g and J are compatible, $g(X) \subset I(X), f(X) \subset J(X)$ and if one of the mappings I and J is continuous then, f, g, I and J have a unique fixed point z.

Proof: Define set-valued mappings *F* and *G* on *X* by $Fx = \{fx\}$ and $Gx = \{gx\}$ for all $x \in X$. If *I* (or *J*) is continuous then the conditions of Theorem 4 are satisfied and it follows that *f*,*g*,*I* and *J* have a unique common fixed point *z*.

Since commuting maps are compatible, we have the following Corollary.

Corollary: 2 Let *f*, *g* and *h* be mappings of a complete metric space into itself, *h* is continuous, fh = hf; gh = hg; $f(X) \cup g(X) \subset h(X)$. If *f*, *g* and *h* satisfy $d(fx, gy) \le maxfd(hx,hy)$, $d(hx, fx), d(hy, gy) - w[max\{d(hx, hy), d(hx, fx), d(hy, gy)]$ for all *x*, *y* in *X*, then *f*, *g* and *h* have a unique common fixed point in *X*. Corollary 2 is the theorem of Liu [7].

REFERENCES:

[1] B. Fisher, Common fixed point theorems for mappings and set-values mappings, J. Univ. Kuwait (Sci) 11 (1) (1984), 15-21.

[2] B. Fisher, Common fixed points of mappings and set-valued mappings on metric spaces, Kyungpook Math. J. 25 (1) (1985), 35-42.

[3] G. Jungck, Compatible mapping and common fixed points, Int. J. Math. Math. Sci. 9 (1986), 771-779.

[4] G. Jungck, Common fixed points for commuting and compatible mappings on compacta, Proc. Amer. Math. Soc. 130 (1988), 977-983.

[5] G. Jungck, Compatible mappings and common fixed points (2), Int. J. Math. Math. Sci. 9 (1988), 285-288.

[6] G. Jungck and B. E. Rhoades, Fixed point theorems for compatible mappings, Int. J. Math. Math. Sci. 26 (3) (1993), 417-428.

[7] Z. Liu, A note on unique common fixed point, Bull. Cal. Math. Soc. 85 (1993), 469-472.

[8] B. E. Rhoades, S. Park and K. B. Moon, On generalizations of Meir-Keeler type contraction maps, J. Math. Anal. Appl. 146 (2) (1990), 482-494.

[9] B. E. Rhoades, Kalishankar Tiwary and G. N. Singh, A common fixed point theorem for compatible mapping, Indian J. Pure Appl. Math., 26 (5) (1993), 403-409.

[10] S. Sess, B. E. Rhoades and M. S. Khan, On common fixed points of compatible mappings in metric space and Banach spaces, Int. J. Math. Math. Sci. 11 (1988), 375-392.

[11] S. L. Singh, K. S. Ha and Y. J. Cho, Coincidence and fixed points of non-linear hybrid contractions, Int. J. Math. Math. Sci. 12 (1989), 247-256.

[12] Kalishankar Tiwary and Uttam Samajpati, Some common fixed point-theorems in complete metric spaces, Jnanabha 34 (2004), 133-139.
