

HYERS-ULAM-RASSIAS STABILITY OF GENERALIZED QUADRATIC FUNCTIONAL EQUATIONS

Manoj Kumar¹ & Ashish Kumar^{2*}

Department of Mathematics, Maharshi Dayanand University, Rohtak-124001, INDIA

E-mail: manoj.ahlawat393@gmail.com, akrmisc@gmail.com

(Received on: 06-02-12; Accepted on: 27-02-12)

ABSTRACT

In this paper, we consider a generalized form of the quadratic functional equations and established the stability in the spirit of D. H. Hyers, S. M. Ulam and Th. M. Rassias, for the function $f: E_1 \rightarrow E_2$, where E_1 is a normed space and E_2 a Banach space.

Keywords: - Hyers-Ulam-Rassias stability, Generalized quadratic functional equation.

MSC 2010:- 39B05, 39B62, 39B82, 30D05.

1. INTRODUCTION

The history of the stability theory of functional equations started with a problem concerning group homomorphisms posed by S.M. Ulam [5] in 1940:

Let G_1 be a group and let G_2 be a metric group with metric $d(\cdot, \cdot)$. Given $\varepsilon > 0$, does there exists a $\delta > 0$, such that if a function $h: G_1 \rightarrow G_2$ satisfies the inequality $d(h(x, y), h(x)h(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $H: G_1 \rightarrow G_2$ with $d(h(x), H(x)) < \varepsilon$ for all $x \in G_1$?

In other words, we are looking for situations when the homomorphisms are stable. i.e. if a mapping is almost a homomorphism, then there exists a true homomorphism near it. If we turn our attention to the case of functional equations, then we can ask the question, How do the solutions of the inequality differ from those of the given functional equations?

In 1941, D. H. Hyers [2] gave the first affirmative answer to the question of S. M. Ulam [5] under the assumption that G_1 and G_2 are Banach spaces. Hyers result was generalized by T. Aoki [10] for additive mappings and by Th. M. Rassias [7] for linear mappings by considering an unbounded Cauchy difference. Th. M. Rassias [7] has provided a lot of influence in the development of what we now call Hyers – Ulam – Rassias stability of functional equations.

The quadratic function $f(x) = cx^2$ satisfies the functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y) \quad (1.1)$$

and therefore the equation is called the quadratic functional equation. The first stability result for the quadratic functional equation (1.1) was proved by J. Skof [3] for the function $f: X \rightarrow Y$ where X is a normed space and Y a Banach space. The result of F. Skof [3] is still true if the relevant domain X is replaced by an abelian group and this was dealt by P. W. Cholewa [4]. This result was further generalized by Th.M. Rassias [8], C. Borelli and G. L Forti [1].

In this paper, we prove the Hyers – Ulam – Rassias stability of the following generalized quadratic functional equation

$$F(x+my) + f(x-my) = 2[f(x) + m^2 f(y)] \quad (1.2)$$

for the mapping $f: E_1 \rightarrow E_2$, where E_1 is a normed space and E_2 a Banach space.

***Corresponding author: Ashish Kumar^{2*}, *E-mail: akrmisc@gmail.com**

2. HYERS-ULAM-RASSIAS STABILITY OF (1.2)

Theorem: 2.1 Let E_1 be a normed space, E_2 be a Banach space and $\zeta: E_1 \times E_1 \rightarrow [0, \infty)$ be a function such that

$$\lim_{n \rightarrow \infty} \frac{\zeta(m^n x, m^n y)}{m^{2n}} = 0 \quad (2.1)$$

for all $x, y \in E_1$. Suppose that a function $f: E_1 \rightarrow E_2$ with $f(0) = 0$, satisfies

$$\|f(x + my) + f(x - my) - 2[f(x) + m^2 f(y)]\| \leq \zeta(x, y) \quad (2.2)$$

for all $x, y \in E_1$. Then there exists a unique quadratic function $T: E_1 \rightarrow E_2$ which satisfies the inequality

$$\|f(y) - T(y)\| \leq \frac{1}{2m^2} \sum_{i=0}^{\infty} \frac{1}{m^{2i}} \zeta(0, m^i y) \quad (2.3)$$

for all $y \in E_1$. The function T is given by

$$T(y) = \lim_{n \rightarrow \infty} \frac{f(m^n y)}{m^{2n}} \quad (2.4)$$

for all $y \in E_1$.

Proof: Let $x = 0$ in (2.2), we get

$$\|2f(my) - 2m^2 f(y)\| \leq \zeta(0, y) \quad (2.5)$$

for all $y \in E_1$, so

$$\left\| \frac{f(my)}{m^2} - f(y) \right\| \leq \frac{1}{2m^2} \zeta(0, y) \quad (2.6)$$

for all $y \in E_1$. Replacing y by my in (2.6) and dividing by m^2 and summing the resulting inequality with (2.6), we get

$$\left\| \frac{f(m^2 y)}{m^4} - f(y) \right\| \leq \frac{1}{2m^4} \zeta(0, my) + \frac{1}{2m^2} \zeta(0, y) \leq \frac{1}{2m^2} \left[\frac{1}{m^2} \zeta(0, my) + \zeta(0, y) \right] \quad (2.7)$$

for all $y \in E_1$. Using induction on a positive integer n , we obtain that

$$\left\| \frac{f(m^n y)}{m^{2n}} - f(y) \right\| \leq \frac{1}{2m^2} \sum_{i=0}^{n-1} \frac{1}{m^{2i}} \zeta(0, m^i y) \leq \frac{1}{2m^2} \sum_{i=0}^{\infty} \frac{1}{m^{2i}} \zeta(0, m^i y) \quad (2.8)$$

for all $y \in E_1$. In order to prove the convergence of the sequence $\left\{ \frac{f(m^n y)}{m^{2n}} \right\}$, we replace y by $m^k y$ and divide inequality (2.8) by m^{2k} to find that for $n, k > 0$,

$$\begin{aligned} \left\| \frac{f(m^n \cdot m^k y)}{m^{2n+2k}} - \frac{f(m^k y)}{m^{2k}} \right\| &= \left\| \frac{f(m^{n+k} y)}{m^{2(n+k)}} - \frac{f(m^k y)}{m^{2k}} \right\| \\ &\leq \frac{1}{m^{2k}} \left\| \frac{f(m^{n+k} y)}{m^{2n}} - f(m^k y) \right\| \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2m^2} \frac{1}{m^{2k}} \sum_{i=0}^{\infty} \frac{\zeta(0, m^{i+k}y)}{m^{2i}} \\ &\leq \frac{1}{2m^2} \sum_{i=0}^{\infty} \frac{\zeta(0, m^{i+k}y)}{m^{2(i+k)}} \end{aligned} \quad (2.9)$$

Since the right hand side of the inequality (2.9) tends to zero as $k \rightarrow \infty$, the sequence $\left\{ \frac{f(m^n y)}{m^{2n}} \right\}$ is a Cauchy

sequence for all $y \in E_1$. Since E_2 is complete, the sequence $\left\{ \frac{f(m^n y)}{m^{2n}} \right\}$ converges to a fixed point $T(y) \in E_1$. So, one can define the function $T: E_1 \rightarrow E_2$ by

$$T(y) = \lim_{n \rightarrow \infty} \frac{f(m^n y)}{m^{2n}}$$

for all $y \in E_1$. To show that T satisfies the equation (1.2) replacing x and y by $m^n x$, $m^n y$ respectively in (2.2) and dividing by m^{2n} , then it follows that

$$\left\| \frac{f(m^n(x+my))}{m^{2n}} + \frac{f(m^n(x-my))}{m^{2n}} - \frac{2[f(m^n x) + m^2 f(m^n y)]}{m^{2n}} \right\| \leq \frac{\zeta(m^n x, m^n y)}{m^{2n}}$$

Taking limit as $n \rightarrow \infty$ and using (2.1), it shows that T satisfies (1.2) for all $y \in E_1$.

Now, let $T': E_1 \rightarrow E_2$ be another quadratic function satisfying (1.2) and (2.3). Then we have

$$\begin{aligned} \|T(y) - T'(y)\| &= \frac{1}{m^{2n}} \|T(m^n y) - T'(m^n y)\| \\ &\leq \frac{1}{m^{2n}} (\|T(m^n y) - f(m^n y)\| + \|T'(m^n y) - f(m^n y)\|) \\ &\leq \frac{1}{m^2} \sum_{i=0}^{\infty} \frac{\zeta(0, m^{i+n}y)}{m^{2(i+n)}} \end{aligned} \quad (2.10)$$

which tends to zero as $n \rightarrow \infty$ for all $y \in E_1$.

So, we can conclude that $T(y) = T'(y)$ for all $y \in E_1$ which shows the uniqueness of the function T . This completes the proof of theorem.

Corollary: 2.1 (Hyers Stability). Let E_1 and E_2 be a real normed space and Banach space, respectively, and let $\varepsilon \geq 0$ be a real number. Suppose that a function $f: E_1 \rightarrow E_2$ with $f(0) = 0$ satisfies

$$\|f(x+my) + f(x-my) - 2[f(x) + m^2 f(y)]\| \leq \varepsilon$$

for all $x, y \in E_1$. Then there exists a unique quadratic function $T: X \rightarrow Y$ defined by $T(y) = \lim_{n \rightarrow \infty} \frac{f(m^n y)}{m^{2n}}$ which

satisfies the equation (1.2) and the inequality $\|f(y) - T(y)\| \leq \frac{\varepsilon}{2(m^2 - 1)}$ for all $y \in E_1$. Further, if for each fixed $y \in E_1$

the mapping $t \rightarrow f(ty)$ from R to E_2 is continuous, then $T(my) = m^2 T(y)$.

Corollary: 2.2 (Rassias Stability) Let E_1 and E_2 be a real normed space and a Banach space, respectively, and let $\varepsilon \geq 0$, $0 < p < 2$ be real numbers. Suppose that a function $f: E_1 \rightarrow E_2$ with $f(0) = 0$ satisfies

$$\|f(x + my) + f(x - my) - 2[f(x) + m^2 f(y)]\| \leq \varepsilon(\|x\|^p + \|y\|^p)$$

for all $x, y \in E_1$. Then there exists a unique quadratic mapping $T: E_1 \rightarrow E_2$ which satisfies the equation (1.2) and the inequality

$$\|f(y) - T(y)\| \leq \frac{\varepsilon}{2(n^2 - m^p)} \|y\|^p$$

for all $y \in E_1$. The function T is given by

$$T(y) = \lim_{n \rightarrow \infty} \frac{f(m^n y)}{m^{2n}}$$

for all $y \in E_1$. Further, if for each fixed $y \in E_1$ the mapping $t \rightarrow f(ty)$ from R to E_2 is continuous, then $T(my) = m^2 T(y)$ for all $m \in R$.

Theorem: 2.2 Let E_1 be a normed space, E_2 be a Banach space and $\zeta: E_1 \times E_2 \rightarrow [0, \infty)$ be a function such that

$$\lim_{n \rightarrow \infty} m^n \zeta\left(\frac{x}{m^n}, \frac{y}{m^n}\right) = 0 \quad (2.11)$$

for all $x, y \in E_1$. Suppose that a function $f: E_1 \rightarrow E_2$ with $f(0) = 0$, satisfies

$$\|f(x + my) + f(x - my) - 2[f(x) + m^2 f(y)]\| \leq \zeta(x, y) \quad (2.12)$$

for all $x, y \in E_1$. Then there exists a unique quadratic function $T: E_1 \rightarrow E_2$ which satisfies the inequality

$$\|f(y) - T(y)\| \leq \frac{1}{2} \sum_{i=0}^{\infty} m^{2i} \zeta\left(0, \frac{y}{m^{i+1}}\right) \quad (2.13)$$

for all $y \in E_1$. The function T is given by

$$T(y) = \lim_{n \rightarrow \infty} m^{2n} f\left(\frac{y}{m^n}\right) \quad (2.14)$$

Proof: Replacing $y = \frac{y}{m}$ and multiplying by m^2 in (2.6) we get

$$\left\| f(y) - m^2 f\left(\frac{y}{m}\right) \right\| \leq \frac{1}{2} \zeta\left(0, \frac{y}{m}\right) \quad (2.15)$$

Again replacing y with y/m and multiplying by m^2 in (2.15).

$$\left\| m^4 f\left(\frac{y}{m^2}\right) - f(y) \right\| \leq \frac{m^2}{2} \zeta\left(0, \frac{y}{m^2}\right) + \frac{1}{2} \zeta\left(0, \frac{y}{m}\right)$$

for all $y \in E_1$. Hence

$$\|T(y) - f(y)\| \leq \frac{1}{2} \sum_{i=0}^{\infty} m^{2i} \zeta\left(0, \frac{y}{m^{i+1}}\right) \quad (2.16)$$

To prove the convergence of the sequence $\left\{ m^{2n} f\left(\frac{y}{m^n}\right) \right\}$, we replace y by $\frac{y}{m^k}$ and multiplying inequality by m^{2k} ,

we have

$$\left\| m^{2n+2k} f\left(\frac{y}{m^{n+k}}\right) - m^{2k} f\left(\frac{y}{m^k}\right) \right\| \leq \frac{1}{2} \sum_{i=0}^{\infty} m^{2(i+k)} \zeta\left(0, \frac{y}{m^{i+k}}\right)$$

It follows from (2.16) that $\left\{ m^{2n} f\left(\frac{y}{m^n}\right) \right\}$, is a Cauchy sequence for all $y \in E_1$. Since E_2 is complete the sequence

$\left\{ m^{2n} f\left(\frac{y}{m^n}\right) \right\}$ converges to a unique point $T \in E_2$. So, we can define the mapping $f: E_1 \rightarrow E_2$ by

$$T(y) = \lim_{n \rightarrow \infty} m^{2n} f\left(\frac{y}{m^n}\right)$$

for all $y \in E_1$. Also by Theorem 2.1. $T: E_1 \rightarrow E_2$ is quadratic mapping. The rest of the proof is similar to the proof of Theorem 2.1.

Corollary: 2.3 (Hyers stability) If a function $f: E_1 \rightarrow E_2$ satisfies $f(0) = 0$ and the inequality

$$\|f(x + my) + f(x - my) - 2[f(x) + m^2 f(y)]\| \leq \varepsilon$$

for all $x, y \in E_1$, then there exists a unique quadratic mapping $T: E_1 \rightarrow E_2$ such that

$$\|f(y) - T(y)\| \leq \frac{q}{2(1 - m^2)}$$

for all $y \in E_1$. The function T is given by $T(y) = \lim_{n \rightarrow \infty} m^{2n} f\left(\frac{y}{m^n}\right)$ for all $y \in E_1$.

Corollary: 2.4 (Rassias stability) If a function $f: E_1 \rightarrow E_2$ with $f(0) = 0$ satisfies the inequality

$$\|f(x + my) + f(x - my) - 2[f(x) + m^2 f(y)]\| \leq \alpha(\|x\|^p + \|y\|^p)$$

for some $p > 2$ and for all $x, y \in E_1$, then there exists a unique quadratic function $T: E_1 \rightarrow E_2$ such that

$$\|f(y) - T(y)\| \leq \frac{1}{2} \frac{\varepsilon}{(m^p - m^2)} \|y\|^p$$

for all $y \in E_1$. The function T is given by $T(y) = \lim_{n \rightarrow \infty} m^{2n} f\left(\frac{y}{m^n}\right)$ for all $y \in E_1$.

REFERENCES

- [1] C. Borelli and G.L. Forti, On a general Hyers-Ulam stability, Int. J. Math. Math. Sci. (18) (1995), pp. 229-236.
- [2] D. H. Hyers, On the Stability of the Linear Functional Equation, Proc. Nat. Acad. Sci. U.S.A. (27) (1941), pp. 222-224.
- [3] F. Skof, Proprietà locali e approssimazione di operatori, Rend. Sem. Mat. Fis. Milano, (53) (1983), pp. 113-129.
- [4] P. W. Cholewa, Remarks on the stability of functional equations, Aequationes Math., (27) (1984), pp. 76-86.

- [5] S. M. Ulam, A Collection of the Mathematical Problems, Interscience Publ. New York, 1960.
- [6] S. Czerwik, On the stability of the quadratic mappings in normed spaces, Abh. Math. Sem. Univ. Hamburg, (62) (1992), pp. 59–64.
- [7] Th. M. Rassias, On the stability of the Linear mapping in Banach spaces, Procc. Of the American Mathematical Society, (72) (2) (1978), pp. 297-300.
- [8] Th. M. Rassias, On the stability of the functional equations in Banach spaces, J. Math. Anal. Appl., (251) (2000), pp. 264–284.
- [9] T. Aoki, On the stability of the linear transformation in Banach spaces, Journal of the Mathematical Society, (2) (1-2) (1950), pp. 64-66.
