

M-CONTINUITY AND ITS DECOMPOSITIONS

¹R. G. Balamurugan, ²S. Vijaya and ³O. Ravi*

¹*Department of Mathematics, Cauvery International School, Manaparai,
Trichy District, Tamil Nadu, India
E-mail: rgbala2010@yahoo.com*

²*Department of Mathematics, Madurai Institute of Engineering and Technology,
Pottapalayam, Sivagangai District, Tamil Nadu, India
E-mail: viviphd.11@gmail.com*

³*Department of Mathematics, P. M. Thevar College, Usilampatti, Madurai District, Tamil Nadu, India
E-mail: siingam@yahoo.com*

(Received on: 04-01-12; Accepted on: 09-02-12)

ABSTRACT

The aim of this paper is to introduce the notions of R -locally m -closed sets and π -locally m -closed sets and some new subsets of minimal spaces and to obtain decompositions of M -continuity.

2010 Mathematics Subject Classification: 54C10, 54C08, 54C05.

Key words and Phrases: R -locally m -closed set, π -locally m -closed set, A_{mr} -set, $A_{m\pi}$ -set, (A, m) -closed set, (A, mr) -closed set, $(A, m\pi)$ -closed set.

1. INTRODUCTION

In [4] Maki introduced the notions of minimal structures and minimal spaces. Popa and Noiri [6] introduced a new notion of M -continuous functions as a function defined between sets satisfying some minimal conditions. In 1970, the notion of generalized closed (briefly, g -closed) sets were introduced and investigated by Levine [3]. Recently, many modifications of g -closed sets have defined and investigated. One among them is mg -closed sets which were introduced by Noiri and studied in [5]. In [5], he also introduced locally m -closed sets in minimal spaces.

In this paper, we introduce the notions of R -locally m -closed sets and π -locally m -closed sets, some new subsets of minimal spaces and obtain decompositions of M -continuity. Also we investigate some properties and characterizations of these sets with some theorems, examples and counter examples.

2. PRELIMINARIES

Definition 2.1[4]: A subfamily $m_x \subset P(X)$ is said to be a minimal structure on X if $\emptyset, X \in m_x$. The pair (X, m_x) is called a minimal space (or an m -space). A subset A of X is said to be m -open if $A \in m_x$. The complement of an m -open set is called m -closed set. We set $m\text{-Int}(A) = \bigcup \{U : U \subset A, U \in m_x\}$ and $m\text{-Cl}(A) = \bigcap \{F : A \subset F, X - F \in m_x\}$.

Lemma 2.2 [6]: Let (X, m_x) be an m -space and $A \subset X$. Then $x \in m\text{-Cl}(A)$ if and only if $U \cap A \neq \emptyset$ for every $U \in m_x$ containing x .

A minimal space (X, m_x) has the property $[B]$ if the union of any family of subsets belonging to m_x belongs to m_x .

Proposition 2.3 [6]: Let (X, m_x) be a minimal space.

- (i) For any two subsets A, B of X , the following properties hold:
(a) $A \subset m\text{-Cl}(A)$ and $A = m\text{-Cl}(A)$ if A is a m -closed set.

Corresponding author: ³O. Ravi, *E-mail: siingam@yahoo.com

- (b) $m\text{-Int}(A) \subset A$ and $A = m\text{-Int}(A)$ if A is an m -open set.
- (c) $A \subset B \Rightarrow m\text{-Cl}(A) \subset m\text{-Cl}(B)$ and $A \subset B \Rightarrow m\text{-Int}(A) \subset m\text{-Int}(B)$.
- (d) $m\text{-Cl}(m\text{-Cl}(A)) = m\text{-Cl}(A)$.
- (e) $(m\text{-Cl}(A))^c = m\text{-Int}(A^c)$ and $(m\text{-Int}(A))^c = m\text{-Cl}(A^c)$.
- (f) $m\text{-Cl}(\phi) = \phi$; $m\text{-Cl}(X) = X$; $m\text{-Int}(\phi) = \phi$; $m\text{-Int}(X) = X$.

(ii) The following are equivalent.

- (a) m_x has the property [B].
- (b) If $m\text{-Int}(A) = A$, then $A \in m_x$.
- (c) If $m\text{-Cl}(B) = B$, then $X - B \in m_x$.

Definition 2.4 [8]: A subset A of a minimal space (X, m_x) is said to be

- (a) regular m -open if $A = m\text{-Int}(m\text{-Cl}(A))$,
- (b) m -semi open if $A \subset m\text{-Cl}(m\text{-Int}(A))$,
- (c) $m\text{-}\pi$ -open if it is the finite union of regular m -open sets of A .

Definition 2.5 [5]: A subset A of a minimal space (X, m_x) is said to be m g-closed if $m\text{-Cl}(A) \subset U$ whenever $A \subset U$ and U is m -open in X .

Definition 2.6[5]: A subset A of an m -space (X, m_x) is said to be locally m -closed if $A = U \cap V$ where U is m -open and V is m -closed.

Lemma 2.7 [8]: For the subsets of a minimal space (X, m_x) satisfying property [B], every $m\text{-}\pi$ -open set is an m -open set but not conversely.

Example 2.8: Let (X, m_x) be a minimal space satisfying property [B], such that $X = \{a, b, c\}$ and $m_x = \{\phi, \{c\}, \{a, b\}, \{a, c\}, X\}$. Then $A = \{a, c\}$ is an m -open set but not an $m\text{-}\pi$ -open set.

Remark 2.9 [8]: The implication in Lemma 2.7 will not hold if m_x does not have property [B] as shown in the following Example 2.10.

Example 2.10: Let (X, m_x) be a minimal space such that $X = \{a, b, c\}$ and $m_x = \{\phi, X, \{a\}, \{b\}\}$. Then $A = \{a, b\}$ is an $m\text{-}\pi$ -open set but not an m -open set.

Lemma 2.11 [8]: For the subsets of a minimal space (X, m_x) , every regular m -open set is an $m\text{-}\pi$ -open set but not conversely.

Example 2.12: Let (X, m_x) be a minimal space such that $X = \{a, b, c\}$ and $m_x = \{\phi, \{a\}, \{b\}, X\}$. Then $A = \{a, b\}$ is an $m\text{-}\pi$ -open set but not a regular m -open set.

Remark 2.13 [1]: For the subsets of a minimal space (X, m_x) , every m -open set is m -semi open set but not conversely.

Definition 2.14 [6]: A function $f: (X, m_x) \rightarrow (Y, m_y)$ is said to be M -continuous if for each $x \in X$ and each $V \in m_y$ containing $f(x)$, there exists $U \in m_x$ containing x such that $f(U) \subset V$.

Lemma 2.15 [6]: For a function $f: (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B], the following are equivalent.

1. f is M -continuous;
2. $f^{-1}(V)$ is m_x -open for every m_y -open set V of Y ;
3. $f^{-1}(K)$ is m_x -closed for every m_y -closed set K of Y .

3. STRONGER FORMS OF LOCALLY m -CLOSED SETS

Definition 3.1: A subset A of an m -space (X, m_x) is said to be

- (a) R -locally m -closed if $A = U \cap V$ where U is regular m -open and V is m -closed,
- (b) π -locally m -closed if $A = U \cap V$ where U is $m\text{-}\pi$ -open and V is m -closed.

Definition 3.2: A subset A of an m -space (X, m_x) is said to be

- (a) m -rg-closed if $m\text{-Cl}(A) \subset U$ whenever U is regular m -open in X and $A \subset U$,
- (b) $m\text{-}\pi$ g-closed if $m\text{-Cl}(A) \subset U$ whenever U is $m\text{-}\pi$ -open in X and $A \subset U$,
- (c) $m\omega$ -closed if $m\text{-Cl}(A) \subset U$ whenever U is m -semi open in X and $A \subset U$.

Lemma 3.3 [9]: For the subsets of an m -space (X, m_x) , the following implications hold.
 m -closed $\Rightarrow m\omega$ -closed $\Rightarrow mg$ -closed.

Lemma 3.4: For the subsets of an m -space (X, m_x) satisfying property [B], we have the following implications.
 mg -closed $\Rightarrow m$ - πg -closed $\Rightarrow m$ -rg-closed.

Lemma 3.5: Let (X, m_x) be an m -space and $A \subset X$. If A is m -closed, then

- (i) A is locally m -closed set but not conversely.
- (ii) A is R -locally m -closed set but not conversely.
- (iii) A is π -locally m -closed set but not conversely.

Remark 3.6: None of the implications in Lemmas 3.3, 3.4 and 3.5 is reversible as seen in the following Examples.

Example 3.7: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{b\}, \{a, b\}, \{a, c\}\}$. Then $A = \{b, c\}$ is an $m\omega$ -closed set but not an m -closed.

Example 3.8: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{c\}\}$. Then $A = \{a\}$ is an mg -closed set but not an $m\omega$ -closed.

Example 3.9: Let (X, m_x) be an m -space satisfying property [B] such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{c\}, \{a, b\}, \{a, c\}\}$. Then $A = \{a\}$ is an m - πg -closed set but not an mg -closed.

Example 3.10: Let (X, m_x) be an m -space satisfying property [B] such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$. Then $A = \{a, b\}$ is an m -rg-closed set but not an m - πg -closed.

Example 3.11: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{a\}, \{b\}\}$. Then

- (i) $A = \{b\}$ is locally m -closed set but not a m -closed set.
- (ii) $A = \{a\}$ is both R -locally m -closed set and π -locally m -closed set but not a m -closed.

Proposition 3.12: Let (X, m_x) be an m -space and A a subset of X .

1. If A is m - π -open, then A is π -locally m -closed set.
2. If A is R -locally m -closed set, then A is π -locally m -closed set.

Proposition 3.13: Let (X, m_x) be an m -space satisfying property [B] and A a subset of X . Then the following holds.

If A is π -locally m -closed set, then A is locally m -closed set.

Remark 3.14: The converses of the above Propositions 3.12 and 3.13 need not be true as shown in the following examples.

Example 3.15: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then $A = \{a, c\}$ is π -locally m -closed set but not m - π -open.

Example 3.16: Let (X, m_x) be an m -space such that $X = \{a, b, c, d\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, X\}$. Then $A = \{a, b, c\}$ is π -locally m -closed set but it is not R -locally m -closed set.

Example 3.17: Let (X, m_x) be an m -space satisfying property [B] such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, X, \{c\}, \{a, b\}, \{a, c\}\}$. Then $A = \{a\}$ is locally m -closed set but not π -locally m -closed set.

Remark 3.18: For the subsets of an m -space (X, m_x) satisfying property [B], by Propositions 3.12 and 3.13, we have the following implications.

R -locally m -closed set $\Rightarrow \pi$ -locally m -closed set \Rightarrow locally m -closed set.

Theorem 3.19: A subset A of an m -space (X, m_x) satisfying property [B] is m -closed if and only if it is

- (i) locally m -closed and mg -closed. [4]
- (ii) R -locally m -closed and m -rg-closed.
- (iii) π -locally m -closed and m - πg -closed.

Proof: (i) Necessity is trivial. We prove only sufficiency. Let A be locally m -closed set and mg -closed set. Since A is locally m -closed, $A = U \cap V$, where U is m -open and V is m -closed. So, we have $A = U \cap V \subset U$. Since A is mg -closed, $m\text{-Cl}(A) \subset U$. Also $A = U \cap V \subset V$ and V is m -closed, then $m\text{-Cl}(A) \subset V$. Consequently, we have $m\text{-Cl}(A) \subset U \cap V = A$ and hence A is m -closed.

(ii) and (iii) It is similar to that of (i).

Theorem 3.20: For a subset A of an m -space (X, m_x) satisfying property $[B]$, the following are equivalent.

- (i) A is m -closed.
- (ii) A is R -locally m -closed and mg -closed.
- (iii) A is R -locally m -closed and $m\text{-rg}$ -closed.

Theorem 3.21: For a subset A of an m -space (X, m_x) satisfying property $[B]$, the following are equivalent.

- (i) A is m -closed.
- (ii) A is π -locally m -closed and $m\omega$ -closed.
- (iii) A is locally m -closed and mg -closed.

Theorem 3.22: For a subset A of an m -space (X, m_x) satisfying property $[B]$, the following are equivalent.

- (i) A is m -closed.
- (ii) A is locally m -closed and $m\omega$ -closed.
- (iii) A is locally m -closed and mg -closed.

Theorem 3.23: For a subset A of an m -space (X, m_x) satisfying property $[B]$, the following are equivalent.

- (i) A is m -closed.
- (ii) A is R -locally m -closed and $m\omega$ -closed.
- (iii) A is π -locally m -closed and mg -closed.
- (iv) A is π -locally m -closed and $m\text{-}\pi g$ -closed.

Theorem 3.24

For a subset A of an m -space (X, m_x) satisfying property $[B]$, the following are equivalent.

- (i) A is m -closed.
- (ii) A is R -locally m -closed and mg -closed.
- (iii) A is R -locally m -closed and $m\text{-}\pi g$ -closed.
- (iv) A is R -locally m -closed and $m\text{-rg}$ -closed.

Remark 3.25

1. The notions of locally m -closed sets and mg -closed sets (resp. $m\omega$ -closed sets) are independent.
2. The notions of π -locally m -closed sets and mg -closed sets (resp. $m\omega$ -closed sets, $m\text{-}\pi g$ -closed sets) are independent.
3. The notions of R -locally m -closed sets and mg -closed sets (resp. $m\omega$ -closed sets, $m\text{-rg}$ -closed sets, $m\text{-}\pi g$ -closed sets) are independent.

Example 3.26

- (i) Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{c\}, \{a, b\}, \{a, c\}, X\}$. Then $A = \{b, c\}$ is both mg -closed set and $m\omega$ -closed set but it is not locally m -closed set.
- (ii) Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{c\}, X\}$. Then $A = \{c\}$ is locally m -closed set but it is not mg -closed set.
- (iii) Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then $A = \{a\}$ is locally m -closed set but it is not $m\omega$ -closed.

Example 3.27

- (i) Let (X, m_x) be an m -space satisfying property $[B]$ such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{c\}, \{a, b\}, \{a, c\}, X\}$. Then $A = \{b, c\}$ is both mg -closed set and $m\omega$ -closed set but it is neither R -locally m -closed set nor π -locally m -closed set. Moreover it is both $m\text{-rg}$ -closed set and $m\text{-}\pi g$ -closed set.

(ii) Let (X, m_x) be an m -space satisfying property [B] such that $X = \{a, b, c, d\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, X\}$. Then $A = \{a\}$ is both R -locally m -closed set and π -locally m -closed set but it is neither mg -closed set nor $m\omega$ -closed set. Moreover it is neither m - rg -closed set nor m - πg -closed set.

4. ON NEW SUBSETS OF MINIMAL SPACES

Definition 4.1 [2]: Let A be a subset of a minimal space (X, m_x) . Then the m -kernel of the set A , is denoted by $\Lambda_m(A)$, is the intersection of all m -open supersets of A .

Definition 4.2[2]: A subset A of a minimal space (X, m_x) is called Λ_m -set if $A = \Lambda_m(A)$.

Definition 4.3 [2]: A subset A of an m -space (X, m_x) is called (Λ, m) -closed if $A = U \cap V$ where U is Λ_m -set and V is m -closed.

Lemma 4.4:

- (i) Every locally m -closed set is (Λ, m) -closed.
- (ii) Every m -closed set is (Λ, m) -closed but not conversely.[2]

Example 4.5: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then $A = \{a\}$ is (Λ, m) -closed set but not m -closed.

Lemma 4.6 [2]: For a subset A of an m -space (X, m_x) satisfying property [B], the following conditions are equivalent.

- (i) A is (Λ, m) -closed.
- (ii) $A = L \cap m\text{-Cl}(A)$ where L is Λ_m -set.
- (iii) $A = \Lambda_m(A) \cap m\text{-Cl}(A)$.

Lemma 4.7: A subset $A \subset (X, m_x)$ is mg -closed if and only if $m\text{-Cl}(A) \subset \Lambda_m(A)$.

Proof: Suppose that $A \subset X$ is mg -closed set. Let $x \in m\text{-Cl}(A)$. Suppose $x \notin \Lambda_m(A)$. Then there exists an m -open set U containing A such that $x \notin U$. Since A is mg -closed set, $A \subset U$ and U is m -open implies that $m\text{-Cl}(A) \subset U$ and so $x \notin m\text{-Cl}(A)$, a contradiction. Therefore $m\text{-Cl}(A) \subset \Lambda_m(A)$. Conversely, suppose $m\text{-Cl}(A) \subset \Lambda_m(A) \subset U$. Therefore A is mg -closed.

Theorem 4.8: For a subset A of an m -space (X, m_x) satisfying property [B], the following conditions are equivalent.

- (i) A is m -closed.
- (ii) A is mg -closed and locally m -closed.
- (iii) A is mg -closed and (Λ, m) -closed.

Proof: (i) \Rightarrow (ii) \Rightarrow (iii) Obvious.

(iii) \Rightarrow (i) Since A is mg -closed, so by Lemma 4.7, $m\text{-Cl}(A) \subset \Lambda_m(A)$. Since A is (Λ, m) -closed, so by Lemma 4.6,

$A = \Lambda_m(A) \cap m\text{-Cl}(A) = m\text{-Cl}(A)$. Hence A is m -closed.

The following two examples show that the concepts of mg -closed sets and (Λ, m) -closed sets are independent.

Example 4.9: Let (X, m_x) be an m -space such that $X = \{a, b, c, d\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, X\}$. Then $A = \{a, c\}$ is (Λ, m) -closed set but not mg -closed.

Example 4.10: Let (X, m_x) be an m -space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{b\}, \{a, b\}, \{a, c\}, X\}$. Then $A = \{b, c\}$ is mg -closed set but not (Λ, m) -closed.

Definition 4.11: Let A be a subset of a minimal space (X, m_x) . Then

- (i) The m - r -kernel of the set A , denoted by $m\text{-r-ker}(A)$, is the intersection of all regular m -open supersets of A .
- (ii) The m - π -kernel of the set A , denoted by $m\text{-}\pi\text{-ker}(A)$, is the intersection of all m - π -open supersets of A .

Definition 4.12: A subset A of a minimal space (X, m_x) is called

- (i) Λ_{mr} -set if $A = m\text{-r-ker}(A)$.
- (ii) $\Lambda_{m\pi}$ -set if $A = m\text{-}\pi\text{-ker}(A)$.

Definition 4.13: A subset A of an m -space (X, m_x) is called

- (i) (Λ, mr) -closed if $A = L \cap F$ where L is Λ_{mr} -set and F is m -closed.
- (ii) $(\Lambda, m\pi)$ -closed if $A = L \cap F$ where L is $\Lambda_{m\pi}$ -set and F is m -closed.

Lemma 4.14: Every m-closed set is $(\Lambda, m\pi)$ -closed but not conversely.

- (i) Every π -locally m-closed set is $(\Lambda, m\pi)$ -closed.
- (ii) Every m-closed set is $(\Lambda, m\pi)$ -closed but not conversely.
- (iii) Every R-locally m-closed set is $(\Lambda, m\pi)$ -closed.

Example 4.15: Let (X, m_x) be an m-space such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then

- 1. $A = \{a\}$ is $(\Lambda, m\pi)$ -closed set but not m-closed.
- 2. $A = \{a\}$ is $(\Lambda, m\pi)$ -closed set but not m-closed.

Lemma 4.16: For a subset A of an m-space (X, m_x) satisfying property [B], the following are equivalent.

- (a) 1. A is $(\Lambda, m\pi)$ -closed .
- 2. $A = L \cap m\text{-Cl}(A)$ where L is $\Lambda_{m\pi}$ -set.
- 3. $A = m\text{-r-ker}(A) \cap m\text{-Cl}(A)$.
- (b) 1. A is $(\Lambda, m\pi)$ -closed.
- 2. $A = L \cap m\text{-Cl}(A)$ where L is $\Lambda_{m\pi}$ -set.
- 3. $A = m\text{-}\pi\text{-ker}(A) \cap m\text{-Cl}(A)$.

Lemma 4.17

- (i) A subset $A \subset (X, m_x)$ is m- π g-closed if and only if $m\text{-Cl}(A) \subset m\text{-}\pi\text{-ker}(A)$.
- (ii) A subset $A \subset (X, m_x)$ is m-rg-closed if and only if $m\text{-Cl}(A) \subset m\text{-r-ker}(A)$.

Theorem 4.18: For a subset A of an m-space (X, m_x) satisfying property [B], the following are equivalent.

- (a) 1. A is m-closed.
- 2. A is m- π -closed and π -locally m-closed.
- 3. A is m- π g-closed and $(\Lambda, m\pi)$ -closed.
- (b) 1. A is m-closed.
- 2. A is m-rg-closed and R-locally m-closed.
- 3. A is m-rg-closed and $(\Lambda, m\pi)$ -closed.

Remark 4.19: By Examples 4.20 and 4.21, we realize that the following concepts are independent.

- 1. $(\Lambda, m\pi)$ -closed sets and m- π g-closed sets.
- 2. $(\Lambda, m\pi)$ -closed sets and m-rg-closed sets.

Example 4.20: Let (X, m_x) be an m-space satisfying property [B], such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then

- (i) $A = \{a\}$ is $(\Lambda, m\pi)$ -closed but not m- π g-closed.
- (ii) $A = \{c\}$ is m- π g-closed but not $(\Lambda, m\pi)$ -closed.

Example 4.21

- (i) Let (X, m_x) be an m-space satisfying property [B], such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{b\}, \{a, b\}, \{a, c\}, X\}$. Then $A = \{b, c\}$ is m-rg-closed but not $(\Lambda, m\pi)$ -closed.
- (ii) Let (X, m_x) be an m-space satisfying property [B], such that $X = \{a, b, c\}$ and $m_x = \{\emptyset, \{a\}, \{b\}, X\}$. Then $A = \{a\}$ is $(\Lambda, m\pi)$ -closed but not m-rg-closed.

5. DECOMPOSITIONS OF M-CONTINUITY

Definition 5.1: A function $f : (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B] is said to be M-g-continuous (resp. M-rg-continuous, M- ω -continuous, M- π g-continuous) if $f^{-1}(A)$ is mg-closed (resp. m-rg-closed, m ω -closed, m- π g-closed) in (X, m_x) for every m-closed set A of (Y, m_y) .

Definition 5.2: A function $f : (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B] is called

- (i) locally M-continuous if $f^{-1}(A)$ is locally m-closed in (X, m_x) for every m-closed set A of (Y, m_y) .
- (ii) R-locally M-continuous if $f^{-1}(A)$ is R-locally m-closed in (X, m_x) for every m-closed set A of (Y, m_y) .
- (iii) π -locally M-continuous if $f^{-1}(A)$ is π -locally m-closed in (X, m_x) for every m-closed set A of (Y, m_y) .

Theorem 5.3: A function $f : (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B] is M-continuous if and only if it is

- (i) locally M-continuous and M-g-continuous.
- (ii) R-locally M-continuous and M-rg-continuous
- (iii) π -locally M-continuous and M- π g-continuous.

Proof: It is an immediate consequence of Theorem 3.19.

Theorem 5.4: Let (X, m_x) be an m-space satisfying property [B]. For a function $f: (X, m_x) \rightarrow (Y, m_y)$, the following are equivalent.

- (1) f is M-continuous.
- (2) f is R-locally M-continuous and M-g-continuous.
- (3) f is R-locally M-continuous and M-rg-continuous.

Proof: It is an immediate consequence of Theorem 3.20.

Theorem 5.5: Let (X, m_x) be an m-space satisfying property [B]. For a function $f: (X, m_x) \rightarrow (Y, m_y)$, the following are equivalent.

- (1) f is M-continuous.
- (2) f is π -locally M-continuous and M- ω -continuous.
- (3) f is locally M-continuous and M-g-continuous.

Proof: It is an immediate consequence of Theorem 3.21.

Theorem 5.6: For a function $f: (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B], the following are equivalent.

- (1) f is M-continuous.
- (2) f is locally M-continuous and M- ω -continuous.
- (3) f is locally M-continuous and M-g-continuous.

Proof: It is an immediate consequence of Theorem 3.22.

Theorem 5.7

Let (X, m_x) be an m-space satisfying property [B]. For a function $f: (X, m_x) \rightarrow (Y, m_y)$, the following are equivalent.

- (1) f is M-continuous.
- (2) f is R-locally M-continuous and M- ω -continuous.
- (3) f is π -locally M-continuous and M-g-continuous.
- (4) f is π -locally M-continuous and M- πg -continuous.

Proof: It is an immediate consequence of Theorem 3.23.

Theorem 5.8: Let (X, m_x) be an m-space satisfying property [B]. For a function $f: (X, m_x) \rightarrow (Y, m_y)$, the following are equivalent.

- (1) f is M-continuous.
- (2) f is R-locally M-continuous and M-g-continuous.
- (3) f is R-locally M-continuous and M- πg -continuous.
- (4) f is R-locally M-continuous and M-rg-continuous.

Proof: It is an immediate consequence of Theorem 3.24.

Definition 5.9: A function $f: (X, m_x) \rightarrow (Y, m_y)$ where m_x satisfies property [B] is said to be (Λ, M) -continuous (resp. $(\Lambda, M\pi)$ -continuous, (Λ, Mr) -continuous) if $f^{-1}(A)$ is (Λ, m) -closed (resp. $(\Lambda, m\pi)$ -closed, (Λ, mr) -closed) in (X, m_x) for every m-closed set A of (Y, m_y) .

Theorem 5.10: For a function $f: (X, m_x) \rightarrow (Y, m_y)$, satisfying property [B], the following are equivalent.

- (1) f is M-continuous.
- (2) f is M-g-continuous and locally M-continuous.
- (3) f is M-g-continuous and (Λ, M) -continuous.

Proof: It is an immediate consequence of Theorem 4.8.

Theorem 5.11: For a function $f: (X, m_x) \rightarrow (Y, m_y)$ satisfying property [B], the following are equivalent.

- (a) 1. f is M-continuous.
2. f is M- πg -continuous and π -locally M-continuous.
3. f is M- πg -continuous and $(\Lambda, M\pi)$ -continuous.
- (b) 1. f is M-continuous.
2. f is M-rg-continuous and R-locally M-continuous.
3. f is M-rg-continuous and (Λ, Mr) -continuous.

Proof: It is an immediate consequence of Theorem 4.18.

REFERENCES

- [1] J. Antony Rex Rodrigo, O. Ravi, R. G. Balamurugan and K. Vijayalakshmi, A unified theory for modifications of \hat{g} -closed sets, International Journal of Advances in Pure and Applied Mathematics, 1 (1) (2011), 53-67.
- [2] F. Cammaroto and T. Noiri, On Λ m-sets and related topological spaces, Acta Math. Hungar., 109(3)(2005), 261-279.
- [3] N. Levine, Generalized closed sets in topology, Rend.Circ.Mat.Palermo(2), 19(1970), 89-96.
- [4] H. Maki, On generalizing semi-open and preopen sets, Report for Meeting on Topological Spaces Theory and its Applications, August 1996, Yatsushiro College of Technology, 13-18.
- [5] T. Noiri, A unified theory for modifications of g-closed sets, Rendiconti del circolo matematico di Palermo, Serie II, Tomo LVI (2007), 171-184.
- [6] V. Popa and T. Noiri, On M-continuous functions, Anal. Univ. "Dunarea de Jos" Galati, Ser. Mat. Fiz. Mec. Teor., (2), 18 (23) (2000), 31-41.
- [7] O. Ravi, R. Senthilkumar, A. R. Thilagavathi and K. Indirani, $m-I_{\pi g}$ -closed sets and $M-I_{\pi g}$ -continuity, submitted.
- [8] E. Rosas, N. Rajesh and C. Carpintero, Some new types of open and closed sets in Minimal Structures – I^1 , International Mathematical Forum, 44 (4) (2009), 2169 – 2184.
